



**Institutionen för skogens produkter och marknader
och Skogsmästarskolan i Skinnskatterberg**

**Natural and cultural continuous cover forests
in Sweden**

– how much remain and how are they managed?

Robert Axelsson



**Institutionen för skogens produkter och marknader
och Skogsmästarskolan i Skinnskatterberg**

**Natural and cultural continuous cover forests
in Sweden**

– how much remain and how are they managed?

Robert Axelsson

*Examensarbete 20 poäng, D-nivå i ämnet biologi
Robert Axelsson*

*Handledare SLU: Per Angelstam
Handledare Skogsstyrelsen: Johan Svensson*

Abstract

Swedish forestry has a long and successful history of developing sustained production of wood. Consequently silviculture is dominated by clear-felling approaches, despite the fact that Swedish forests are ecologically and culturally diverse. A considerable proportion of our naturally dynamic forests were Continuous Cover Forests (CCF). CCFs were also a part of the pre-industrial cultural landscape. Both the natural and the cultural landscape CCFs are today threatened and their area extent does not satisfy the new dimensions of sustainable forest management as biodiversity conservation and socio-cultural values.

In this study I estimate the past and present amount of two natural and one cultural CCF types in two study areas using several different approaches at two different spatial scales. The results are compared with the silvicultural practices in the two areas located around two important biophysical and socio-cultural transition zones in northwest and south-central Sweden.

My results show that in our present landscape 9 to 10% of our forested land is potential wet and dry CCF sites. In addition to this there is a portion of high altitude CCF with moist oceanic climate in the Scandinavian Mountains. However, of all potential CCF sites in the southern study area only 0 to 6% remains. In the northern study area the proportion left was slightly higher at 0 to 17% of the potential, except for the high altitude CCF where the decline was less. It is clear that this difference between the study areas is due to the later arrival of the timber frontier in the north. Present management practices on different site types indicates that the situation will be the same as in the south in a few decades since Swedish forest management exercises limited consideration in different site types to maintain CCFs. To match the natural diversity of forests and to satisfy all three dimensions of sustainable forest management there is a clear need for developing alternative forest management methods in Sweden.

Sammanfattning

Den svenska skogsnäringen har sen länge utvecklat en uthållig timmer produktion. Trakthyggesbruk är den dominerande skogsbruksmetoden trots att svenska skogar har en stor ekologisk och kulturell mångfald. En stor del av den svenska skogen bestod naturligt av olika skogstyper med kontinuerlig krontäckning, dvs. skogstyper som sällan drabbades av storskaliga störningar som tex. brand. Dessa skogar som naturligt blev gamla var också en del av det förindustriella landskapet. Både den naturliga och kulturellt betingade skogen med kontinuerligt krontäcke är idag hotad och den kvarvarande arean motsvarar inte vad som skulle kunna kallas hållbart.

Jag har uppskattat den naturliga och nuvarande arealen av två naturliga och en kulturellt betingad skogstyp med kontinuerlig krontäckning. Detta har gjorts med flera olika metoder i två studieområden med flera datakällor i två olika rumsliga skalor. Resultaten har jämförts med skogsbruksmetoderna i de två studieområdena som är valda i två viktiga biogeografiska och socio-kulturella övergångszoner i nordvästra och södra sverige.

Mina resultat visar att dagens landskap innehåller 9 till 10 % fuktiga-våta och torra ståndorter passande för skog av typ med kontinuerlig krontäckning. Dessutom finns en andel mark på höga höjder i fjällkedjan med denna karaktär på grund av det oceana klimatet. Av all potentiell mark med skog av kontinuerlig krontäckningstyp så är bara 0-6% gammal. I norr så är andelen något högre med 0-17% av potentialen. På höga höjder i fjällkedjan så återstår en större andel. Data visar klart att detta beror på den senare ankomsten av timmerfronten i norr. Nuvarande skötsel på olika ståndorter indikerar att situationen i norr kommer att vara densamma som i söder om några decennium. Detta eftersom skogsbruket inte tar tillräcklig hänsyn och anpassar brukandet till ståndorten i tillräckligt stor utsträckning för att behålla dessa skogstyper. För att matcha mångfalden i de svenska skogarna så finns det helt klart ett behov av att utveckla alternativa skogsskötselmetoder i Sverige.

Key words

continuous cover forest, continuous cover forestry, forest dynamics, forest management

Contents

<i>Introduction</i>	1
<i>Study areas</i>	5
Landscape scale analyses.....	5
Stand scale analyses	7
<i>Methods</i>	8
Landscape scale analyses.....	8
Sample plot analyses.....	8
Coarse-grained spatially explicit analyses	8
Cultural landscapes.....	9
Stand scale analyses	9
<i>Results</i>	10
Landscape scale.....	10
Cultural landscape	13
Stand scale	13
<i>Discussion</i>	15
<i>Continuous cover forests are rare</i>	15
Remote sensing as a tool to estimate and identify remaining continuous cover forests	16
Gap between policy, forest management and continuous cover forests	17
Conservation of continuous cover forests	18
A need to improve the thematic resolution of forest management plans	19
<i>Conclusion</i>	20
The way forward	20
The future for our continuous cover forests and forestry	21
<i>References</i>	22

Introduction

In most developed countries the historical use of forests has evolved through several more or less distinct phases (Björklund, 1984; Angelstam and Arnold, 1993; Drushka, 2003; Williams, 2003; Burton et al., 2003). In general three historical phases can be identified, (Östlund and Zackrisson, 2000): (1) The pre-industrial agricultural phase with a low intensity use of the forests, often including animal husbandry with cattle grazing the forests. (2) The “diverse uses” phase where the usage of a range of different resources was diversified and intensified. This phase included local and regional small-scale production of for example potash, tar and charcoal together with extraction of larger trees. (3) The industrial phase, where the use of the forest was even further intensified and often less diverse. This phase initially often resulted in unsustainable exploitation of the wood resources. The realisation that harvesting rates were larger than the regeneration and growth rates in the forest was an important driver towards the development of a sustainable timber yield (Schuler, 1998; Drushka, 2003; Kardell, 2004).

As forestry in Sweden developed after WW2, there was a need for improved and more economic efficient management regime. Large forest tracts were in poor condition due ineffective regeneration and poor forest management during the exploitative phase. In the 1948 forestry act the importance of industrial forestry for the country was emphasised, and measures of existing knowledge on how to optimise production levels to improve and optimise production to ensure a long-term sustainable yield timber and pulpwood were taken. Clear-felling is an economically efficient method that is quite simple to practice using mechanised methods (Matthews, 1989). As a consequence this method became the main silvicultural approach in Sweden since about 1950 (Ekelund and Hamilton, 2001). An important part of the development of economic sustainable forestry has been the implementation of effective means of regeneration of forest after harvesting.

Around 1970 a discussion of ecological issues began in society, which also affected the forest sector. In Sweden, clear-felling methods (Anon., 1974) triggered increased environmental considerations (Anon., 2002b; Angelstam, 2003a). In 1983 the United Nations (UN) appointed the so called Brundtland commission to propose strategies on

how to improve human well-being in the short term without threatening the local and global environment in the long term (WCED, 1987). This report triggered several forest-related events such as the development of the Forest principles and the UNFF (UN, 1992, 2000).

In Sweden's forest policy from 1993 it is clearly stated that economic and environmental-ecological aspects of forestry are of equal importance (Anon., 2001a, 2002a). The environmental quality goals of Sweden have reinforced this (Anon., 2001b). Finally, during the last few years, also socio-cultural values have been acknowledged (Anon., 2002a). This is a trend also in the rest of Europe (Malcolm et al., 2001; Mason, 2003). Thus, all three pillars of the concept sustainable development are now included in the national policies related to the management of forests in Sweden. The development of the concept sustainable forest management (SFM) has thus entered a fourth phase representing post-modern values (Bolshakov, 2000).

As a consequence, the selection and combination of silvicultural methods and landscape planning is widely discussed (e.g., Angelstam and Pettersson, 1997; Fries et al., 1997, 1998; Raivio et al., 2001). Forestry has since the 19th century reduced many elements and structures as well as simplified the dynamics of natural Swedish forests to a minimum (Östlund et al., 1997). An evaluation of the contemporary Swedish forest policy was established in 1993, and in the subsequent 1994 Forestry Act it was concluded that there is a need for developing alternatives to the clear-felling approaches (Anon., 2002b), such as continuous cover forestry. Initially the focus was on emulating natural disturbance regimes for the maintenance of biodiversity (Rülcker et al., 1994; Angelstam, 1998). Alternative forest management approaches has also been advocated to satisfy socio-cultural values such as recreation and tourism both in Sweden (Lindhagen, 1996; Rydberg and Falck, 2000; Rydberg, 2001) and internationally (Konijnendijk, 1997, 1999, 2003; Wenner, 2000). Management recommendations related to climate change, storm and other new risks include similar conclusions (Anon., 1999; Blennow, 2004; Chao et al., 2005).

From an ecological perspective, scientists agree that forests are formed by different kinds of disturbance dynamics (Sprugel, 1991; Kuuluvainen, 1994, 2002; Angelstam, 1998; Bergeron et al., 2002; Korpilahti and Kuuluvainen, 2002; Angelstam and

Kuuluvainen, 2004). This means that, depending on local and regional factors, a natural forest landscape may contain several types of dynamic systems (Angelstam, 1998). The result is a very diverse forest landscape when it comes to tree age distribution, structures and species (Kuuluvainen et al., 1998, 2002; Pennanen, 2002). Following the disturbance regime paradigm Swedish boreal and temperate natural forests can be divided into three main groups (for details see Falinski, 1986; Rülcker et al., 1994; Fries et al., 1997; Angelstam and Kuuluvainen, 2004): (1) Succession dynamics on mesic sites with large-scale disturbance regimes, resulting in largely even-aged stands with successions of young to old-growth deciduous, coniferous and mixed stands. (2) Multi-cohort dynamics on dry and poor sites where low-intensity fires often kill younger trees and leave some older and larger trees. Typically dominated by Scots pine *Pinus silvestris* or Oak *Quercus robur* stands contain trees of several age classes. (3) Gap phase dynamics with small-scale disturbances, but as a rule not fire, on wet sites and in humid climates (Angelstam and Kuuluvainen, 2004). Typically Norway spruce *Picea abies* and other shade tolerant species dominate. Here some trees die of age, disease or because they are weak and can not stand strong winds or other kinds of biotic and abiotic stress. Eventually a few trees fall and create small gaps. In a naturally dynamic landscape continuous cover forests dominated in multi-cohort and gap phase dynamics, and occurred in parts of landscape's mesic sites by chance not subject to stand-replacing disturbances.

In analogy with natural forest disturbances, human use of the landscape often gave rise to cultural continuous cover forests when tree covered grasslands were either mowed or used as grazing grounds for domestic cattle. In addition, people collecting winter fodder used trees from a range of species in the cultural landscape. This often resulted in large and old slow growing deciduous trees (Nolbrant, 1998; Ekstam and Forshed, 1996). During the 20th century, with forest management practices including cleaning of deciduous trees, the cultural continuous forests provided rescue habitats for many species and have thus been vital for their survival (Tucker and Evans, 1997; Kirby and Watkins, 1998; Mikusinski et al., 2003).

The evolution of the sustainable forest management concept implies a need to evaluate the extent to which the silvicultural methods used match the different management objectives including forest production, biodiversity conservation and

socio-cultural functions (Angelstam, 2003b). In Sweden today about 96% of forests are managed by clear-felling methods (variable retention and seed tree) (Anon., 2002b). By contrast, estimates of the amount of forest land hosting continuous cover forests in boreal forest range from 10% (Anon., 2005c) to 30% (Rülcker et al., 1994) and as high as more than 50% (Pennanen, 2002). In addition a proportion of natural succession forests also had continuous cover characteristics (Pennanen, 2002). The lower end of this range refers to today's situation with anthropogenic landscape alterations due to for example forest drainage, accumulation of nutrients and non-natural forest dynamics, while upper end is an estimate of the natural situation. In addition a portion of especially southern Sweden's forests consisted of cultural continuous cover forests in the form of wooded grasslands. Finally, alternatives to clear-felling are desired to satisfy recreational needs (Konijnendijk, 1997, 1999, 2003; Lindhagen, 1996; Rydberg and Falck, 2000; Wenner, 2000; Rydberg, 2001). In countries of economic transition also livelihood issues require alternative to clear-felling (Antrop, 2005).

There is thus overwhelming evidence that the current low diversity in silvicultural approaches is unlikely to match the current ecological and socio-cultural dimensions of SFM. Countries having developed the current broad view on the SFM concept earlier than Sweden have also initiated the development of new silvicultural approaches. To mitigate problems with natural hazards countries in the Alps developed continuous cover methods already several hundred years ago (Dengler, 1944, 1990). In much of Western Europe such methods are advocated for forests dominated by shade-tolerant trees. Similarly, in Great Britain, recent forest policy statements envisage a much greater use of CCF methods (Mason, 2003).

The aim of this paper is (a) to improve the understanding of the extent to which there is a mismatch between the natural and cultural dynamics of forests on the one hand, and today's management practices on the other; (b) to estimate the remaining amount of continuous cover forest in two study areas. I estimate the potential amount of cultural and natural continuous cover forest land at different spatial scales using different methods. Four kinds of continuous cover forests, namely multi-cohort, gap phase due to ground wetness or due to high altitude humid oceanic climate and cultural wooded grasslands are studied (see also Anon., 2004). To cover the range of

boreal forest conditions in Sweden I focus on two regions located in the major transition zones in northern and south-central Sweden, “the cultivation line” (Swedish: odlingsgränsen) the northern and high altitude border for cultivation of land (Hahn, 1998; Lundmark, pers. com.) and ”Limes Norrlandicus” (Swedish: norrlandsgränsen) (Selander, 1955; Fransson, 1965) the natural south border of the boreal forest biome (Sjörs, 1956; Aldentun, 1997) in Sweden. Landscapes located in these regions represent a wide range of ecological properties, history of forest use, and types of governance found in the boreal forest region. At this landscape scale both sampling plot data and spatially explicit coarse-grained analyses were used. Secondly, I evaluate the extent to which continuous cover stand level characteristics are found at present on sites of different types and on land managed by different forest owner categories in these two regions. Finally, trends, potential gaps between ecological and socio-cultural on the one hand, and management dimensions of continuous cover forests on the other, are discussed.

Study areas

Landscape scale analyses

There are large regional differences in the history of development of silvicultural systems (Eliasson, 2002; Angelstam, 2003b; Kardell, 2004; Holmberg, 2005). This means that different forest history phases can often be found at the same time in different regions (Angelstam et al., 1997). As illustrated by my two study areas. Sweden is a good example of this.

The southern study area (Figure 1) in Bergslagen located around the “Limes Norrlandicus” (Selander, 1955; Fransson, 1965) include both upland areas with a long land use history and lowland areas where considerable areas have been cleared for agricultural purposes. In the Bergslagen region in south-central Sweden forestry started several hundred years ago, driven by the mining industry (Wieslander, 1936). During the 17th and 18th centuries, large parts of the world’s iron production originated from Bergslagen. Due to intensive produce charcoal for the iron industry the forest rotation time was short and stands were often harvested at less than 50 years of age (Ek, 1995). When the iron industry declined in the late 19th century the practice of clear-felling with high costs for regeneration was questioned (Wallmo, 1897). After WW1 the criticism grow stronger and resulted in a period of single and group tree

selection before the breakthrough of mechanised forestry with clear-felling as the main harvesting method in the 1950s (Ekelund and Hamilton, 2001). The long history of development with industry, forestry and farming in Bergslagen resulted also in a cultural landscape with continuous cover forests. Land owners in Bergslagen range from non-industrial private to forest commons, the Swedish church and large companies. The ownership pattern in Bergslagen is fragmented with about 70% owned by owners with properties less than 1000 hectares in size. Only a few percent of the land is owned by the government.

The northern study area (Figure 2) was located in western Västerbotten and southwest Norrbotten in the northern part of Sweden around the “Cultivation Line” (Hahn, 1998; Lundmark, pers. com.). The area covers a gradient from north boreal to sub-alpine mountain forest. In this area the history of industrial forestry is relatively short. In Västerbotten county the main force affecting forests and forestry was the arrival of the timber frontier in the late 19th century associated with the export of wood to markets abroad (Angelstam et al., 2004b). In fact, large parts of the forests close to the mountains have still not been subject to intensive forest management (Forsell and Axelsson, 1990; see Table 2). Landowner patterns are different in this study area with less fragmentation and only about 25% owned by owners with properties less than 1000 hectares in size. In this study area land is to a large extent owned by the government and large forest companies. Included in this study area is the Vilhelmina Model Forest (Svensson et al., 2004).




Figure 1. The figure shows the two study areas on a map of Sweden. Bergslagen is located in mid-central Sweden and Västerbotten in northern Sweden. The dark grey squares are the 50x100 km squares used for the mixed spatially explicit methods analysis.

Cultural landscapes

The county of Västmanland is a central part of the southern study area in Bergslagen and was thus chosen for estimating the present amount of cultural landscape continuous cover forest. These kinds of culturally created wooded grasslands were never a characteristic feature in western Västerbotten county.

Stand scale analyses

For the stand scale analysis of the extent to which forest stand characteristics differ among site types (dry, mesic, moist/wet) a sample of local landscapes represented by stands or properties with the full range of site types was collected. Additionally, for the northern study area, where gap phase dynamics may be due to both ground wetness and humid oceanic climate, estates with oceanic climate were included.

Methods

Landscape scale analyses

Sample plot analyses

All sample plots from a 100-km wide band centred on the southernmost 350 km of the cultivation line and the Limes Norrlandicus (Figure 1) were extracted from the national forest inventory database (Anon., 2002c). In the northern study area it was assumed that continuous cover forests should be present at higher altitudes due to the moist oceanic climate. Data on precipitation was used to extract sample plots in this climatic zone (Ångström, 1958). In this analysis only meteorological data on precipitation was used to identify the wet sites. The age distributions of the forest on different potential continuous cover sites and mesic sites were compared. An estimate of the present amount of continuous cover forest was made by using the oldest age class i.e. forest stand 140 years old or older.

Coarse-grained spatially explicit analyses

Within the study areas two squares of 50x100 km were chosen for a coarse-grained spatially explicit analysis. A topographic wetness index (TWI) (Beven and Kirby, 1979; Rodhe and Seibert, 1999; Zinko, 2004; Güntner et al., 1999; Zinko et al., 2005) was used to estimate the area proportion of naturally moist and wet areas (Figure 2). Soil moisture is a result of several factors such as precipitation, evaporation, topography, lateral flow, soil types and vegetation (Moore et al., 1991; Wilson et al., 2004). In a limited area like the two 50x100 km squares precipitation and evaporation is assumed to be spatially uniform over a longer time scale. The model used was based on the wetness index developed by Beven and Kirby (1979). The TWI has been further developed and verified towards identification of wetter areas. Results from the TWI modelling are typically digital thematic maps with land classified in different wetness indices ranging from wet to dry. From the distribution of wetness classes I chose an amount of moist and wet areas, and combined them with digital map data on rich soil types (sediment type soils) to obtain an estimate of the potential areas for wet-rich continuous cover forests (Figure 2). The estimates were conservative to make sure I did not overestimate the proportion of this forest type. To identify the dry and poor sites only soil type information (sand-gravel and rocks with absent or very thin soil layer) were used. In the northern study area data on yearly average precipitation

was used to identify areas with high altitude humid oceanic climate. All remaining forest land was classified as being mesic sites. Finally, forest age distributions on the different site types were estimated using nearest neighbour (kNN) classified satellite image data (Tomppo, 1991; Franco-Lopez et al., 2001; McRoberts et al. 2002, Reese et al. 2003, 2005).

Cultural landscapes

The present amount of the cultural type of continuous cover forest was extracted as all cultural grasslands with a tree cover of 10% (Anon., 2005b) or more from a recent Swedish national inventory (Anon., 2005a).

Stand scale analyses

For stand scale analysis data was extracted from forest management plans (Swedish: skogsbruksplan, grön skogsbruksplan). I contacted personally and requested data from all large forest owners and some private forest owners in each study area. I asked the forest companies for data from 5 test areas containing a large sample of each continuous cover forest type. The private forest owners were identified by combining soil type and property maps at the National Board of Forestry or by just getting a list of forest owners from their staff working in the field. All private forest owners were asked if they had a forest management plan and if they were willing to let us use the data in it for my study. Often forest owners also advised us where I could find other private properties with these forest types. The collected data sets were very diverse both in size and quality in terms of variables allowing site type classification. The diversity in variables forced us to use different approaches to identify the site types having a potential to host continuous cover forests. Where ground moisture or ground layer vegetation was present in the data sets this was used. In other cases combinations of ground carriage capacity and site type index (SI), in some cases only the SI was used and in a few cases the forest owner provided a list of the stands with the searched characteristics (Anon., 1985a, 1985b, 1985c; Hägglund and Lundmark, 1987, 1999, 2003). Large parts of the collected datasets were not used since the variables presented were not sufficient to identify the site types.

Results

Landscape scale

The owner categories in the two study areas were diverse (Table 1). There was a clear difference in amount land owned by small owners (less than 1000 ha) and large forest owners between the two study areas.

Table 1. Size of study areas and distribution of land among different landowners. See figure 1 for an overview of the areas. Data from Markägarkartan (Landowner map) produced by Metria for the Swedish Environmental Protection Agency (Anon., 2005d).

	Bergslagen	Västerbotten
Total study area (ha)	5 105 001	4 757 969
Government (%)	2.3	37.7
Municipalities (%)	1.2	0.3
The Swedish church (%)	1.9	0.1
Commons (%)	0.6	5.4
Sveaskog	8.6	20.2
Bergvik skog väst AB (%)	15.7	-
Holmen skog AB (%)	0	0.8
Korsnäs AB (%)	2.5	-
SCA (%)	-	12.1
Other Corporations (%)	1.5	0.6
Other companies and commonly owned (%)	0.3	0.1
Landowners with less than 1000 ha (%)	65.4	22.7

The analysis based on sample plot data from the Swedish national forest inventory data shows that the potential amount of continuous cover forest land with old forest was very small (Table 2). There was more old forest left on all site types in the northern study area than in the southern. The result from the analysis of National Forest Inventory indicated less forest in the youngest age class for wet sites compared to mesic and dry sites (Table 2). Potential remaining CCF on dry and wet sites was 9-10% in the southern study area and about 22% in the northern with an additional 31-32% of high altitude oceanic climate forest (Table 2).

Table 2. Landscape scale forest age distribution (%) on different site types based on national forest inventory data in study areas Bergslagen and Västerbotten. Note that the “Amount site type” shows numbers in percent of all forest land and that the following lines show the age distribution of the different site types. The site types dry, mesic and wet (moist-wet) include all forest land (100%) and are based on the National Forest Inventory parameter “ground moisture” while climate is overlapping. Climate can thus include dry, mesic and wet land. Here climate means areas with a yearly precipitation of more than 800 mm in the Swedish mountain range. Other land is land where the national forest inventory variable could not be measured or estimated. The national forest inventory has different sample densities in the two areas.

Site type	Bergslagen			Västerbotten			
	dry	mesic	wet	dry	mesic	wet	climate
Amount of site type	5.5	90.7	3.8	5.8	90.5	3.8	3.2
Young Forest (age 0-40 years)	49.5	48.5	32.9	46.6	38.1	24.8	22.3
Growing Forest (41-80)	17.6	25.5	29.7	18.3	16.7	28.5	3.6
Production Forest (81-110)	19.8	15.6	24.3	10.0	12.4	9.7	12.3
Old prod. Forest (111-140)	6.4	6.4	6.4	14.6	17.8	23.2	29.4
Old Forest (141-)	5.9	1.1	3.5	10.4	12.8	11.7	31.5
Other forest land	0.7	2.9	3.1	0.0	2.2	2.2	0.9
SUM	100	100	100	100	100	100	100

Table 3. Forest age distribution on different site types in Bergslagen and Västerbotten based on mixed spatially explicit methods from two 50x100 km squares in the study areas (see figure 1). All numbers are given in percent (%). Note that the “Amount site type” shows the age distribution of the different site types. The category dry is based on soil type only and the category wet is based on a wetness index and the soil type. The site types dry, mesic and wet (moist-wet) include all forest land (100%) while climate is overlapping. Climate land can thus include dry, mesic and wet land. Here climate means areas with a yearly precipitation of more than 800 mm in the Swedish mountain range.

Site type	Bergslagen			Västerbotten			
	dry	mesic	wet	dry	mesic	wet	climate
Amount site type	12.9	81.6	0.3	5.8	88.9	0.4	2.4
Young Forest (age 0-40 years)	36.7	40.9	54.8	40.6	44.9	50.6	57.8
Growing Forest (41-80)	42.3	41.1	32.7	22.1	24.5	25.0	7.5
Production Forest (81-110)	20.6	17.7	12.2	24.0	20.5	18.0	12.6
Old prod. Forest (111-140)	0.3	0.3	0.3	13.0	9.9	6.3	20.0
Old Forest (141-)	0.01	0.01	0.00	0.4	0.4	0.2	2.0
SUM	100	100	100	100	100	100	100

According to the Topographic Wetness Index the southern study area was slightly drier than the northern study area (Figure 2). As indicated by the amount of younger forests the recent management of the different site types has been very similar in both study areas. For older age classes the results indicate lower amounts in the southern study area than in the northern. For sites located at high altitudes with moist oceanic climate it can be noted that there are larger parts with old production forest and old forests remaining than for any other forest type (see Table 3). The results from the coarse-grained spatially explicit analysis show only the age distribution on the different site types since the amount of the different site types has been chose from a calculated topographic wetness index (Figure 2). As a consequence this analysis says nothing about the proportion of the different site types in the landscape.




Figure 2. Total Wetness Index (TWI) in the two 50x100 km large squares in the study areas calculated from altitude and water bodies. The graph shows the distribution of land area in different wetness classes. In this graph all land except water bodies is included. The TWI ranges from dry (low numbers) to wet (high numbers).

Cultural landscape

The present amount of wooded grasslands with 10% or more tree cover (Anon., 2005b) in the county of Västmanland was estimated to 0.4-0.5% (about 2000 hectares) of today's forested land.

Stand scale

The forest management data from Non Industrial Private Forest (NIPF) owners and from large forest owners is very diverse in size (see Table 4). The stand scale analysis shows that there was more old forest left in the northern study area (see Table 5a and b).

Table 4. The number of stands and size of the total sample of different site/CCF types included in the stand scale study for different forest owners in the two study areas. All numbers in hectares (ha).

Site type	Poor-Dry		Mesic		Moist/Wet		Climate	
	stands	ha	stands	ha	stands	ha	stands	ha
Bergslagen								
Forest owner A	79	382	101	627	96	400	-	-
Forest owner B	665	3824	951	3298	263	1160	-	-
Private	66	44	267	486	49	143	-	-
Västerbotten								
Forest ownerA	132	1801	1458	19945	-	-	501	5552
Forest owner B	-	3020	-	51751	-	13576	-	-
Forest owner C	34	172	481	2888	43	144	-	-
Private	29	193	226	1648	31	132	76	544

Table 5a. Stand scale forest age distribution (%) estimates based on forest management plans for large forest owners and private forest owners. A and B are two large forest owners and Priv. is a group of private forest owners.

Site type	Bergslagen								
	dry			mesic			wet		
	A	B	Priv.	A	B	Priv.	A	B	Priv.
Young Forest (age 0-40 years)	67.9	42.2	31.9	73.5	22.4	52.1	43.7	46.4	56.8
Growing Forest (41-80)	16.1	9.3	18.4	17.2	26.9	15.7	12.9	50.7	38.9
Production Forest (81-110)	8.1	16.9	44.0	5.0	27.9	31.6	32.1	2.8	4.3
Old production Forest (111-140)	7.9	24.1	5.8	4.3	18.2	0.6	8.2	-	-
Old Forest (141-)	-	7.5	-	-	4.6	-	3.0	-	-

Table 5b. Stand scale forest age distribution estimates based on forest management plans for large forest owners and private forest owners. A and B are two large forest owners and Priv. is a group of private forest owners.

Västerbotten										
Site type	dry				mesic					
	A	B	C	Priv.	A	B	C	Priv.	A	B
Young Forest (age 0-40 years)	19.6	31.2	45.1	79.5	30.3	50.9	40.7	30.8	-	42.4
Growing Forest (41-80)	1.2	35.3	45.1	16.5	3.6	17.0	12.5	30.7	-	15.8
Production Forest (81-110)	32.4	11.7	10.8	0.5	9.0	10.8	4.2	9.2	-	14.7
Old production Forest (111-140)	29.4	12.7	0.0	2.1	51.1	13.6	15.2	13.5	-	19.7
Old Forest (141-)	17.3	9.2	15.3	1.3	15.0	7.6	27.4	15.8	-	7.3

Discussion

Continuous cover forests are rare

In Sweden continuous cover forests and forestry have recently been discussed by the National Forest Administration (Anon., 2004). The proposed definition is that the area extent of individual stands should be at least 0.25 ha and should have had a continuous forest cover with no major changes of tree species for at least 300 years. Further, the forest should have held a minimum of 30 m³/ha of living trees. In mixed deciduous forests there should be at least one species with over 10 m³/ha. The exception is cultural forests where the definition is not as strict.

The results from this study, together with the assumption that most of the wettest and driest sites were dominated by continuous cover forests, suggest that these forest types have declined sharply, and are still declining. The remaining amount of such forests in the southern study area today is somewhere between 0 and 6% of the potential (Table 6). In the northern study area the proportion left was slightly higher at 0 to 17 % of the potential, except for the high altitude continuous forest where the decline was even less (Table 6). For the county of Västmanland I estimated the past (about 1910) amount of cultural continuous cover forests to be 2-3% (> 10 000 hectares) of the presently forested area using old agricultural statistics (Anon., 1898; Anon., 1916) to be able get a rough estimate of the decline. My estimate shows a decline of 80% for this forest type (Table 6). The present management as seen by looking at the amount

of forest in the youngest age class shows that continuous cover forest sites to a large degree are managed in the same way as mesic sites (Table 7).

Table 6. Percent of area with no old forest on potential continuous cover forest sites with different analysis, scales and data sets.

Site type	Bergslagen			Västerbotten		
	Dry	Wet	Cultural	Dry	Wet	Climate
National Forest Inventory	94.1	96.5	-	89.6	88.3	68.5
Remote sensing	99.99	99.99	-	99.6	99.8	98.0
Agricultural statistics	-	-	80.9	-	-	-
Stand scale	92.5	97.0	-	89.2	81.8	64.5

Table 7. Young forest (0-40 years) (%) on different site types with different analysis methods. For the stand scale analysis the mean values from the different owners/owner categories has been used.

Site type	Bergslagen			Västerbotten			
	Dry	Mesic	Wet	Dry	Mesic	Wet	Climate
National Forest Inventory	49.9	48.5	32.9	46.6	12.8	28.8	22.3
Remote sensing	36.7	40.9	54.8	40.6	44.9	50.6	57.8
Stand scale	49.6	49.4	49.0	49.6	15.4	8.8	9.4

A main difference between the two regions studied is that the timber frontier arrived later in Västerbotten than in south-central Sweden, and especially in Bergslagen (Angelstam et al., 2004b). This is also clearly visible in the sample-plot and coarse-grained spatially explicit analyses (Table 2 and 3). However, because today's management seems to be similar among site types and owners, given enough time, the age distributions on different site types will probably look very similar in the two areas within a few decades.

Remote sensing as a tool to estimate and identify remaining continuous cover forests

One consequence of the paucity of continuous cover forests today is that they are hard to identify using remote sensing techniques. This is clearly shown if results from the sample-plot and coarse-grained spatially explicit analyses are compared (Table 2 and 3). The large differences in the oldest age classes in the sample-plot and coarse-

grained spatially explicit analysis are a result of the rarity of continuous cover forests and differences in the two landscapes. In the southern study area the remaining amount of old forest is smaller and more fragmented while in the northern study area the landscape consists of larger homogenous elements. Small fragments of old forest stands are hard or impossible to detect using the kNN analyzing method. In Bergslagen the forest is denser and the canopy closes at an earlier age. After the canopy has closed it is hard to see any differences between stands using remote sensing methods. Other methods like multispectral analyses of satellite images might be better options for identifying the last remaining biologically old forests.

Gap between policy, forest management and continuous cover forests

While contacting private forest owners to obtain forest management data a clear majority reacted negatively when they heard the words “continuous cover forestry”. This seemed to be due to a misconception that the current forest policy trend would force them to use continuous cover forestry on all their land. Continuous cover forestry is a concept that could be used in stands and areas where for some reason the desired values would not be maintained if all trees were cut. This could be due to ecological reasons like hosting species adapted to a continuous cover forest on the site types described in this study. It could also be since the forest have other functions than timber production such as protecting a water resource, protecting infrastructures such as roads and residential areas, reducing air and noise pollution, providing opportunities for recreation and tourism. Continuous cover forestry methods that retain more or less of the forests natural characteristics should be used on these sites.

Several forest companies in this study have expressed a clear interest in continuous cover forestry. Why is that? Is it because they want methods that mean they can produce some timber on sites that would else not be suited for forestry? Or is the reason to develop new silvicultural regimes? During the study I had contacts with forest managers that did not understand the difference between selection and selective cutting and its implication for the forests ecological values. Hence, if forest managers due to poor knowledge use alternatives to clear-felling whereby they select and cut all the most valuable trees the result will be more or less as bad as if all trees were cut, thus resorting to high-grading approaches commonly used in the mid 19th century.

The Swedish national forest statistics indicates that 96% of our forests are managed by clear-felling methods (variable retention and seed tree) (Anon., 2002b). This could mean that at the most 4% of the forests are managed with methods that attempt to mimic natural disturbance regimes. However, my study shows no sign that this is the case. Instead it shows that forestry methods are not as diverse as our forests and that our last remaining continuous cover forests will soon disappear or be small isolated patches in the landscape if management methods will not change. The change might already have started on wet sites. One forest manager in a large forest company told us that management has changed on some of the wet sites the last 15 years. This change might also be indicated in the results from the sample-plot analysis (Table 2.).

Conservation of continuous cover forests

If I like to conserve a sufficient amount of continuous cover forests and the connected biodiversity there is clearly a need for alternative management methods. For most habitats it is assumed to be critical when an 80% loss is reached (Andrén, 1994; Farig, 1997; Angelstam et al., 2004a). However, for continuous cover forests the critical number might be different since they more or less by definition are very fragmented also in a natural landscape. The very small areas with remaining old forest will need urgent protection with appropriate management if they are not protected already. For at least a part of the remaining areas with these site types restoration is needed. It will make a big difference for restoration purposes if the last remaining continuous cover forests could be used as core areas in the restorative process. The restoration can be achieved by protecting some areas. In other areas forestry methods that are based on the idea to emulate natural dynamic and thus maintain the characteristic structures of different forest types should be used. Both rich and wet sites as well as forests in humid macroclimate will sooner or later recover if left, even if the right management probably could speed up the process. If the management method could mimic natural processes some timber harvesting could take place.

By contrast, the conservation of cultural and poor-dry site continuous cover forests usually requires special management to maintain their biodiversity values. The cultural continuous cover forest needs to be kept open and cleaned from most of the coniferous growth. Ideally most of this could be done with grazing domestic animals.

It is for ecological reasons very important that we maintain this forest type at least until the remaining part of our forests will hold natural amounts of deciduous trees (Mikusinski et al., 2003). Scandinavian forests suffer from nitrogen deposition and the production capacity has clearly increased during the last 50 years (Binkley and Högberg, 1997; Solberg et al., 2004). Along with the fact that we have had a long time period without forest fires this means an accumulation of organic material that adds to making sites richer, which is a threat to the poor-dry continuous cover forest type. Naturally frequent low intensity forest fires burnt the dry lichens covering the ground. Most pioneer plants were kept away by this kind of low intensity disturbance and nutrients in the shape of soot particles often left the sites with wind and rain. The sites were thus maintained as poor-dry, to the benefit of specialised species (Berglind, 2004). Today forest fires are more or less absent, and thus this kind of continuous cover forest type is severely threatened. To conserve this dry site continuous cover forests some kind of selection cutting combined with burning could be used. Despite the results in this study this might be our most threatened continuous cover forest type since it does need active management to be maintained.

A need to improve the thematic resolution of forest management plans

My contacts with large forest companies showed that not all of them have the needed knowledge about continuous cover forests and forestry. Here I can see the need for more information. The thematic information in most forest management plans were also not enough for proper planning of the management of continuous cover forest sites. Given the current definition of sustainable forest management this is not acceptable. Most of the data have been collected once to calculate production oriented parameters. To meet society's current sets of values forest management plans need to be standardised and become more complete so they could be used as important tools for planning of ecological considerations and used to reach sustainable forest management. Ultimately they should develop into landscape management plans, including both terrestrial and aquatic dimensions.

To meet the current definition of SFM there is clearly a need for a broader set of management approaches than have been applied so far (Sayer and Maginnis, 2005). To specify the relative ambitions in terms of wood production and biodiversity management in forest stands with different site conditions in Sweden, the National

Board of Forestry, forest owners' associations and Sveaskog Co. use a system for management ambition attributes to be used in addition to the traditional site and wood production information in forest management plans (Anon., 2003; Ingemarsson, 2001). Currently four management attributes imply a gradient in the relative importance of wood production and biodiversity management: PG (production goal with general environmental considerations); PF (production goal with reinforced environmental considerations); NS (nature conservation goal with management); NO (nature conservation goal based on no management). One problem is that a forest management plan contains information on one property only while planning for ecological values need to consider a larger scale and where actual values are present or could be restored. The management ambition attributes system used by the National Board of Forestry, forest owners' association and Sveaskog Co. might be proper for very large properties or forest companies. For smaller properties this is however not an optimal approach (Ask and Nilsson, 2005).

Conclusion

The way forward

The multiple approaches employed in this study need to be developed further. First, I propose an extensive analysis of the potential amount of continuous cover forests using the available Swedish national forest inventory data and other relevant data. By combining several variables the definition of different forest types could be more specific than in this pilot study. The Swedish national kNN dataset (Reese et al., 2003, 2005) covers most parts of our country and a new data set will be available in a year or two. It is possible to use the data set as in this study with a modelled topographic wetness index with expanded study areas. Even a full country analysis is possible. Another interesting approach would be to use vegetation maps for areas where they are available. Several counties have produced these kinds of maps. Here one problem might be the diverse origin of these vegetation maps and thus potentially different quality. On the other hand the dry-poor and wet-rich sites should be quite easy to detect.

A second important thing to look at is the long-term trend for these kinds of forests. For at least a few different areas (where data is available) older datasets should be used to describe past trends over time, and estimate the trajectory into the future. This

information will be very useful when making decisions on how to conserve the remaining continuous cover forests, and to develop new.

For the cultural continuous cover forest data on the present situation as well as data on areas that could be restored is available for the whole country. Here it would be useful with some new approaches to estimate the past amount.

For forest management plans it is more or less impossible to collect large amounts of high quality data from non industrial private forest owners. Instead an enhanced co-operation with the larger forest companies, other large forest owners and the National Property Board should be the way forward. The aim should be to use their full forest management datasets to identify all stands on the different site types and compare the age distributions. One problem could possibly be that different forest companies have different parameters in their stand descriptions. The problem could be minimised by combining several parameters for a successful identification of the site types.

The future for our continuous cover forests and forestry

To fulfil the present forest policy the combination of different forest management methods used should allow the naturally occurring forest organisms to maintain viable populations. This means that sites naturally having large-scale disturbance regimes potentially could be managed using intensive methods like clear-felling combined with variable retention. However, the different continuous cover forest sites naturally were dominated by small-scale disturbance regimes. Most organisms in these forest types are adapted to this and would suffer from large stand replacing disturbances. Here new management approaches need to be implemented (Fries et al., 1997).

One problem appears to be that forest managers generally do not understand the differences between selection and selective cutting and its implications to continuous cover forests. The result is that these sites are managed for short-term economic gains instead of aiming to reach sustainable forest management. Long-term effective forestry with sustainable management of our forests all values can only be reached if owners and land managers collaborate according to their ability, and apply a diversity of management approaches (protection, management and restoration) at multiple spatial and temporal scales. Here I see a need for learning about how continuous cover

forestry is applied in regions with different forest ecosystems, land use history and systems for governance such as in Russia and the Alps.

Another challenge is limited knowledge on what kind of disturbances our flora and fauna needs and/or can manage. Some organisms need the disturbance or secondary effects from the disturbance, and for other organisms the same disturbance is a problem. Additionally we need better knowledge about thresholds for many species habitat requirements in terms of quality, quantity and configuration of habitats. Too much ecological care is costly and too little means that we lose some of our forests values. The same argument can be made for socio-cultural values.

References

- Aldentun, Y., 1997. Vegetationsregioner i Sverige: en historisk betraktelse. Redogörelse (Skogforsk 1997:6). Skogforsk, Uppsala.
- Andrén, H., 1994. The effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos* 71, 355–366.
- Angelstam, P., 1998. Maintaining and restoring biodiversity by developing natural disturbance regimes in European boreal forest. *Journal of Vegetation Science*, 9 (4): 593-602.
- Angelstam, P., 2003a. Forest biodiversity management - the Swedish model. In: Lindenmayer, D.B., Franklin, J.F. (Eds.) *Towards Forest Sustainability*, CSIRO Publishing, Canberra, and Island Press, Washington, pp. 143-166.
- Angelstam, P., 2003b. Reconciling the linkages of land management with natural disturbance regimes to maintain forest biodiversity in Europe. In: Bissonette, J.A. and Storch, I. (Eds.). *Landscape ecology and resource management: linking theory with practice*. Island Press, Covelo CA and Washington, D.C., pp. 193-226.
- Angelstam, P., Arnold, G.W., 1993. Contrasting roles of remnants in old and newly impacted landscapes: lessons for ecosystem reconstruction. In: Saunders, D.A., Hobbs, R.J., Ehrlich, P.R. (Eds.), *Reconstruction of Fragmented Ecosystems: Global and Regional Perspectives*. Surrey Beatty, pp. 109-125.
- Angelstam, P., Pettersson, B., 1997. Principles of present Swedish forest biodiversity management. *Ecological Bulletins* 46, 191-203.

- Angelstam, P., Anufriev, V., Balciauskas, L., Blagovidov, A., Borgegård, S.-O., Hodge, S., Majewski, P., Ponomarenko, S., Shvarts, E., Tishkov, A., Tomialojc, L., Wesolowski, T., 1997. Biodiversity and sustainable forestry in European forests - how west and east can learn from each other. *Wildlife Society Bulletin*, 25 (1), 38-48.
- Angelstam, P., Kuuluvainen, T., 2004. Boreal forest disturbance regimes, successional dynamics and landscape structures – a European perspective. *Ecol. Bull.*, 51, 117-136.
- Angelstam, P., Dönn-Breuss, M., Roberge, J.-M., (Eds.) 2004a. Targets and tools for the maintenance of forest biodiversity. *Ecol. Bull.* 51, 510 pp.
- Angelstam, P., Mikusinski, G., Fridman, J., 2004b. Natural forest remnants and transport infrastructure – does history matter for biodiversity conservation planning? *Ecol. Bull.*, 51, 149-162.
- Anon., 1898. Beskrifningar till kartorna öfver Vermlands län upprättade i Rikets Ekonomiska Karteverk åren 1887-97. Kungliga boktryckeriet, Stockholm.
- Anon., 1916. Beskrifningar till konceptkartorna öfver Västmanlands län upprättade i Rikets almäna kartverk åren 1909-1914. Kungliga boktryckeriet, Stockholm.
- Anon., 1974. Kalhyggen. Jordbruksdepartementet Ds Jo 1974:2. Stockholm.
- Anon., 1985a. Fälthäfte i bonitering Örebro och Västmanlands län, National board of forestry, Jönköping.
- Anon., 1985b. Fälthäfte i bonitering Kopparbergs och Gävleborgs län, National board of forestry, Jönköping.
- Anon., 1985c. Fälthäfte i bonitering Västerbottens län, National board of forestry, Jönköping.
- Anon., 1999. State of the world's forests (State of food and agriculture). Food and agriculture organization of the United Nations, Rome.
- Anon., 2001a. Skogspolitiken idag– en beskrivning av den politik och övriga faktorer som påverkar skogen och skogsbruket (Skogsstyrelsen Rapport 8B 2001). National board of forestry, Jönköping.
- Anon., 2001b. Svenska miljömål – delmål och åtgärdsstrategier (Regeringens proposition 2000/01:130. pp. 137-147.). The Swedish government, Stockholm.
- Anon., 2002a. En samlad naturvårdspolitik (Regeringens skrivelse 2001/02:173). The Swedish government, Stockholm.

- Anon., 2002b. Skogsvårdsorganisationens utvärdering av skogspolitikens effekter SUS 2001 (Meddelande I – 2002). National board of forestry, Jönköping.
- Anon., 2002c. Instruktion för fältarbetet vid riksskogstaxeringen. Swedish university of agricultural sciences, Umeå.
- Anon., 2003 Målklassning – hjälper skogsägare att bruka skogen långsiktigt (brochure), National board of forestry, Jönköping.
- Anon., 2004. Kontinuitetsskogar – en förstudie (Meddelande 1 2004). National board of forestry, Jönköping
- Anon., 2005a. Ängs- och betesmarksinventeringen 2002-2004 (Jordbruksverkets Rapport 2005:1). Swedish board of agriculture, Jönköping.
- Anon., 2005b. Global forest resources assessment update 2005 - Terms and definitions. Forestry Department, Food and Agriculture Organization of the United Nations, Rome.
- Anon., 2005c. September 16. MarkInfo [WWW database]. URL <http://www-markinfo.slu.se/index.phtml>
- Anon., 2005d. Frekvensanalys av skyddsvärd natur – Förekomst av värdekärnor i skogsmark (Rapport 5466). Swedish environmental protection agency, Stockholm.
- Antrop, M., 2005. Why landscapes of the past are important for the future. *Landscape and Urban Planning*, 70, 21-34.
- Ask, P., Nilsson, S.G., 2005. Det privata skogsbrukets landskap – En studie av uthålligt brukande i Stenbrohultsområdet. SUFOR, Alnarp.
- Bergeron, Y., Leduc, A., Harvey, B.D., Gauthier, S., 2002. Natural fire regime: a guide for sustainable management of the Canadian boreal forest. *Silva Fenn.*, 36, 81-95.
- Beven, K.J., Kirkby, M.J., 1979. A physically based variable contributing area model of basin hydrology. *Hydrol. Sci. Bull.*, 24, 43-69.
- Berglund, S.-Å., 2004. Area-sensitivity of the sand lizard and spider wasps in sandy pine heath forests - umbrella species for early successional biodiversity conservation? *Ecol. Bull.*, 51, 189-207.
- Binkley, D., Högberg, P., 1997. Does atmospheric deposition of nitrogen threaten Swedish forests? *Forest Ecology and Management*. 92, 119-152.
- Björklund, J., 1984. From the Gulf of Bothnia to the White sea - Swedish direct investments in the sawmill industry of Tsarist Russia. *Scand. Econ. Hist., Rev.* 32, 18-41.

- Blennow, K., 2004. Osäkerhet och aktiv riskhantering - Aspekter på osäkerhet och risk i sydsvenskt skogsbruk. SUFOR, Lund.
- Bolshakov, N.M., 2000. The recreational development of forests. Ministry of Education of the Russian Federation, Syktyvkar Forest Institute of St. Petersburg State Forest Technical Academy. Syktyvkar.
- Burton, P.J, Messier, C., Smith, D.W., Adamowicz, W.L. 2003. Towards Sustainable Management of the Boreal Forest. NRC research press, Ottawa. pp. 1039.
- Chao, G., Wang, H., Dai, L., Wu, G., Li, Y., Lang, R., Song, B., 2005. Integrating stand and landscape decisions for multi-purposes of forest harvesting. *Forest Ecology and Management*, 207, 233–243.
- Dengler, A., 1944. *Waldbau*. Springer Verlag, Berlin.
- Dengler, A., 1990. *Waldbau auf ökologischer Grundlage. II Baumartenwahl, Bestandesgründung und Bestandspflege* (6th ed. revised by E. Röhrig and H.A: Gussone). Verlag Paul Parey, Hamburg. 314 pp.
- Drushka, K., 2003. *Canada's Forests: A History*. Forest History Society, Durham.
- Ek, I., 1995. *Skinnskattebergs revir*. Hultebo Tryckeri AB, Skinnskatteberg.
- Ekelund, H., Hamilton, G., 2001 *Skogspolitisk historia (Skogsstyrelsen Rapport 8A 2001)*. National board of forestry, Jönköping.
- Ekstam, U., Forshed, N. 1996. *Äldre fodermarker*. Naturvårdsverkets förlag, Stockholm.
- Eliasson, P., 2002. *Skog, makt och människor: en miljöhistoria om svensk skog 1800-1875 (PhD Dissertation)*. Lund University, Lund.
- Falinski, J.B., 1986. *Vegetation dynamics in temperate lowland primeval forest. Ecological studies in Bialowieza Forest*. W. Junk. Dordrecht. 537 pp.
- Farig, L. 1997. Relative effects of habitat loss and fragmentation on population extinction. *Journal of Wildlife Management*, 61(3), 603-610.
- Forsell, H., Axelsson, B., 1990. *Sambruksformer för markanvändningsintressen inom Vilhelmina kommun (Inventering Handlingsprogram)*. Regional board of forestry mellannorrland, Lantbruksnämnden, Västerbotten county, Vilhelmina municipality, 105 pp.
- Franco-Lopez, H., Ek, A.R., Bauer, M.E., 2001. Estimation and mapping of forest stand density, volume, and cover type using the k-nearest neighbor method. *Remote Sensing of Environment*, 77 (3), 251-274.
- Fransson, S., 1965. The borderland. *Acta Phytogeographica Suecica*, 50, 167-175.

- Fries, C., Johansson, O., Pettersson, B., Simonsson, P., 1997. Silvicultural models to maintain and restore natural stand structures in Swedish boreal forests. *For. Ecol. Manage.*, 94, 89-103.
- Fries, C. et al., 1998. A review of conceptual landscape planning models for multiobjective forestry in Sweden. *Can. J. For. Res.*, 28, 159-167.
- Güntner, A., Uhlenbrook, S., Seibert, J., Leibundgut, C., 1999. Multi-criterial validation of TOPMODEL in a mountainous catchment. *Hydrological Processes*, 13 (11), 1603-1620.
- Hahn, T., 1998. Samernas markrättigheter i Sverige (Fakta Skog Nr 11 1998). Swedish university of agricultural sciences, Uppsala.
- Holmberg, L-E., 2005. Skogshistoria år från år 1177-2005 - Skogspolitiska beslut och andra viktiga händelser i omvärlden som påverkat Skogsvårdsorganisationens arbete (Skogsvårdsstyrelsen Rapport 5 2005). National board of forestry, Jönköping.
- Hägglund, B., Lundmark, J.-E., 1987. Handledning i bonitering med Skogshögskolans boniteringssystem (Del 1 Definitioner och anvisningar). National board of forestry, Jönköping.
- Hägglund, B., Lundmark, J.-E., 1999. Handledning i bonitering med skogshögskolans boniteringssystem (Del 3 Markvegetationstyper – Skogsmarksflora). National board of forestry, Jönköping.
- Hägglund, B., Lundmark, J.-E., 2003. Handledning i bonitering med Skogshögskolans boniteringssystem (Del 2 Diagram och tabeller). National board of forestry, Jönköping.
- Ingemarsson, F., 2001. Gröna planer (Skogsstyrelsen Rapport 8C 2001). National board of forestry, Jönköping.
- Kardell, L., 2004. Från baggböleri till naturvård (Svenskarna och skogen). Skogsstyrelsens förlag, Jönköping. pp. 303.
- Kirby, K.J., Watkins, C., (Eds.) 1998. The ecological history of European forests. CAB International, Wallingford, UK.
- Konijnendijk, C.C., 1997. Urban forestry: Overview and analysis of European forest policies. Part 1: Conceptual framework and European urban forestry history (Working Paper 12). European Forest Institute, Joensuu.
- Konijnendijk, C.C., 1999. Urban forestry: Overview and analysis of European forest policies. Part 2: Comparative analysis of policies and concepts in Europe.

- Contemporary urban forestry policymaking in selected cities and countries of Europe (Working Paper 20). European Forest Institute, Joensuu.
- Konijnendijk, C.C., 2003. A decade of urban forestry in Europe. *Forest Policy and Economics*, 5, 173–186.
- Korpilahti, E., Kuuluvainen, T., (Eds.) 2002. Disturbance dynamics in boreal forests: defining the ecological basis of restoration and management of biodiversity. *Silva Fennica*, 36 (1), 1-447.
- Kuuluvainen, T., 1994. Gap disturbance, ground microtopography, and the regeneration dynamics of boreal coniferous forests in Finland: a review. *Annales Zoologici Fennici*, 31, 35-51.
- Kuuluvainen, T., 2002. Natural variability of forests as a reference for restoring and managing biological diversity in boreal Fennoscandia. *Silva Fennica*, 36 (1), 97-125.
- Kuuluvainen, T., Syrjänen, K., Kalliola, R., 1998. Structure of a pristine *Picea abies* forest in north-eastern Europe. *Journal of Vegetation Science*, 9, 563-574.
- Kuuluvainen, T., Mäki, J., Karjalainen, L., Lehtonen, H., 2002b. Tree age distributions in old-growth forest sites in Vienansalo wilderness, eastern Fennoscandia. *Silva Fennica*, 36, 169-184.
- Lindhagen, A., 1996. Forest Recreation in Sweden. Four Case Studies Using Quantitative and Qualitative Methods (PhD dissertation). Inst. F. skoglig landskapsvård, Swedish University of Agricultural Sciences, Uppsala.
- Malcolm, D.C., Mason, W.L., Clarke, G.C., 2001. The transformation of conifer forests in Britain – regeneration, gap size and silvicultural systems. *Forest Ecology Management*, 151, 7–23.
- Mason, W.L., 2003. Continuous cover forestry: developing close to nature forest management in conifer plantations in upland Britain. *Scottish Forestry*, 57, 141-149.
- Matthews, J.D., 1989. *Silvicultural systems*. Oxford University Press, Oxford.
- McRoberts, R.E., Nelson, M.D., Wendt, D.G., 2002. Stratified estimation of forest area using satellite imagery, inventory data, and the k-nearest neighbours technique. *Remote Sensing of Environment*, 82 (Issues 2-3), 457-468.
- Mikusinski, G., Angelstam, P., Sporrang, U., 2003. Distribution of deciduous stands in villages located in coniferous forest landscapes in Sweden. *Ambio*, 33 (8), 520-526.

- Moore, I.D., Grayson, R.B., Ladson, A.R., 1991. Digital terrain modeling – a review of hydrological, geomorphological, and biological applications. *Hydrological Processes*, 5, 3-30.
- Nolbrant, P., 1998. Hamlade träd och solitärträd. In: *Skötselhandbok för gårdens natur och kulturvärden*. Swedish Board of Agriculture, Jönköping.
- Pennanen, J., 2002. Forest age distribution under mixed-severity fire regimes - a simulation-based analysis for middle boreal Fennoscandia. *Silva Fennica*, 36 (1), 213-231.
- Raivio, S., Normark, E., Pettersson, B., Salpakivi-Salomaa, P., 2001. Science and the management of boreal forest biodiversity – forest industries' views. *Scandinavian journal of forest research*. 16, Supplement 3, 99-104.
- Reese, H., Nilsson, M., Granqvist Pahlén, T., Hagner, O., Joyce, S., Tingelöf, U., Egberth, M., Olsson, H., 2003. Countrywide estimates of forest variables using satellite data and field data from the National Forest Inventory. *Ambio*, 32 (8), 542-548.
- Reese, H., Granqvist Pahlen, T., Egberth, M., Nilsson, M., Olsson, H., 2005. Automated estimation of forest parameters for Sweden using Landsat data and the kNN algorithm. In: *Proceedings, 31st International Symposium on Remote Sensing of Environment*.
- Rodhe, A., Seibert, J., 1999. Wetland occurrence in relation to topography - a test of topographic indices as moisture indicators. *Agricultural and Forest Meteorology*, 98-99, 325-340.
- Rydberg, D., Falck, J., 2000. Urban forestry in Sweden - A review. *Landscape and Urban Planning*, 47 (Issue 1-2), 1-18.
- Rydberg, D., 2001. *Skogens sociala värden (Skogsstyrelsen Rapport 8J 2001)*. National board of forestry, Jönköping.
- Rülcker, C., Angelstam, P., Rosenberg, P., 1994. *Ekologi i skoglig planering - förslag på planeringsmodell i Särnaprojektet med naturlandskapet som förebild (SkogForsk Redogörelse nr. 8.)*. Skogforsk, Uppsala.
- Sayer, J., Maginnis, S., 2005. *Forests in Landscapes - Ecosystem Approaches to Sustainability*. Earthscan Publications Ltd., London. pp. 348.
- Schuler, A., 1998. Sustainability and biodiversity - forest historical notes on two main concerns of environmental utilisation. In: *Bachmann, P., Köhl, M., Päivinen, R.*

- (Eds.), *Assessment of Biodiversity for Improved Forest Planning*. Kluwer Academic Publishers, Dordrecht, pp. 353–560.
- Selander, S., 1955. *Det levande landskapet i Sverige*. Albert Bonniers förlag, Stockholm.
- Sjörs, H., 1956. *Nordisk växtgeografi*. Albert Bonniers förlag, Stockholm.
- Solberg, S., Andreassen, K., Clarke, N., Tørseth, K., Tveito, O.E., Strand, G.H., Tomter, S., 2004. The possible influence of nitrogen and acid deposition on forest growth in Norway. *Forest Ecology and Management*, 192, 241–249.
- Sprugel, D.G., 1991. Disturbance, equilibrium and environmental variability: what is "natural" vegetation in a changing environment? *Biological Conservation*, 58, 1-18.
- Svensson, J., Fries, C., Jougda, L., 2004. Synthesis of the model forest concept and its application to Vilhelmina model forest and Barents model forest network (Rapport 6 2004). National board of forestry, Jönköping.
- Tomppo, E., 1991. Satellite imagery-based national inventory of Finland. *International Archives of Photogrammetry and Remote Sensing*, 28, 7-1, 419-424.
- Tucker, G.M., Evans, M.I, 1997. *Habitats for birds in Europe: a conservation strategy for the wider environment* (BirdLife Conservation Series no. 6.). BirdLife International, Cambridge, England.
- UN (United Nations), 1992. Report of the United Nations conference on environment and development (Annex III) - non-legally binding statement of principles for a global consensus on the management, conservation and sustainable development of all types of forests. United Nations, Rio de Janeiro.
- UN (United Nations Economic and Social Council), 2000. Report of the fourth session of the Intergovernmental Forum on Forests. United Nations, New York.
- Wallmo, U., 1897. *Rationell skogsafverkning. Praktiska råd till såväl större som mindre enskilde skogsägare, samt svar på en fråga för dagen*. CE Frizes Kongl. Hofbokhandel, Stockholm, pp 288.
- Wenner, M., 2000. *Public participation in forestry in Europe and North America* (Report of the team of specialists on participation in forestry). International Labour Office, Geneva.
- WCED, 1987. *Our Common Future*. Oxford University Press, Oxford
- Wieslander, G., 1936. The shortage of forest in Sweden during the 17th and 18th centuries. *Sveriges Skogsvårdsförbunds Tidskrift*, 34, 593-633.

- Williams, M., 2003. Deforesting the earth. Chicago university press, Chicago.
- Wilson, D.J., Western, A.W., Grayson, R.B., 2004. Identifying and quantifying sources of variability in temporal and spatial soil moisture observations. *Water Resources Research*, 40.
- Zinko, U., 2004. Plants go with the flow: predicting spatial distribution of plant species in the boreal forest (PhD dissertation). Swedish University of Agricultural Sciences, Umeå.
- Zinko, U., Seibert, J., Dynesius, M., Nilsson, C., 2005. Plant species numbers predicted by a topography based groundwater-flow index. *Ecosystems*, 8 (4), 430-441.
- Ångström, A., 1958. Sveriges klimat. General stabens litografiska anstalt, Stockholm.
- Östlund, L., Zackrisson, O., Axelsson, A.-L., 1997. The history and transformation of a Scandinavian boreal forest landscape since the 19th century. *Can. J. For. Res.*, 27, 1198-1206.
- Östlund, L., Zackrisson, O., 2000. The Forest History of Boreal Sweden – A Multidisciplinary Approach! In: Agnoletti, M. (eds.) *Methods and Approaches in Forest History*. IUFRO, Florence, Italy.

Publikationer från Institutionen för skogens produkter och marknader, Sveriges lantbruksuniversitet (SLU)

Rapporter

1. Persson, E. et al. 2002. Storage of spruce pulpwood for mechanical pulping. Part 1. Effects on wood properties and industrially produced pulp. Department of Forest Products and Markets, SLU, Uppsala
2. Pape, R. 2002. Rödkärna i björk – uppkomst, egenskaper och användning. *Red heart in birch – origin, properties and utilization*. Institutionen för skogens produkter och marknader, SLU, Uppsala
3. Staland, J. Navrén, M. & Nylinder, M., 2002. Resultat från sågverksinventeringen 2000. Institutionen för skogens produkter och marknader, SLU, Uppsala
4. Beck-Friis, M., et al. 2002. Skoglig logistik – Supply Chain Management i svensk skogssektor. Institutionen för skogens produkter och marknader, SLU, Uppsala
5. Orvér, M. 2002. Stickprovsmätning av skogsråvara – en praktisk handledning. Institutionen för skogens produkter och marknader, SLU, Uppsala
6. Lönnstedt, L. & Rosenqvist, H. 2002. Skatternas inverkan på skogsfastigheternas prisutveckling – Några hypoteser. Institutionen för skogens produkter och marknader, SLU, Uppsala
7. Hugosson, M. & Ingemarson, F. 2003. Depicting management ideas of private forest owners' – An assessment of general trends in Sweden based on new theoretical ideas. Department of Forest Products and Markets, SLU, Uppsala
8. Lind, T., et al. 2003. Storage of spruce pulpwood for mechanical pulping. Part 2. Effects of different sprinkling parameters on wood properties and pulp produced using a laboratory grinder. Department of Forest Products and Markets, SLU, Uppsala
9. Tascón Claro, Á. 2003. Pulpwood debarking. Department of Forest Products and Markets, SLU, Uppsala
10. Hultåker, O., Bohlin, F. & Gellerstedt, S. 2003. Ny entreprenad i skogen – bredda för bättre arbetsmiljö och lönsamhet. *New services for contracting in forestry – diversifying for better work environment and profitability*. Institutionen för skogens produkter och marknader, SLU, Uppsala
11. Bohlin, F. & Mårtensson, K. 2004. Askåterföring till skog, vardande blir verklighet? Institutionen för skogens produkter och marknader, SLU, Uppsala
12. Lönnstedt, L. & Nordvall, H.O. 2004. *The Japanese pulp and paper industry – An analysis of financial performance 1991-2001*. Institutionen för skogens produkter och marknader, SLU, Uppsala
13. Vestlund, K. & Hugosson, M. 2004. Produktutveckling för lönsammare sågverk – teori och ett praktikfall. *Product development for more profitable sawmilling -theory and a case study*. Institutionen för skogens produkter och marknader, SLU, Uppsala
14. Eriksson, P. 2004. Pilotstudie av drivningssystemet Besten och Kuriren – Slutavverkning med förarlös skördare manövrerad från skotare. Institutionen för skogens produkter och marknader, SLU, Uppsala
15. Edlund, J., Lindström, H. & Nilsson, F. 2004. Akustisk sortering av grantimmer med hänsyn till utbytets hållfasthet. Institutionen för skogens produkter och marknader, SLU, Uppsala
16. Roos, A. 2005. Forskning om marknadsorienterad innovation och produktutveckling inom svensk trävaruindustri – En kunskapsöversikt. *Research on market-oriented innovation and product development in the Swedish wood products industry – An overview*. Institutionen för skogens produkter och marknader, SLU, Uppsala
17. Wallin, A., & Nylinder, M. 2005. Träd- och virkesegenskaper hos två kloner av mikroförökad masurbjörk. Institutionen för skogens produkter och marknader, SLU, Uppsala
18. Hultåker, O. & Bohlin, F. 2005. Skogsmaskinentreprenörers diversifiering – Empiriska resultat och en tolkningsmodell. *Forest machine contractors' diversification – Empirical findings and a model*. Institutionen för skogens produkter och marknader, SLU, Uppsala
19. Edlund, J., Lindström, H. & Nilsson, F. 2005. Successiv uttorkning av stockar – inverkan på elasticitetsmodul. Institutionen för skogens produkter och marknader, SLU, Uppsala
20. Pivoriūnas, A. 2005. *Cooperation Among Private Forest Owners: Lithuania as a Case Study*. Licentiate thesis. Department of Forest Products and Markets, SLU, Uppsala
21. Tobisch, R., Hultåker, O., Walkers, M. & Weise, G. 2005. *Improvements of ergonomic assessment procedures for forest machines – A comparative evaluation of three established test methods*. Förbättringar av ergonomiska bedömningsystem för skogsmaskiner – En jämförande utvärdering av tre etablerade testmetoder. *Verbesserungen von ergonomischen Beurteilungsverfahren für Forstmaschinen – Eine vergleichende Bewertung von drei eingeführten Prüfmethoden*. Institutionen för skogens produkter och marknader, SLU, Uppsala
22. Roos, A., et al. 2005. *Workshop proceedings – Nordic Workshop on International Forest Processes*. Nordiskt forskarmöte om internationella skogliga processer 16-17 September, 2004. The Royal Swedish Academy of Agriculture and Forestry, Stockholm. Institutionen för skogens produkter och marknader, SLU, Uppsala
23. Roos, A., Törrö, M. & Rönneberg, J. 2005. *China's forest sector – A literature review*. Institutionen för skogens produkter och marknader, SLU, Uppsala

24. Lidén, E. 2005. *Benchmarks for good work organisation and successful implementation processes – Background to and working process of WORX*. Department of Forest Products and Markets, SLU, Uppsala
25. Vik, T. 2005. *Working conditions for forest machine operators and contractors in six European countries*. Department of Forest Products and Markets, SLU, Uppsala
26. Østensvik, T., et al. 2005. *Work exposure and complaints in a sample of French and Norwegian forest machine operators – A comparative field study within the ErgoWood programme*. Department of Forest Products and Markets, SLU, Uppsala
27. Jonsson, M. 2005. Lagring av barkat timmer. *Storage of debarked saw logs*. Institutionen för skogens produkter och marknader, SLU, Uppsala

Uppsatser

1. Eriksson, L. & Woxblom, L. 2002. Privatskogsbruk i Norrlands inland på 2000-talet. Institutionen för skogens produkter och marknader, SLU, Uppsala
2. Lewark, S. 2005. *Scientific reviews of ergonomic situation in mechanized forest operations*. Department of Forest Products and Markets, SLU, Uppsala
3. Bigot, M., et al. 2005. *Implementation and socio-economic impact of mechanisation in France and Poland – Synthesis*. Department of Forest Products and Markets, SLU, Uppsala
4. Walker, M. Tobisch, R. & Weise, G. 2005. *The Machine Operator Current Opinions and the Future Demands on Technical Ergonomics in Forest Machines*. Department of Forest Products and Markets, SLU, Uppsala
5. Kumm, J. 2005. *Implementation plan for ErgoWood. Research Notes No. 5*. Department of Forest Products and Markets, SLU, Uppsala

Examensarbeten

1. Törrö, M. 2002. Förändringar i skogsbranschens organisation på 1990-talet. Institutionen för skogens produkter och marknader, SLU, Uppsala
2. Svensson, H. 2002. Skogsbruksplanens betydelse för aktiviteten hos privata skogsägare i Älvdalen. Institutionen för skogens produkter och marknader, SLU, Uppsala
3. Sundblad, K. & Ekström, M. 2002. En marknadsundersökning om regelvirke – kvaliteter och kunduppfattningar. Institutionen för skogens produkter och marknader, SLU, Uppsala
4. Alvehus, A. 2002. Förslag till skötselplan för Uppsala högar och Tunåsen -ett exempel på medbestämmande planering. Institutionen för skogens produkter och marknader, SLU, Uppsala
5. Rosén, J. 2002. Kalkning och vitaliseringsgödsling. Institutionen för skogens produkter och marknader, SLU, Uppsala
6. Eriksson, J. 2002. Integration mellan skog & förädlingsindustri. Institutionen för skogens produkter och marknader, SLU, Uppsala
7. Paulsson, J. 2002. Den icke-monetära nyttans betydelse för prisbildningen på skogsfastigheter. Institutionen för skogens produkter och marknader, SLU, Uppsala
8. Paulmann, L. 2002. Julgransodlingar i Sverige – utbud, efterfrågan och lönsamhet. *Christmas tree plantations in Sweden - supply, demand and profitability*. Institutionen för skogens produkter och marknader, SLU, Uppsala
9. Hultåker, O. 2002. Skogsentreprenad idag och i framtiden – En kvalitativ studie av skogsmaskinentreprenörers verksamhet och framtidsvisioner. *Forest Contracting Today and in the Future – A qualitative Study of Logging Contractors' Activities and Their Visions of the Future*. Institutionen för skogens produkter och marknader, SLU, Uppsala
10. Ericsson, P. 2002. Skogsägares intresse för uppdatering av Gröna planer. Institutionen för skogens produkter och marknader, SLU, Uppsala
11. Warngren, K. 2002. Askåterföring värt besväret? – En fallstudie av följderna av Stora Enso's försöksverksamhet med askåterföring. *Ash recycling worth the trouble? – A case study on the consequences of Stora Enso's research and trials with ash recycling*. Institutionen för skogens produkter och marknader, SLU, Uppsala
12. Henriksson, J. 2003. Förändrad aptering av massaved från 3- till 4-meters längder vid gallring inom Södra. En systemanalys av effekter från avverkning till levererad virkesråvara. *Changed cross cut instruction of pulpwood from 3- to 4-meter lengths in thinning at Södra, a Swedish Forest Owner Association*. Institutionen för skogens produkter och marknader, SLU, Uppsala
13. Beck-Friis, M. 2003. Förskolors inställning till och användning av stadens natur. Institutionen för skogens produkter och marknader, SLU, Uppsala
14. Backman, M., 2003. Analys av orsak till nedklassning av granträvaror. Underlag för övergång till tväsidig sortering och automatsortering. Institutionen för skogens produkter och marknader, SLU, Uppsala
15. Backman, M. 2003. Analys av orsak till nedklassning av granträvaror. Underlag för övergång till tväsidig sortering och automatsortering. Institutionen för skogens produkter och marknader, SLU, Uppsala
15. Håkansson, B. 2003. Mobilt internet för skogsbruket med CDMA2000 i 450 MHz – bandet. Institutionen för skogens produkter och marknader, SLU, Uppsala
16. Jansson, J. 2003. Köpare av skogsfastigheter i Småland år 2000-2001 – En undersökning hur den privata ägarstrukturen ser ut i Sverige. *Buyer of forest properties in Småland the year 2000-2001 – A study of the private forestry holdings Sweden*. Institutionen för skogens produkter och marknader, SLU, Uppsala
17. Viklund, M. 2003. Hinder för svenskt trä inom den italienska byggbranschen i allmänhet och produktsegmenten fönster och dörrar i synnerhet. Institutionen för skogens produkter och marknader, SLU, Uppsala
18. Nilsson, F. 2003. Förbättrat råvaruutnyttjande vid kvalitetssortering av timmer – Utvärdering av analysprogrammet Stockholmen för automatiserad timmersortering i dimensions- och kvalitetsklasser hos BARO WOOD AB. *Improved quality*

- sorting of saw logs – Evaluation of the analyse program Stockholmen and the quality sorting of saw logs at BARO WOOD AB.* Institutionen för skogens produkter och marknader, SLU, Uppsala
19. Andersson, P. 2003. Omfattningen av icke avverkade områden i samband med slutavverkning. *The extent of non-cut areas at final cut operations.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 20. Fransila, J. 2003. Besökarstudie i Kilsbergens rekreatiomsområden – En metod för att utveckla rekreatiomsjöggheter på Sveaskogs marker. *Visitor survey in the recreation areas of Kilsbergen – A method to develop opportunities for recreation in the forests of Sveaskog.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 21. Eriksson, U. 2003. En intervju och enkätstudie av besökare i tre tätortsnära skogsområden i Stockholmstrakten. *Interviews and surveys in three urban forest areas in the Stockholm region.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 22. Blomqvist, L. 2003. Invandrare i tätortsnära natur – Kvalitativa intervjuer angående natursyn och nyttjande samt förslag till åtgärder. *Immigrants in nature close to urban settings – Qualitative interviews concerning views and utilization and proposed measures to increase usage.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 23. Nordin, H. 2003. Virkets formförändring och dess betydelse vid postning. Institutionen för skogens produkter och marknader, SLU, Uppsala
 24. López, J. 2003. *Forest fires and fire management in Sweden; a comparison with Spain.* Department of Forest Products and Markets, SLU, Uppsala
 25. Samuelsson, S. 2003. Uppfattningar om tryckved bland träbearbetande företag i Sverige. *Perception of compression wood among sawmills and wood-manufacturing companies in Sweden.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 26. Sjölander, H. 2003. Ändamålsanpassad TINA-sortering av sågtimmer. *Enduse orientated gamma-ray sorting of sawlogs.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 27. Toikkanen, C. 2003. Rekryteringsstrategier för företag inom skogssektorn – en undersökning om hur skogsbrukande och träförädlade företag bygger sitt arbetsgivarvarumärke. Institutionen för skogens produkter och marknader, SLU, Uppsala
 28. Svedberg, P. 2003. Hur uppfattas pcSKOG AB och pcSKOG-gård av privata skogsägare? En undersökning av en programvara för privatskogsbruket. *How are pcSKOG AB and pcSKOG-gård apprehended by private forest-owners? A study of a software for private forest estates.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 29. Bauer, M. 2003. Den geografiska, funktionella och processororienterade organisationen; En fallstudie av Holmen Skog, SCA Skog och Sydkraft Vattenkraft. Institutionen för skogens produkter och marknader, SLU, Uppsala
 30. Althoff, D. 2004. Sambandet mellan bostadsbyggandet och konsumtionen av sågade barrträraror i några av Europas länder. Institutionen för skogens produkter och marknader, SLU, Uppsala
 31. Lindow, K. 2004. Ekonomisk konsekvensanalys av sprickor. I samband med avverkning och sågverksproduktion. Institutionen för skogens produkter och marknader, SLU, Uppsala
 32. Eriksson, H. & Kreij, E. 2004. Möjliga strategier för Holmens framtida skogsägande. Institutionen för skogens produkter och marknader, SLU, Uppsala
 33. Kogler, F. 2004. Färsk ved till Hallstaviks pappersbruk. *Fresh wood to Hallstaviks papermill.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 34. Forsbäck, M. 2004. Direktmarknadsföringens alternativ – En fallstudie för Logosol AB. *Direct marketing alternatives – A case study at Logosol.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 35. Jansson, A. 2004. Privata markägares attityder och inställningar till förnyrningsfrågor – En studie utförd i Mälardalen. Institutionen för skogens produkter och marknader, SLU, Uppsala
 36. Arvidsson, C. 2004. Attityder hos råvaruleverantörer till ett sågverksföretag – En fallundersökning av leverantörer till J.G. Anderssons Söner AB i Kronobergs län. *Attitudes among primary product suppliers to a sawmilling company – A case study among of suppliers to J.G Andersson's Söner AB in Kronobergs län.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 37. Berggren, A. 2004. Modeller för brösthöjdsålder för tall och gran. *Prediction models for breast height age for Scots Pine and Norway Spruce.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 38. Lundin, M. 2004. En studie av besöksantalet i tre tätortsnära skogar i Stockholmsområdet med hjälp av Radio Beam Counter – Ett räkneverk baserat på radiovågsteknik. *A study of the number of visitors in three urban woods in the Stockholm area using Radio Beam Counter technique.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 39. Sigurdh, M. 2004. Mekaniserad plantering med Eco-Planter i södra Sverige. *Mechanized planting with Eco-Planter in southern Sweden.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 40. Gunnarsson, F. & Mårtenson, C. 2004. Vilka mål och behov har olika typer av skogsägare kring sitt skogsägande? *Which goals and needs have different types of forest owners?* Institutionen för skogens produkter och marknader, SLU, Uppsala
 41. Carlsson, P. 2005. Möjligheter att öka effektiviteten och det ekonomiska utfallet av barkhanteringen vid Seskarö sågverk. *Possibilities to increase the efficiency and profitability regarding the bark handling at Seskarö sawmill.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 42. Lundquist, J. 2005. Kommunägd skog i Sverige – en enkät- och intervjustudie av de tätortsnära skogarnas ekonomiska och sociala värde. *Municipality owned forest in Sweden – a questionnaire and interview study of social and economic values of the urban forests.* Institutionen för skogens produkter och marknader, SLU, Uppsala
 43. Selmeryd, O. 2005. Efterfrågan av grova sågade dimensioner och hyvlade produkter bland Wallnäs AB:s kunder – En marknadsundersökning. Institutionen för skogens produkter och marknader, SLU, Uppsala
 44. Norström, D. & Gustafsson, K. 2005. *Latvian logging companies – present state and development needs.* Skogsavverkningsföretag i Lettland – dagsläge och utvecklingsmöjligheter. Institutionen för skogens produkter och marknader, SLU, Uppsala

45. Delavaux, H. 2005. *Cultivation of trees as a way to achieve diversification for smallholdings in Nicaragua*. Institutionen för skogens produkter och marknader, SLU, Uppsala
46. Göransson, P. 2005. Värdering för markåtkomst vid järnvägs- och motorvägsbyggnation En fallstudie av intrångsvärdering i området mellan Örebro och Arboga. *Valuation of ground rights when building railway and highway – A case study of infringement valuation in the area between Örebro and Arboga*. Institutionen för skogens produkter och marknader, SLU, Uppsala
47. Eriksson, M. 2005. Sveaskogs möjligheter att utveckla träbränslevarksamheten i Västerbotten och södra Norrland. *Sveaskog's possibilities to increase the wood fuel activity in Västerbotten and southern Norrland*. Institutionen för skogens produkter och marknader, SLU, Uppsala
48. Andersson, L. & Kumm, E. 2005. *Estonian logging companies - An exploratory survey of the Estonian logging companies*. Institutionen för skogens produkter och marknader, SLU, Uppsala
49. Prejer, B. 2005. Utveckling av ett skogsbolags kontaktstrategi. En kvalitativ intervjustudie bland större privata virkesleverantörer. *Development of the contact strategy of a forest company. A quality study among large timber suppliers*. Institutionen för skogens produkter och marknader, SLU, Uppsala
50. Johansson, P. 2005. Affärsupplägg biobränsle Västerbotten - En undersökning av större biobränsleanvändares syn på biobränslemarknaden i Västerbotten. *Business conditions for bio energy in Västerbotten – A survey of larger bio energy consumers' views of the bio energy market in Västerbotten*. Institutionen för skogens produkter och marknader, SLU, Uppsala
51. Andersson, C. 2005. Bioenergi från röjningsgallringar, en jämförande studie av fyra flödeskedjor från avlägg till förbrukare. Institutionen för skogens produkter och marknader, SLU, Uppsala
52. Ek, K. & Furness-Lindén, A. 2005. Syns vi – finns vi!? – Marknadsföringsstrategier för Svenska FSC. *Marketing Strategies for FSC Sweden*. Institutionen för skogens produkter och marknader, SLU, Uppsala
53. Loré, J. 2005. Tillämpning av naturvårdsavtal. *Application of nature conservation agreements*. Institutionen för skogens produkter och marknader, SLU, Uppsala
54. Vidmo, M. 2005. Röjningsförbandets betydelse för avverkningsekonomin i södra Sverige. Institutionen för skogens produkter och marknader, SLU, Uppsala
55. Bager, H. 2005. *An inventory of Non- Wood Forest Products used by people living in the buffer zone of a national park in the Amazonian Peru – assessment on subsistence and ecology*. Institutionen för skogens produkter och marknader, SLU, Uppsala
56. van Soest, M. 2005. *The European sawmill industry in a global competitive market: perspectives with regard to Monterey pine plantations in the Southern hemisphere*. Institutionen för skogens produkter och marknader, SLU, Uppsala
57. Wahn, J. 2005. Strategisk/Taktisk vägplan. Institutionen för skogens produkter och marknader, SLU, Uppsala
58. Blicharska, M. 2005. *Using a Swedish forest biodiversity assessment under Polish conditions*. Institutionen för skogens produkter och marknader, SLU, Uppsala
59. Lennartsson, A. 2005. Val av tidpunkt för markberedning vid naturlig förnyring under skärm av *Pinus sylvestris* i Svealand. *Timing of scarification when using natural regeneration in seed tree stands of Pinus sylvestris in Central Sweden*. Institutionen för skogens produkter och marknader, SLU, Uppsala
60. Bergh, J. 2006. Vad tycker skogsägare om virkesinköpare och inköpsorganisationer? *Private forest owners' opinion about forest purchaser and wood supply organisations*. Institutionen för skogens produkter och marknader, SLU, Uppsala
61. Ureña Lara, F.J. 2006. *Spanish Woodworking Industry – Geographical structure, Export and Import*. Institutionen för skogens produkter och marknader, SLU, Uppsala
62. Åkesson, J. 2006. Prislisteoptimering för ett sågverk – Jarlträ AB. *Optimization of timber price lists for a sawmill – Jarlträ AB*. Institutionen för skogens produkter och marknader, SLU, Uppsala
63. Mörner, G. 2006. Kinas intåg på skogsvarumarknaden – Idag och i framtiden. Institutionen för skogens produkter och marknader, SLU, Uppsala
64. Frank, N. 2006. Underröjning i förstagallring. *Cleaning of understorey trees before thinning*. Institutionen för skogens produkter och marknader, SLU, Uppsala
65. Karlsson, P. & Sylén, O. 2006. Skogsmaskinens bränsleförbrukning. *Forest machines' fuel consumption*. Institutionen för skogens produkter och marknader, SLU, Uppsala
66. Karlsson, F. 2006. Privata markägares reflektioner med hänseende till den minskade röjningsaktiviteten – så kan skogsvårdsstyrelsen anpassa sitt arbete. *Family foresters' thoughts concerning the decreasing activity in precommercial thinning – how the Swedish forestry board can adjust its work*. Institutionen för skogens produkter och marknader, SLU, Uppsala
67. Axelsson, R. 2006. *Natural and cultural continuous cover forests in Sweden – how much remain and how are they managed?* Institutionen för skogens produkter och marknader, SLU, Uppsala