



Sveriges lantbruksuniversitet  
Swedish University of Agricultural Sciences

**Faculty of Veterinary Medicine and Animal  
Science**

# Effects of extruded linseed on methane emission and milk production of dairy cows

Effekter av extruderat linfrö på metanemission och mjölkproduktion hos mjölkkor

Stina Blomdahl

**Degree project • [45]hp**

Master in Animal Science

Department of Animal Nutrition and Management

Swedish University of Agricultural Sciences Uppsala 2019



# Effects of extruded linseed on methane emission and milk production of dairy cows

## Effekter av extruderat linfrö på metanemission och mjölkproduktion hos mjölkkor

Stina Blomdahl

**Supervisor:** Kjell Holtenius, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management  
**Assistant supervisor:** Maria Åkerlind, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management  
**Examiner:** Rolf Spörndly, Swedish University of Agricultural Sciences, Department of Animal Nutrition and Management

**Credits:** [45] credits  
**Level:** Advanced E  
**Course title:** Degree project in Animal Science  
**Course code:** EX0653  
**Programme/education:** Agronomprogrammet - Husdjur  
**Course coordinating department:** Department of Animal Nutrition and Management

**Place of publication:** Uppsala  
**Year of publication:** 2019  
**Cover picture:**  
**Online publication:** <https://stud.epsilon.slu.se>

**Keywords:** Extruded linseed, methane output, fatty acid, milk yield



## Innehållsförteckning

|  |    |
|--|----|
| Abbreviations .....  | 1  |
| Abstract.....  | 3  |
| Sammanfattning.....  | 3  |
| 1. Introduction and background.....  | 4  |
| 2. Literature review.....  | 5  |
| 2.1 Why is methane production a problem? .....                                 | 5  |
| 2.2 Why is methane produced in rumen?.....                                     | 5  |
| 2.2.1 Methane production from cows .....                                       | 5  |
| 2.2.2 Methanogenesis.....  | 5  |
| 2.3 Possibilities to reduce methane output<br>through feed ratio .....         | 6  |
| 2.3.1 Dietary fat effect on methane output in<br>general .....                 | 7  |
| 2.3.2 Linseeds.....  | 8  |
| 2.4. Effects of linseed feed ration on milk<br>yield and fat composition ..... | 9  |
| 2.4.1 Milk yield .....   | 9  |
| 2.5 Milk fat composition .....   | 9  |
| 2.5.1 Lipids.....  | 9  |
| 2.5.2 Dietary fat effect on milk fat composition.....                          | 10 |
| 2.6 How is methane measured from cows?.....                                    | 11 |
| 2.6.1 Methane measurements .....   | 11 |
| 2.6.2 Influencing factors of methane measuring<br>.....                        | 13 |
| 3. Materials and Methods.....  | 13 |
| 3.1 Experimental design and herds .....  | 13 |
| 3.1.1 Treatment feed.....  | 14 |
| 3.1.2 Selected cows.....   | 14 |
| 3.2 Measurements and sampling .....  | 15 |
| 3.2.1 Analyses on feed and milk .....  | 15 |
| 3.2.2 Methane measurement .....  | 16 |
| 3.2.3. Indirect methane estimations with<br>Visiolait.....                     | 17 |
| 3.2.4 Indirect methane estimations by NorFor .....                             | 17 |
| 3.3 Statistical calculations .....   | 17 |

|   |    |
|---|----|
| 4. Results .....  | 18 |
| 4.1 Descriptive data.....   | 18 |
| 4.2 Milk yield and milk composition .....                                     | 20 |
| 4.2.1 Milk yield .....  | 20 |
| 4.2.2 Protein.....  | 20 |
| 4.2.3 Unsaturated fatty acids in milk .....                                   | 20 |
| 4.2.4 Saturated milk fatty acids .....  | 21 |
| 4.3 Methane output.....   | 22 |
| 4.3.1 Estimated methane value through fatty<br>acid analyses, Visiolait ..... | 23 |
| 4.3.2 Estimated methane value through NorFor<br>.....                         | 23 |
| 5. Discussion.....  | 25 |
| 5.1 Effect of Extruded linseed.....   | 25 |
| 5.1.1 Milk Yield.....   | 25 |
| 5.1.2 Milk fat.....   | 25 |
| 5.1.3 Methane output .....  | 26 |
| 5.2.1 Herds and experimental design .....                                     | 26 |
| 5.2.2 Number of cows .....  | 27 |
| 5.2.3 Sampling .....  | 27 |
| 5.2.4 Feed.....   | 28 |
| 5.3 Future .....  | 28 |
| 6. Conclusion .....   | 29 |
| 7. Acknowledgements.....  | 29 |
| 8. References.....  | 30 |

## Abbreviations

AMS-automatic milking system

CIP-Competitive and innovation frame work programme

CO<sub>2</sub>-eq-carbon dioxide equivalence

*de novo*- “from the new”

DIM-days in milk

DM-dry matter

DMI-dry matter intake

EL-extruded linseed

FA-fatty acid

FFA-free fatty acids

FPCM-fat and protein corrected milk

FTIR-Fourier Transform Infra-red analyse. Analyse distributed by Agrosom via Eurofins Steins laboratory.

FAT<sub>tot</sub>- fat concentration in milk for cows in testgroup. Analyse from FTIR

FA<sub>pr</sub>- protein concentration in milk for cows in testgroup. Analyse from FTIR

ECM-Energy corrected milk

GC-Gas *Chromatography*

GHG-greenhouse gas

GLM-General linear model

*in vitro*-experiment done in laboratory

*in vivo*-experiment done inside the body

LCFA-long-chain fatty acids

LMD-laser methane detector

MF-milk fat content from official Swedish milking recording scheme distributed via VäxaSverige

MP-milk protein content

MY-milk yield

MUFA-monounsaturated fatty acid

NDF-Neutral detergent fibre

Per-period

PUFA-poly unsaturated fatty acids

PMR-partly mixed ration

SFA-saturated fatty acids

TMR-total mixed ration

UFA-unsaturated fatty acids

VFA-volatile fatty acids

VLDL-very-low density lipoprotein

## Abstract

The agriculture sector is one of the major sources to produce greenhouse gases of today. One big contributor is methane (CH<sub>4</sub>) generated during rumen digestion of the feed carbohydrates. One approach to lower CH<sub>4</sub> output from the agriculture sector, is to change the diet for the cows. Diets with extruded linseed (EL) have in earlier studies showed to have a lowering effect on CH<sub>4</sub> output from cows. Previous studies have shown correlations between CH<sub>4</sub> output and fatty acid (FA) profile of milkfat. EL has also shown in previous studies to influence the FA profile in milkfat. The objects of this study were to evaluate if CH<sub>4</sub> output can be reduced if diets are supplemented with extruded linseed, if CH<sub>4</sub> increases the concentration of unsaturated FA in milk fat, and the effect of EL on milk yield.

In this study a total of 177 cows from five herds were individually scanned for CH<sub>4</sub> output, milk fatty acid profile and milk yield. Cows were studied while fed a control diet, during the treatment diet with EL and when going back to the control diet again. EL diets had no effect on milk yield but increased the unsaturated FA in the milk. EL also decreased the CH<sub>4</sub> output from the cows with 22% from the first control period to the treatment diet, however there was no increase of CH<sub>4</sub> output between treatment period and second control period. Conclusion of this master thesis is that Extruded linseed did not affect the milk yield. EL had an effect on milk fat composition. Due to the unreliable measuring method of methane this study could not point out an obvious decrease of CH<sub>4</sub> due to EL.

Key words: extruded linseed, methane output, fatty acid, milk yield.

## Sammanfattning

Lantbruket är en betydande källa till dagens globala utsläpp av växthusgaser. Växthusgasen metan produceras i kons våm när fodret smälts. Ett sätt att minska dessa utsläpp från lantbrukssektorn är att förändra foderstaten för boskapen. Att inkludera extruderat linfrö (EL) i foderstaten har i tidigare försök visat sig minska metanutsläppet. Tidigare försök har även visat att det finns en korrelation mellan metanutsläpp och mjölkfettets fettsyreprofil samt att EL påverkar mjölkfettets fettsyreprofil. I denna studie har totalt 177 kor från 5 gårdar medverkat för att undersöka ELs effekt på mjölkavkastning, fettsyrasammansättningen i mjölken och metanutsläpp från kor. Kornas metanutsläpp mättes totalt tre gånger under testperioden. Första gången under en kontrollfoderstat andra gången under behandlingsfoderstat med EL och slutligen under ytterligare en kontrollfoderstat. Denna studie visar att EL inte hade någon effekt på mjölkavkastningen. EL ökade de omättade fettsyrorna och minskade de mättade fettsyrorna i mjölken. EL minskade metan utsläppen per kg ECM med 22 % mellan första kontrollperioden och den följande perioden med behandlingsfoderstaten. Dock erhöles ingen förändring i metanutsläpp då korna åter fick kontrollfoderstaten. Slutsatsen av denna

studie är att EL hade en effekt på mjölkfettsyraprofilen. På grund av mätmetoden kunde inte studien visa på ett tydligt minskat metanutsläpp beroende på EL.

## 1. Introduction and background

Global warming is a big threat for the world today. The underlying reason is the increasing concentration of greenhouse gases (GHG) in the atmosphere. These GHG are among others carbon oxide (CO<sub>2</sub>), CH<sub>4</sub> and nitrous oxide (NO<sub>2</sub>) that “traps” the heat in the atmosphere and results in global warming (Lovett et al., 2005; IPCC, 2007 (Climate Change 2007: Synthesis Report). An increased global temperature has already resulted in increased melting of ice and rising of average sea levels (IPCC 2007). Output of GHG have a natural or an anthropogenic origin (Bousquet 2006). Example of natural GHG output is volcanic eruptions and wetlands (IPCC Arturo 2007 chapter 2). Leading anthropogenic GHG output is energy supply, 25.9 %, industry, 19.4 %, forestry and deforestation, 17.4 % and on fourth place agriculture sector with 13.5 % (IPCC 2007). The global dairy sector alone stands for of 4 % of the total GHG emissions however large differences exists between different parts of the world (FAO, 2010).

There are many potential ways to decrease GHG emissions from the agricultural sectors. They involve further intensifications of livestock production by breeding, feeding and management that results in using less animals (FAO, 2006).

Feeding strategies aiming at lowering CH<sub>4</sub> include the use of fat supplements (as reviewed by Toprak (2015) and Knapp et. al (2013). The effects on CH<sub>4</sub> output vary depending on fat source (Chilliard et al, 2009).

Unsaturated fats of linseeds have shown to decrease ruminal CH<sub>4</sub> output depending on how the linseeds are processed (Martin et al, 2008). Fat supplements have further shown to affect the milk fat composition, and a positive correlation between a differed milk fat composition and a decreased CH<sub>4</sub> output (Chilliard et al. 2009).

The objects of this study were to evaluate if linseed supplemented to feed could; decrease CH<sub>4</sub> output, increase unsaturated FA in milk fat and increase the milk yield.

Hypotheses:

- 1) Extruded linseed increases the proportion of unsaturated fat in milk fat and decreases the saturated fat.
- 2) Extruded linseed increases the milk yield.
- 3) Extruded linseed supplementation reduces CH<sub>4</sub> emissions per unit milk.

## 2. Literature review

### 2.1 Why is methane production a problem?

CH<sub>4</sub> is a smell and colourless gas which is produced during breakdown of organic material (Nationalencyklopedien, 2016). The production is done by microorganisms only in anaerobic environments like soil, swamps and sediment and in the rumen of cattle (Sjaastad, 2010). CH<sub>4</sub> is one of the major greenhouse gases and have the warming effect of 28 (±40%) (IPCC, 2013) times the warming effect of CO<sub>2</sub> on a 100-year time horizon (FAO, 2010: IPCC2013). Different greenhouse gases are often expressed as CO<sub>2</sub>-equivalents (-eq), where the GHG is multiplied with its warming potential (IPCC, 2007).

There are many ways to express GHG in literature. One way, which is used by FAO, is to calculate GHG emissions per product or unit. This is also called the carbon footprint (FAO, 2010).

The average carbon footprint of global milk yield is calculated to 2.4 kg CO<sub>2</sub>-eq per kg of fat and protein corrected milk (FPCM) (FAO 2010). However, this figure differs due to differences in production circumstances. Regions as Sub-Saharan Africa have a carbon footprint of 7.5 kg CO<sub>2</sub>-eq per kg FPCM compared to 1-2 kg CO<sub>2</sub>-eq per kg FPCM in industrialised regions like Sweden (FAO, 2010). The biggest contribution, 40 %, of GHG from cattle is the enteric gases (Greber et al. 2013 cited in Danielsson, 2016).


### 2.2 Why is methane produced in rumen?

#### 2.2.1 Methane production from cows

Every day the rumen produces approximately 2000-4000 litres of different gases. These gases are in order of quantity: 40 % (CO<sub>2</sub>), 30-40 % of CH<sub>4</sub>, 5 % hydrogen (H<sub>2</sub>) and small amounts of oxygen and nitrogen (Sjaastad et al., 2010: Mc Donald, 2011). Gases in the rumen are produced by microbes that ferments the plant carbohydrates. The gases are emitted from the cow through belching, also called eructation. A large portion of the gas is inhaled into the lungs and are then exhaled throughout the nose and mouth (Sjaastad, 2010). Not all gas produced in the rumen exits through eructation. Around 20 % of CH<sub>4</sub> leaves the body through feces (Sjaastad et al, 2010). Enteric emissions are however unavoidable in the process of turning grass and grains into food for humans (Sjaastad, 2010).


### 2.2.2 Methanogenesis

There are three classes of microbes in the rumen; bacteria, protozoa and fungi. These microbes, and their enzymes, break down dietary carbohydrates to volatile fatty acids (VFA) mainly propionate, acetate and butyrate (Figure 1).



**Figure 1.** fermentation of glucose to VFA (Boadi et al., 2004; Moss et al., 2000)

The produced VFA is used by the cow as energy and for milk synthesis. The H<sub>2</sub> is accumulating in the rumen and limits further fermentation of feed. However, the methanogenic bacteria use H<sub>2</sub> for CH<sub>4</sub> production (figure 2) and the excessive H<sub>2</sub> is removed from rumen through eructation (McAllister & Newbold, 2008).



**Figure 2.** Methanogenesis (Boadi et al., 2004)

There are several types of methanogens that uses other substrate than H<sub>2</sub> for the methanogenesis. The substrate used besides H<sub>2</sub> is example formate and methanol (Liu & Whitman, 2008; Lang et al., 2015). Earlier trial show that type of rumen methanogens can be a reason of variations in CH<sub>4</sub> output. Danielsson (2012) has shown that there might be differences in CH<sub>4</sub> output between individual cows.

Although CH<sub>4</sub> production is an efficient way to reduce H<sub>2</sub> in rumen the process is energy demanding. An alternate way of H<sub>2</sub> could bring, besides reducing CH<sub>4</sub> output, an improved feed efficiency. Acetate and butyrate production produce H<sub>2</sub>, however propionate production uses H<sub>2</sub> (Figure1). This dietary arrangement altering the VFA toward a higher production of propionate is one way to decrease CH<sub>4</sub> output (Johnsson & Johnsson 1995).

### 2.3 Possibilities to reduce methane output through feed ratio

There are a few feed factors that have a significant effect on enteric CH<sub>4</sub> output. One is the positive relationship between DMI and CH<sub>4</sub> output is generally accepted. This is due to the increased amount of substrate in rumen that is fermented (Moss et al., 1995; review by Ramin & Huhtanen, 2013). However, cows with high milk yield generally also have a high DMI

and thus CH<sub>4</sub> output is diluted per unit of product. A high digestibility is correlated with a lowered CH<sub>4</sub> output. Also, here is an increased digestibility connected to high producing cows and gives therefore a lower CH<sub>4</sub> output per unit.

Type of feed affects the CH<sub>4</sub> output. Feed ratios with a high proportion of starch rich concentrate is found to be correlated to a lower CH<sub>4</sub> output and high milk yield (Johnson & Johnson 1995). Starch rich diets have shown to increase propionic acid in rumen (Penner et al., 2009). Production of propionic acid works as a H<sub>2</sub> sink compared to acetic and butyrate acid production as mentioned above. Other factor effecting CH<sub>4</sub> output is how the forage is processed. Silage compared to hay decreases the enteric CH<sub>4</sub>. The fermentation process of silage was found to give a variation in acids in the silage. Silage high in total acids or high in acetate will lower the CH<sub>4</sub> production from cows (Ramin & Huhtanen, 2014).

Feed supplements in form of lipids is reviewed as one of the leading strategies to reduce CH<sub>4</sub> output (Beuchemin et al., 2008; Ramin & Huhtanen, 2013).

### 2.3.1 Dietary fat effect on methane output in general

Fats in different forms and volumes have shown to influence and decrease enteric CH<sub>4</sub> output. Beauchemin et al (2008), gathered data from 17 studies and found that for each supplemented percentage of fat comes a decrease of CH<sub>4</sub> by 5.6 %. However, diets containing higher amount than 5% fat affects the cow by a reduced appetite, decreased fermentation of feed and diminished the motility of the fore stomachs (Sjaastad, 2010). Comparing the supplemented fats, unsaturated fats had a bigger suppressing effect on CH<sub>4</sub> output compared to other FA (Patra 2013). The C18:3 had marked inhibitory effect, compared to other FA, on CH<sub>4</sub> production (Patra 2013). The background for the suppressing effect is explained by that unsaturated fats are saturated in rumen and act thus as a H<sub>2</sub> sink. However, the bio hydration does not lower the accumulated H<sub>2</sub> as efficient as the production of CH<sub>4</sub> (Czwerkawski et al, 1986 quoted in Johnsson & Johnsson 1995). Van Zijderveld (2011) concludes that saturation of dietary fat of 20 % result in a CH<sub>4</sub> decrease of 1.6 % output (Van Zijderveld et al., 2011). Bacterial type in cow's rumen can influence the fats depressing effects. Gram positive methanogens was shown to be affected by fat supplementation, although no gram-negative methanogens were affected (Galbraith et al., 1971).

The underlying mechanism of lipids decreasing effects on CH<sub>4</sub> output is versatile and not yet fully understood. Main reasons found in literature are CH<sub>4</sub> inhibition through enhanced propionic acid production (Machmyller 2013; Patra 2013), bio hydration of unsaturated FA as an alternative H<sub>2</sub> sink (Czwerkawski et. al, 1966), inhibition of protozoal production (Johnson & Johnson 1995), reduced fibre digestion, and fat to be toxic to

methanogens (Harfoot et al, 1974; review by Topak 2016). It is through the attachment of fat on the fibre and the methanogens that disfavour the methanogens (Machmüller et al., 2003). This is due to that the fat on the membrane of the methanogenic bacteria prevents nutrients to enter. However fat fed in form of free fatty acids (FFA) instead of triacyl glycol have shown to be important for the fat to be able to attach to bacteria (Harfoot et al., 1974)

### 2.3.2 Linseeds

Linseed *Linum usitatissimum* contains a high part of unsaturated FA (table 1) were 18:3n-3 represents more than 50% of total FA (Chilliard et. al, 2007; table 1). The ratio between omega 3 and omega 6 in linseed fat is approximately 3.6:1 (table 1). Linseed used in experiments differ in form; whole linseed, crushed linseed, linseed oil or extruded linseed (EL). General process of extrusion of linseed are done through short period of cooking followed by crushing of seeds under heat and pressure and quickly heating the crushed linseeds. In this way the content of cyanide is lowered (Martin et al., 2008).

A feeding *in vivo* study where linseed was supplemented to the feed showed a decreased CH<sub>4</sub> output/day. Depending on how the linseeds was processed a reduction of CH<sub>4</sub> output was observed varying from -12 % with crude linseed, -38 % with extruded linseed and -64 % with linseed oil (Martin et. al, 2008). Data expressed as CH<sub>4</sub> per fat corrected milk (FCM) showed a decrease of 23 % due to EL. However, a decrease in DMI (-3.1kg/day) and NDF intake (-6.8 % on average) due to EL was also found in that study (Martin et al. 2008).

In a study from 2011, methane output was determined in 40 cows kept in respiration chamber for five days. No effects of extruded linseed were found compared to control diet. Also, no effect was found in energy balance, digestibility of fibre (NDF) compared to control diet. It was concluded that the importance of confirmation of *in vitro* data with *in vivo* data (Van Zijderveld et al, 2011). In other studies, dietary fat (sunflower oil) was found to decrease digestion of NDF (McGinn et al., 2004).

**Table1.** Fatty acid composition (g/100g fatty acids) of some common fats and oils. Fatty acid composition of experimental feed Easylin 100/30 (Modified table from McDonald, 2011; Agrosom, 2015).

| Fatty acid | Rapeseed | Soya bean | Red clover | Ryegrass | Linseed | Butterfat | Easylin 100/30 |
|------------|----------|-----------|------------|----------|---------|-----------|----------------|
| 16:0       | 4        | 10        | 15.4       | 12       | 6       | 31        | 4              |
| 18:0       | 1        | 4         | 2.3        | 2        | 3       | 10        | 3.7            |
| 20:0       | 1        | Tr        | -          | -        | -       | -         | -              |
| 22:0       | Tr       | Tr        | -          | -        | -       | -         | -              |

|         |    |    |      |    |    |    |    |
|---------|----|----|------|----|----|----|----|
| 16:1    | 2  | Tr | 0.1  | 2  | -  | 2  | -  |
| 18:1n-9 | 54 | 25 | -    | 15 | 17 | 23 | 18 |
| 18:2n-6 | 23 | 54 | 20.8 | 68 | 13 | 2  | 16 |
| 18:3n-3 | 10 | 7  | 59.5 | -  | 55 | <1 | 58 |

## 2.4. Effects of linseed feed ration on milk yield and fat composition

### 2.4.1 Milk yield

There are conflicting results in literature of the effect of linseeds on milk yield. Supplementary fat was found to increase milk yield (Patra 2013), linseed oil and extruded linseeds (EL) have been found to increase kg milk yield per day (Bu et al. 2007; Zaucht et al., 2010). Other results show a decrease in the milk yield (Chilliard et al 2009; Gunthier et al 2005) Other studies show no effect on milk yield (Beuchermin 2009).


In an analysis where many studies with different fat supplements were used, an increase of milk yield was found with increasing fat supplement. However, a decrease of milk yield was found with fat supplements over 3.9-6 % (Amlan K. Patra, 2013). In a study where 5.7 % fat of dietary DM was used a drop of 20 % was found for cows feed EL compared to control group (Martin et al. 2008).

## 2.5 Milk fat composition

### 2.5.1 Lipids

Lipids have a crucial role in the body of animals. Lipids are components in the cell membrane, absorber of some vitamins, energy storage and are high in energy and contribute with energy for the animal. Lipids are divided into 4 main groups: FA, glycerides, nonglyceride lipids and complex lipids (Denniston et al., 2010).

Main constituents of the milk fat fraction are triacyl glycols (figure 3), where glycerol bind three FA (McDonald et al 2011). There are more than 500 different FA in milk (Patton & Jensen 1975).



**Figure 3.** Modified picture from Harvey et al., 2011. A triacyl glycol with single and one double bound.

Short-chain FA consist of a carbon chain of 4 to 8 carbons, medium-chain FA consist of 10 to 14 carbons and long-chain FA contains 16 or more carbons (Bauman & Griinari, 2003). The length and saturation have an impact on the melting point of the FA. It increases with increased number of carbon as well as with saturation level (Deniston et al.). Milk contains mainly medium-chain FA with parts of 16 carbons and the milk FA are foremost saturated. Part of C18:3 in cowmilk is 0.2 % (Lindmark & Mårtensson, 2012).

Linoleic acid, C 18:2, and  $\alpha$ -linolenic acid, C 18:3, are essential since they cannot be synthesized by the animal itself (Harvey & Ferrier, 2011). Instead these FA need to be ingested with the feed the feed. Linseed based feed consists of a high part of essential FA as well as grass (table 1).

### 2.5.2 Dietary fat effect on milk fat composition

Lipids consumed by the cow differ in structure depending of feed source. Lipids in plants often consist of glycolipids where fat in grass have high amounts of polyunsaturated FA (PUFA) (Dewhurst et al 2003). Fat in cell membrane consists of phospholipids and fat stored in the seed consists of triglycerides. The fats in oil seed plants also consists of PUFA. Lipids in dietary fat are degraded by ruminal microbes in rumen through hydrolysis. Freed FA high in PUFA in rumen are bio hydrogenated by ruminal microorganisms where hydrogen is added to the PUFA in a saturation process (McDonald et al, 2011). Almost all linoleic acid (C18:2) and linolenic acid (C18:3) are saturated and converted into stearic acid (C18:0) in rumen (rewired by Doreau & Ferlay 1994). However not all FA is completely saturated but intermediates in form of mono unsaturated FA (MUFA) and PUFA are created. The saturated and unsaturated FA are passed on to duodenum where it is extracted to the blood. The fat is transported in chylomicrons and very-low-density lipoproteins (VLDL) (Chilliard et. al, 2007) in the blood to the udder where it is used for milk fat synthesis (Gonthier et al, 2005).

The main part of VFA is absorbed across the ruminal epithelium and released into the portal blood. Propionate is then converted to glucose in the liver. The main fraction of butyrate is metabolised to the ketone body  $\beta$ -hydroxybutyrate in the ruminal epithelium and released into the portal blood (Sjaastad et al, 2010). Acetate and  $\beta$ -hydroxybutyrate are precursors for *de novo* synthesis of milk fat in the mammary gland. Roughly half of milk FA are produced *de novo* and acetate and  $\beta$ -hydroxybutyrate are the principal precursors. However, the relative contribution from the *de novo* synthesis is generally lower in the early stage of lactation when the cow mobilises long FA from the adipose tissue. The other 50 % of milk fat is produced from preformed FA. Mobilised FA also contributes to the preformed FA. The FA are, in the alveolar cell, esterified into triacyl glycol and transported into the milk by fat globuli

(Sjaastad et al, 2010). Short and medium-chain FA originates from *de novo* synthesis and long chain FA originates from preformed FA. FA with C 16 derives from both *de novo* and preformed FA (Bauman & Griinari, 2003; Lindmark Månsson et al. 2006).

Linseed has, in earlier studies, shown to have an impact on milk fat composition. A diet with linseed increases the unsaturated FA (UFA), C18:3, in the milk fat and decreases the saturated FA (SFA), C16:0, acids in the milk (Zachut et al 2010; Chilliard et al., 2009). At the same time negative correlations ( $r=-0.86$  to  $-0.90$ ) have been found between increased CH<sub>4</sub> output and elevated levels of unsaturated FA in milk fat (Chilliard et. al, 2009). The elevation of UFA in milk fat is explained by literature as an increased amount of FA passing to the small intestine. The underlying mechanism is the higher uptake of UFA from linseeds in the small intestine (Chilliard et al., 2009). Supporting results was found in a study where EL in diet resulted in decreased levels of medium chain FA and increased levels of long chain FA in plasma (Gonthier et al 2005). However, the milk fat increase of C18:3 was found to be negatively correlated with milk fat percentage and milk yield (Zachut et al., 2010). A risk with diets supplemented with PUFA is a low milkfat syndrome where a heavy decrease of milk fat can be seen. Highest drop in milk fat depression is seen with diets low in fibre and with supplemented PUFA (Griinari et al., 1997).

## 2.6 How is methane measured from cows?

### 2.6.1 Methane measurements

CH<sub>4</sub> output is represented as a mass of gas that is released by the cow per unit of time. The released amount of gas varies in time depending on the cows belching. This means that the measuring device should be able to measure both factors (Larios et al., 2016).

CH<sub>4</sub> output from cows can be measured in different ways, both *in vivo* and *in vitro*. The most accurate *in vivo* technique for CH<sub>4</sub> output “the golden standard” is the respiration chamber. The animal is placed in a closed chamber where exhaust air with its gas is sampled with a decided time cycle (Changunda et al., 2013; Ramin & Huhtanen, 2014; Hristov et al., 2015). This method is often used for evaluation of other CH<sub>4</sub> measuring methods (Johnsson & Johnsson 1995; Danielsson et al., 2017).

A widely used technique for measuring methane is the tracer gas technique with sulphur hexafluoride. A permeation tube with sulphur hexafluoride is placed in cows rumen releasing the gas successively with a known releasing rate. Greenhouse gases and sulphur hexafluoride exhaled from the cow is sucked into hoses and transported into a stainless-steel collection sphere. The ratio between sulphur hexabromide and CH<sub>4</sub> is determined by a gas chromatograph. Before the

measurements begins the animals need to be trained to wear the sampling equipment on the head. The technique enables measurement on a large number of animals on farm as well as many times/day/animal (Hristov et al., 2015; Boadi et al., 2001).

Other *in vivo* measuring techniques are the GreenFeed-system which is based on an automatic head-chamber system measuring the exhaust air from the cow's muzzle. Here the individual cow is sampled for CH<sub>4</sub> and CO<sub>2</sub>, during feeding in a portal feed trough. Measurements are done by infra-red sensor and optionally with tracer gas CH<sub>4</sub> ratio. The sampling is done during a period of days or weeks resulting in a calculation of CH<sub>4</sub> output per day. Due to many measurements per day this method catches the fluctuations of methane production over the day as well as numerous animals can be measured measurement compared to the respiration chamber technique. Measurement of background greenhouse gas can also be made. Tag reader register the animal and data of that specific animal, ex. milk yield, are transferred to the analysing station. Before the measuring begins the animals need to be trained to eat from the portal feed trough (Hristov et al., 2015; Zimmerman et al., 2012).

There is also *In vitro* methods were CH<sub>4</sub> is measured from containers of feed during oxygen free environment. Indirect estimations of CH<sub>4</sub> exist were correlations with DMI is used for calculations of CH<sub>4</sub> output (Ramin & Huhtanen, 2012 & 2013 ).

CH<sub>4</sub> output can also be estimated by laser methane detection technique (LMD) which is used in this study. The technique was developed for measurement of CH<sub>4</sub> leaks from gas pipes and started to be used in trials of CH<sub>4</sub> measurements of cows by Changunda et al. in 2009. The LMD measures the CH<sub>4</sub> in the exhaust air from the cow with its laser diode. The method relies on the fact that CH<sub>4</sub> absorbs a specific wavelength of infrared rays. The LMD measures the diffused light back and measures the absorption of the light. The LMD measures the average CH<sub>4</sub> gas density between the detector and the target. One measurement is made every 0.5 second over a duration of a few minutes.

CH<sub>4</sub> output is reported as the product of concentration of CH<sub>4</sub> cloud (measured in parts per million) and path length (measured in meters). Best application is outdoors in strong sunshine. The LMD calibrates itself automatically each time it is turned on (Crowcon production voucher, 2013). With this type of measurement it is possible to measure many cows on farm. The accuracy is however lower compared to respiration chamber which was found in a study of Ricci et al., 2014. In the study, measurements were done on 72 steers, after feeding, with one 4-minute measurement/ animal /day repeated over 3 days. The week after the CH<sub>4</sub> output was measured in a respiration chamber. The CH<sub>4</sub> sample/cow was corrected for background CH<sub>4</sub> measurement where data lower than the lowest measurement was subtracted from then deleted. There was a

strong linear relationship ( $R^2=0.86$ ) between LDM measurement and the respiration chamber (Ricci et al., 2014) and the relationship of  $r=0.47$  ( $p=0.001$ ) for dairy cows was found by Changunda et al., 2013.

### 2.6.2 Influencing factors of methane measuring

CH<sub>4</sub> output was found to be affected by lactation status (lactating or dry cows) and the cow's activity. Depending on the activity of the cow, the pattern of eructation differs. Cows have the highest CH<sub>4</sub> output during rumination, lower CH<sub>4</sub> output during feeding and lowest while staying idle. These facts can influence the results of CH<sub>4</sub> measurements if not corrected for in the experiment (Changunda, 2009).

Time for inurement of cows on the new diet can affect the measurements. In an earlier study an 8 days' time of inurement of linolenic acid in diet appears to be required for the CH<sub>4</sub> measurements to stabilise. A 12 day of re-establish initial CH<sub>4</sub> values was needed for the cows after finished test diet (Zwerkawski, 1966).

## 3. Materials and Methods

This study was initiated by the feed company Agrosom and was executed as part of an EU-project named "Rolling out of an innovative enteric methane emissions measurement method for cattle" (ECO-methane). The project was performed within the EU-programme "Competitive and innovation frame work programme" (CIP). The design of this project is almost similar to that of the ECO-methane project (ECO-methane, 2014) however there were some modifications of the material and method of this work. These are mentioned under measurements and sampling.

### 3.1 Experimental design and herds

The study was executed on five conventional dairy herds in Sweden. Three of the participating herds were located in Kalmar län, one in Halland and one in Uppland. Descriptive data of participating herds are presented in table 2.

The study was divided into three periods (Per) where Per 1 was control, Per 2 was treatment, and Per 3 was control. This is also called a switch-back design. During Per 1 all cows were fed control diets, equal to the feeding strategy on the farm. During Per 2 the diets were supplemented with the test feed EL. The EL was fed as an ingredient of commercial compound feed in the ration (table 2 and 3). The goal was to feed each cow 1000 grams of EL per day. Per 3 was a second control period where EL was excluded from the feed ratios and cows returned to their normal feed ratios.

Table 2. Descriptive data of participating herds. PMR=partly mixed ration. TMR= total mixed ration.

| Herd: | Breed                        | Milking system  | Feeding system                           | Feeding system for Extruded Linseed |
|-------|------------------------------|-----------------|--|-------------------------------------|
| 1     | Holstein                     | AMS             | PMR, -feeding wagon                      | Concentrate in AMS                  |
| 2     | SRB and Holstein             | AMS             | -PMR, transport band. -Feeding stations. | Concentrate in feeding stations     |
| 3     | SRB                          | Milking parlour | TMR two different mixes                  | Concentrate in mixing wagon         |
| 4     | SRB, Holstein and crossbreed | AMS             | PMR -feeding wagon                       | Non pelleted in Mixing wagon        |
| 5     | Holstein                     | AMS             | -PMR -feeding wagon. -feeding stations   | Concentrate in feeding station      |

### 3.1.1 Treatment feed

The treatment feed containing linseed, Easylin 100/30, was produced in Bremen, Germany. Easylin 100/30 contained proximately 60 % extruded linseeds and 40% rape seed meal. Amount of crude fat in Easylin was 277 g per kg dm. Total fatty acids in Easylin 100/30 was 887 g/kg crude fat. The amount of linolenic acid was 53.2 g/100g (Agrosom, 2016; NorFor, 2018).

The feed rations at the herds were not corrected for different DMI, solubility of feed, parts of carbohydrate, silage process or amount of fat intake.

### 3.1.2 Selected cows

A test group of 20 cows was created for each herd and period. Criteria for selection for the cows were; days in milk (DIM) within 30-180 days, a health record for at least 2 months back, 25% of the cows was primiparous and 75% were multiparous. These criteria were in accordance with the ECO-methane project (2014). The final selection of the 20 individuals was done at the farms. A list of all potential cows from the herd within the qualifications was created. The first 20 cows from the list that we found in the feeding area was selected to be measured for CH<sub>4</sub> and included in the test group.

## 3.2 Measurements and sampling

The measurements started in December 2014 and ended in May 2015. None of the cows were on pasture during the trial. The following measurements were done once in every period at every herd as in ECO-project; one direct CH<sub>4</sub> measurement of LMD on the selected cows, one questionnaire per herd directed to the farmer, collection of data from milk sampling and of information of sick and treated cows.

Additional measurements outside the ECO-project sampling, only for this master thesis, were; one-day feeding control per herd and period. For that I collected data on feed consumption and milk yield of all lactating cows for the day we visited the herd and the number of cows in the lactating group. The amount of consumed feed took into account the amount of roughage and concentrate fed and leftover feed from feeding bunker. The leftovers were weighed or estimated by the farmer. Roughage feed sample were taken at the herds the same day as I visited the herd roughage for dry matter measurements. Other analyses on nutrient composition of the feeds were collected from the farmers' own feed analyses. The one-day feeding control was calculated in the computer tool IndividRAM (Växa Sverige, Eskilstuna, Sweden) which is based on the feed evaluation system NorFor (Volden, 2011).

### 3.2.1 Analyses on feed and milk

Roughage feed samples were analysed for DM (at 60 °C) at the SLU feed laboratory (Uppsala, Sweden). This sample was used for the one-day feeding control on herd level. The milk samples, collected monthly according to the official Swedish milking recording scheme, were also used for analyses for FA composition. The farmer performed the milk recording for the herd during normal procedures and labelled the sample tubes with cow number for the selected cows (same cows that was measured for CH<sub>4</sub>). The labelled tubes were singled out at the Eurofins Steins laboratory (Jönköping, Sweden) and milk samples were analysed on FA with a Fourier Transform Infra-red (FTIR) instrument. Individual milk samples were analysed for the FA: C16:0, C18:0, C18:1, MUFA, PUFA, sum of SFA and sum of UFA, sum of all FA (FA<sub>tot</sub>) and protein (FA<sub>pr</sub>). Deviation for the FTIR method vary for SFA from -0.07 % to + 0.07 % and for UFA from -0.14 % to +0.12 % (Kaylegian et al., 2009).


Data on milk yield (MY), energy corrected milk (ECM), fat content (MF) and protein content (MP) was collected both on individual cows and on herd level in each period. These data were provided by Växa Sverige through the Swedish milking recording scheme (Eskilstuna Sverige).

The milk fat and protein were analysed by the CombiFoss 6000 equipment (Eurofins Steins, Jönköping, Sweden). According to Nils-Erik Larsson (personal communication), corrections was done for milk fat and protein

content using standard procedures. The corrections were done for herds with milk sampling once per day for the variables; breed, parity, DIM and ratio between the milk recording for the sampling and the milk recording of the complete day. For herds where milk recording, and sampling was done twice per day no corrections were done.

### 3.2.2 Methane measurement

Direct CH<sub>4</sub> measurement was done with LMD from the trademark Crowcon, production name TGE laser methane mini-Green (LMm-G) (Crowcon production voucher, 2013). The LMD green laser point was focused on the cow's nose during the measurement. Each individual measurement lasted between 3 and 4 minutes resulting in a minimum of 360 individual measurements per cow where one new measurement was done every 0.5 second (figure 3). The LMD measurements was performed by a well-trained person from France.



**Figure 3.** An example of a CH<sub>4</sub> measurement from one individual cow. The x-axis shows the total of 499 measuring points during 250 seconds. The y-axis shows CH<sub>4</sub> output in ppm, when a laser methane detector is pointed towards a cow's nostrils.

The person was standing on the feeding bunker at approximately two meters distance from the cow while he directed the LMD device towards the cow's nostrils. CH<sub>4</sub> output was measured while the cow was eating. Methane measurements of concentration was specified as parts per million. Each methane measurement lasted for 180 seconds and a total of 360 methane measurements was made under this time for each cow. A mean of these 360 measurements was calculated for each cow. The individual measured values were then divided by the individual ECM, from milk recording, creating an individual momentary methane output ppm per kg ECM.

### 3.2.3. Indirect methane estimations with Visiolait

Indirect CH<sub>4</sub> estimation was done on herd level by a feeding ratio tool called Visiolait which estimated the daily CH<sub>4</sub> output in grams per litre of milk of each herd. A patented equation (Chesneau, 2008) was used for the estimate. The equation included information of FA profile, amount of milk yield from tank milk kg per cow and day and DIM of herd. The CH<sub>4</sub> values were used for comparison through correlations with the direct CH<sub>4</sub> measurement.

### 3.2.4 Indirect methane estimations by NorFor

Indirect CH<sub>4</sub> estimations on herd level were done by NorFor based one-day feeding control calculations. The CH<sub>4</sub> value was estimated on herd level by grams per cow and day. The intention was to create correlations between NorFor estimations and mean of direct CH<sub>4</sub> LMD measurement per kg ECM and CH<sub>4</sub> estimation of Visiolait.

$$CH_4 = \frac{1.39 * DMI - 0.091 * FA}{55.65} * 1000$$

**Figure 4.** Estimation of CH<sub>4</sub> for dairy cows where CH<sub>4</sub>=methane g per day, DMI=total feed intake kg DM per day, FA=fatty acid in the diet grams per kg DM, number 55.65 converts CH<sub>4</sub> energy MJ to gram (Nielsen et al. 2013).

## 3.3 Statistical calculations

An average of DIM was calculated for selected cows in the test groups on each farm and period. Collected data for these cows was used in statistical calculations in Minitab 17.0. Corrections for DIM, parity, and health and activity were included in the experimental design. For correction of data see statistical calculations further down.

The statistical model ANOVA: General Linear model (GLM) was used to calculate the statistical difference between herds (Herd) and between periods (Per) for the variables DIM, ECM, MY, MF, MP, FAT<sub>tot</sub>, FA<sub>pr</sub>, C16:0, and C18:0, C18:1, MUFA, PUFA, SFA, UFA, CH<sub>4</sub> of LMD, Ch<sub>4</sub> of LMD per kg ECM and amount of EL (table 6). Variables that potentially could have a significant effect was included in the calculation. If there was no significant effect, the variable was withdrawn from the calculation, such a variable was Herd. Herd and Per were fixed in the model. A pairwise comparison with a confidential level of 95% was used to locate the significant difference between specific periods (table 6). Means of variables were also generated through the pairwise comparisons with confidential level of 95%. The mean values were then used to create bar graphs, in Excel, for each variable. Descriptive charts for DIM, EL and number of cows was done in Minitab or excel. No correlations lower than 0.5 are reported in this paper unless required for special reasons.


For the non-significant Herd values, a new GLM was done where Herd was erased from the model.

Correlation between LMD measurements and CH<sub>4</sub> from Visiolait and CH<sub>4</sub> from NorFor was done.

## 4. Results

### 4.1 Descriptive data

Number of cows in the testgroups (figure 5), DIM, daily DMI did not differ significantly between periods ( $p=0.34$ ) or between herds (exception of herd 5 with lower daily DMI compared to the other herds) (table 6; table 3).



**Figure 5.** Number of cows at each herd and for every period in the final test group.

Ingestion of extruded linseed differed between herds. However, no statistic calculation of this was made (table 3). Results from one day feeding control are shown in tables 3 and 4.

**Table 3.** Average daily feed intake per cow on herd level and chemical composition of diets in the five herds. Amounts of feed and composition were provided from the herdsman. Amount of linseed was calculated from the amount of Easylin 100/30 supplied in the diets during period 2.

|                                | Period 1 | Period 2 | Period 3 |
|--------------------------------|----------|----------|----------|
| <b>Herd 1</b>                  |          |          |          |
| No. of cows in test group      | 13       | 18       | 12       |
| No. of lactating cows          | 517      | 519      | 522      |
| Total dry matter intake, kg DM | 22.4     | 23.0     | 21.4     |
| <b>Herd 2</b>                  |          |          |          |
| No. of cows in test group      | 16       | 14       | 16       |

|                                |      |      |      |
|--------------------------------|------|------|------|
| No. of lactating cows          | 138  | 137  | 136  |
| Total dry matter intake, kg DM | 23.6 | 20.8 | 18.8 |
| Roughage intake, kg DM         | 12.6 | 11.3 | 8.8  |
| Concentrate intake, kg DM      | 11.0 | 9.5  | 10.0 |
| Expelled linseed, g feed       | 0    | 1065 | 0    |
| NDF, g/kg DM                   | 353  | 340  | 332  |
| Crude protein, g/kg DM         | 172  | 177  | 189  |
| Crude fat, g/kg DM             | 46   | 39   | 48   |
| <b>Herd 3</b>                  |      |      |      |
| No. of cows in test group      | 19   | 11   | 5    |
| No. of lactating cows          | 188  | 185  | 191  |
| Total dry matter intake, kg DM | 27.4 | 29.6 | 27.8 |
| Roughage intake, kg DM         | 12.1 | 13.6 | 11.5 |
| Concentrate intake, kg DM      | 15.3 | 16.0 | 16.3 |
| Expelled linseed, g feed       | 0    | 924  | 0    |
| NDF, g/kg DM                   | 389  | 328  | 360  |
| Crude protein, g/kg DM         | 197  | 185  | 182  |
| Crude fat, g/kg DM             | 58   | 55   | 53   |
| <b>Herd 4</b>                  |      |      |      |
| No. of cows in test group      | 18   | 11   | -    |
| No. of lactating cows          | 138  | 135  | -    |
| Total dry matter intake, kg DM | 21.5 | 25.3 |      |
| Roughage intake, kg DM         | 14.1 | 11.4 |      |
| Concentrate intake, kg DM      | 7.4  | 13.9 |      |
| Expelled linseed, g feed       | 0    | 1000 | 0    |
| NDF, g/kg DM                   | 286  | 315  |      |
| Crude protein, g/kg DM         | 195  | 184  |      |
| Crude fat, g/kg DM             | 47   | 52   |      |
| <b>Herd 5</b>                  |      |      |      |
| No. of cows in test group      |      | 13   | 11   |
| No. of lactating cows          | -    | 375  | 375  |
| Total dry matter intake, kg DM |      | 23.8 | 28.1 |
| Roughage intake, kg DM         |      | 13.9 | 18.6 |
| Concentrate intake, kg DM      |      | 9.9  | 9.6  |
| Expelled linseed, g feed       | 0    | 1126 | 0    |
| NDF, g/kg DM                   |      | 347  | 343  |
| Crude protein, g/kg DM         |      | 190  | 188  |
| Crude fat, g/kg DM             |      | 44   | 42   |

**Table 4.** Milk yield and milk composition of all lactating cows of the five herds. Data was provided by Växa Sverige

|                                      | Period 1 | Period 2 | Period 3 |
|--------------------------------------|----------|----------|----------|
| <b>Herd 1</b>                        |          |          |          |
| No. of cows                          | 597      | 613      | 615      |
| Days in milk                         | 150      | 206      | 234      |
| Day of sampling from previous period | -        | 35       | 14       |
| Milk yield, kg                       | 25.3     | 24.5     | 24.7     |
| Energy corrected milk, kg            | 25       | 24.3     | 24.3     |

|                                      |      |      |        |
|--------------------------------------|------|------|--------|
| Milk fat, %                          | 3.8  | 3.9  | 3.9    |
| Milk protein, %                      | 3.5  | 3.5  | 3.4    |
| <b>Herd 2</b>                        |      |      |        |
| No. of cows                          | 138  | 138  | 137    |
| Days in milk                         | 127  | 223  | 244    |
| Day of sampling from previous period | -    | 70   | 3      |
| Milk yield, kg                       | 29.9 | 29.6 | 31.2   |
| Energy corrected milk, kg            | 30.5 | 31.0 | 32.1   |
| Milk fat, %                          | 4.1  | 4.2  | 4.1    |
| Milk protein, %                      | 3.6  | 3.7  | 3.6    |
| <b>Herd 3</b>                        |      |      |        |
| No. of cows                          | 194  | 185  | 187    |
| Days in milk                         | 186  | 249  | 222    |
| Day of sampling from previous period | -    | 49   | 7      |
| Milk yield, kg                       | 26.1 | 27.2 | (26.0) |
| Energy corrected milk, kg            | 27.8 | 28.6 | (27.7) |
| Milk fat, %                          | 4.3  | 4.3  | (4.1)  |
| Milk protein, %                      | 3.7  | 3.6  | (3.6)  |
| <b>Herd 4</b>                        |      |      |        |
| No. of cows                          | 135  | 130  | -      |
| Days in milk                         | 228  | 255  | -      |
| Day of sampling from previous period | -    | 24   | -      |
| Milk yield, kg                       | 30.7 | 32.0 | -      |
| Energy corrected milk, kg            | 30.8 | 31.7 | -      |
| Milk fat, %                          | 4.0  | 4.0  | -      |
| Milk protein, %                      | 3.4  | 3.4  | -      |
| <b>Herd 5</b>                        |      |      |        |
| No. of cows                          | -    | 375  | 413    |
| Days in milk                         | -    | 208  | 248    |
| Day of sampling from previous period | -    | 28   | 24     |
| Milk yield, kg                       | -    | 31.5 | 29.4   |
| Energy corrected milk, kg            | -    | 32.3 | 30.6   |
| Milk fat, %                          | -    | 4.1  | 4.3    |
| Milk protein, %                      | -    | 3.6  | 3.5    |

## 4.2 Milk yield and milk composition

### 4.2.1 Milk yield


Calculations based on data from cows in test groups showed that Milk Yield ( $P=0.55$ ) and ECM ( $P=0.24$ ) did not differ significantly between treatment period and control periods (table 6).

### 4.2.2 Protein

Protein content in milk, MP, increased significantly ( $P=0.006$ ) from 3.4 in Per 1 to 3.5 in Per 2. Protein, FApr, did not increase significantly.


### 4.2.3 Unsaturated fatty acids in milk

PUFA, increased significantly ( $P < 0.0001$ ) from period 1, 0.13g per 100g, to period 2, 0.15g per 100g, and decreased mean numerically from period 2 to period 3, 0.13g per 100g (Figure 6; table 6).



**Figure 6.** Mean and their SE of PUFA for period1, 2 and 3.

Herd 1 had significantly ( $P = 0.008$ ), higher mean of unsaturated acids compared to herd 3 and 4. There was a significant ( $P = 0.005$ ) decrease of mean of UFA from period 2, 1.15 g per 100g, to period 3, 1.03 g per 100g. There was a numerical increase of unsaturated acids from Per 1 to Per 2, however not significant (Figure 7; table 6).



**Figure 7.** Mean and SE Mean of UFA for period.

Oleic acid, C 18:1 ( $P = 0.36$ ) and MUFA ( $P = 0.83$ ) did not differ significantly between periods.

#### 4.2.4 Saturated milk fatty acids

Saturated FA did not differ significantly between periods ( $P = 0.64$ ) or between herds ( $P = 0.28$ ) (table 6). Palmitic acid, C16:0, tended to decrease ( $P = 0.055$ ) from period 1 to period 2. A small numerically increase was seen in period 3 (figure 8; table 3).




Figure 8. Mean and SE Mean for palmitic acid (C16:0) in periods.

### 4.3 Methane output




Figure 9. Boxplot of CH<sub>4</sub> measurements by LMD on individual cows (ppm per kg ECM) for all herds (Bes) and periods (Per). \* and \*\* indicate significance level on Per within Bes (\* P<0.05; \*\* P<0.01)

Mean of momentary CH<sub>4</sub> ppm per daily ECM yield differed significantly (P< 0.0001) between period 1 and 2, between period 1 and 3. Mean of CH<sub>4</sub> per ECM decreased significantly (p<0.001) from Period 1, 4.9, to period 2, 3.8 which corresponds to a reduction of 22 %. Mean of CH<sub>4</sub> per ECM increased, however only numerically, from Per 2 to Per 3, 3.8. (Figure 10).




Figure 10. Mean and SE Mean of CH<sub>4</sub> ppm per kg ECM for periods.

### 4.3.1 Estimated methane value through fatty acid analyses, Visiolait

There was no significant ( $p=0.16$ ) relationship ( $r= 0.46$ ) between LMD measured CH<sub>4</sub> ppm production and Visiolait estimated CH<sub>4</sub> production g per litre milk. Data from Per 1 and Per 2 was used (table 5).

### 4.3.2 Estimated methane value through NorFor

There was no significant ( $p= 0.53$ ) correlation ( $r= 0.53$ ) between measured CH<sub>4</sub> by LMD and the estimated CH<sub>4</sub>, production gram per kg ECM, done by NorFor one-day feed control. Data from Per 1 and Per 2 was used (Table 5).

**Table 5.** CH<sub>4</sub> data for the five herds. CH<sub>4</sub> measurement of laser methane detector (LMD) and CH<sub>4</sub> estimation by Visiolait (estimated through fatty acid composition) are based on cows in the test group, while CH<sub>4</sub> estimation by NorFor (calculated in a one-day feeding control) is based on all lactating cows.\* shows missing values

|                                  | Per1 | Per2 |
|----------------------------------|------|------|
| <b>Herd 1</b>                    |      |      |
| CH <sub>4</sub> LMD, ppm per ECM | 3.80 | 3.70 |
| Visiolait, g per litre milk      | *    | 15.0 |
| NorFor g per kg ECM              | 505  | 520  |
| <b>Herd 2</b>                    |      |      |
| CH <sub>4</sub> ppm per ECM      | 7.17 | 3.41 |
| Visiolait, g per litre milk      | 16.3 | 16.0 |
| NorFor g per kg ECM              | 535  | 476  |
| <b>Herd3</b>                     |      |      |
| CH <sub>4</sub> ppm per ECM      | 4.82 | 3.99 |
| Visiolait g per litre milk       | 18.3 | 16.1 |
| NorFor g per kg ECM              | 613  | 674  |
| <b>Herd 4</b>                    |      |      |
| CH <sub>4</sub> ppm per ECM      | 3.69 | 3.30 |
| Visiolait g per litre milk       | *    | 15.3 |
| NorFor g per kg ECM              | 487  | 568  |
| <b>Herd 5</b>                    |      |      |
| CH <sub>4</sub> ppm per ECM      | *    | 4.53 |

|                            |   |     |
|----------------------------|---|-----|
| Visiolait g per litre milk | * | 16  |
| NorFor g per kg ECM        | * | 549 |

**Table 6.** Effects of EL on milk yield, milk composition and CH<sub>4</sub> output for herd and Period (Per). Mean from Basic statistics. LSmean for Per 1, 2 and 3 is created from individual data from the same period. When no p-value under herd, the variable herd was not used in the statistical calculation due to that it was not significant.

| Variable                      | Periods |      |      | P-value |       | Significance level |         |         |
|-------------------------------|---------|------|------|---------|-------|--------------------|---------|---------|
|                               | 1       | 2    | 3    | Herd    | Per   | Per 1-2            | Per 2-3 | Per 1-3 |
| DIM days                      | 103     | 99   | 100  | 0.000   | 0.993 |                    |         |         |
| MY <sub>1</sub> kg per day    | 37      | 39   | 39   | 0.001   | 0.553 |                    |         |         |
| MF <sub>1</sub> , %           | 3.9     | 3.9  | 3.9  | 0.001   | 0.657 |                    |         |         |
| MP <sub>1</sub> , %           | 3.4     | 3.5  | 3.4  | 0.000   | 0.006 | sig                |         |         |
| ECM <sub>1</sub> kg/d         | 36      | 39   | 39   | 0.018   | 0.237 |                    |         |         |
| FATot <sub>2</sub> g per 100g | 3.7     | 3.8  | 3.6  | -       | 0.363 |                    |         |         |
| FApr <sub>2</sub> g per 100g  | 3.3     | 3.4  | 3.3  | -       | 0.098 |                    |         |         |
| C16:0 <sub>2</sub> g per 100g | 1.1     | 1.0  | 1.0  | -       | 0.055 |                    |         |         |
| C18:0 <sub>2</sub> g per 100g | 0.37    | 0.40 | 0.35 | -       | 0.003 | sig                | sig     |         |
| C18:1 <sub>2</sub> g per 100g | 0.65    | 0.67 | 0.63 | 0.026   | 0.356 |                    |         |         |
| MUFA <sub>2</sub> g per 100g  | 0.96    | 0.98 | 0.96 | -       | 0.833 |                    |         |         |
| PUFA <sub>2</sub> g per 100g  | 0.13    | 0.15 | 0.13 | 0.004   | 0.000 | sig                |         |         |
| SFA <sub>2</sub> g per 100g   | 2.5     | 2.4  | 2.4  | -       | 0.641 |                    |         |         |
| UFA <sub>2</sub> g per 100g   | 1.0     | 1.2  | 1.0  | 0.008   | 0.005 |                    | sig     |         |
| LMD, ppm                      | 170     | 140  | 140  | 0.000   | 0.001 | sig                |         | Sig     |
| LMD, ppm per kg ECM           | 4.9     | 3.8  | 3.8  | 0.001   | 0.000 | sig                |         | Sig     |

Sig= significant. DIM=days in milk. MY= milk yield. MF= milk fat. MP= milk protein. ECM= energy corrected milk. FATot= Total fatty acids in milk. C16\_0= palmitic acid. C18\_0= saturated fatty acid with 18 carbons. C18\_1=unsaturated fatty acid with one double bound. MUFA= monounsaturated acids. PUFA= long chain polyunsaturated fatty acids. SFA=saturated fatty acids. UFA= unsaturated fatty acids. LMD= Laser Methane Detector. Measurements of methane output of concentration measured in parts of million. LMD/ECM= Mean CH<sub>4</sub> output per ECM. 3 Minutes= CH<sub>4</sub> measurement, duration of 3 minutes. 1=data from official Swedish milking recording scheme was distributed via VäxaSverige. 2= data from FTIR was distributed via Agrosom.

## 5. Discussion

### 5.1 Effect of Extruded linseed

A lowered DMI due to an intake of EL (Martin et al., 2008) was not found in this study. Due to that the quantity of EL used in this study, from 2.4 to 4.1 % of DMI per herd, to previous studies where a quantity of 7.9 % (Zachut, 2010), 12.6% (Gonthier, 2005) 15% (Chilliard, 2009) 16.6 % (Akrim, 2007) was used, the percentage of EL in this study can be considered low. The unchanged DMI in this study concludes that a quantity of EL up to 4.1 % will not affect the DMI.

#### 5.1.1 Milk Yield

EL did not increase MY or ECM in the present study. This is in accordance with other studies Chilliard *et al*, 2009; Martin *et al*, 2008; Petit Côrtes 2010. The hypothesis that EL increases MY could not be confirmed.

Chilliard et al. (2009) observed a decrease in MY due to a high amount of EL.

The amount of fed EL in the present experiment was lower than other studies, thus comparisons with other studies must be taken with caution.

#### 5.1.2 Milk fat

In this study EL had no significant effect on milk fat yield between periods (table 6). The result corresponds with former studies with feed linseeds (Gonthier *et al*, 2005; Beuchemin *et al* 2009). However other studies with EL in feed, showed a decreased fat yield in milk. This was thought to be a result of a high fat supplement of 5% of DMI that resulted in a lower digestibility of feed (Martin et al, 2008). This concludes that a quantity of up to 4.1 % of EL will not effect milk fat yield.

EL did increase content of PUFA in milk between Per 1 and per 2 (table 6). Due to the high deviation of measurement of UFA (Kaylegian et al., 2009), it must be emphasised that the FTIR method is not accurate to predict FA or groups of such acids with low concentrations. The concentration of PUFA in milk fat is generally less than 5%. Never the less the present result is in accordance with other studies (Petit & Côrtes, 2010; Gonthier et al, 2005; Zaucht et al., 2010) where infrared analyse was used for fat samples (Gonthier et al., 2005; Zaucht et al., 2010). Due to that both period and herd had a significant effect on PUFA, the treatment effect that is seen between periods can be due to the farm effect. EL also tended ( $P=0.055$ ) to decrease saturated FA in form of C 16:0 from Per 1 to Per 2 and increased numerically from Per 2 to Per 3 (figure 8). This is not in accordance with other studies where the decrease during EL-feed ratio have been significant. Also, the increase after finished treatment of C 16:0 has been significant in other studies. The fact that sampling was done close after transition from Per 2 to per 3 with the

result that the cows only consumed the control feed for a few days is probably the reason for the results.

### 5.1.3 Methane output

The result of the present study showing a decreased CH<sub>4</sub> output from Per 1 to Per 2 indicates that EL might have a lowering effect on CH<sub>4</sub> production. The result is in accordance with results of other studies of Martin et al. (2008) and Beauchemin et al. (2009). The reduction is difficult to compare with other studies of CH<sub>4</sub> output since the CH<sub>4</sub> measurements in this trial were only performed during a single activity. However, methane measurements of cows during feeding was made by Changunda 2013 with a result of 284 ppm. Also, a respiration chamber measurement of methane was done with the result of a mean of 356.3 ppm. Measurements of this study, 140 and 170 ppm, is comparatively low to Changundas measurements. Due to the big divergent levels in this study the data measurements cannot be considered to be accurate or be trustworthy. The reason for this difference could be environmental factors at the measuring occasion such as low light, wind draught or differences in measuring device. The small numeric increase of CH<sub>4</sub> from Per 2 to per 3 was not expected result and is not in accordance with other studies. The short time between the end of Per 2 and sampling in Per 3 (table 4 and 6) could explain a lingering treatment effect in Per 3. This is also in accordance with a study where the adaptation time for cows on new feed was 21 days (Gonthier, 2005). The reason for the short adaptation time in Per 3 was the upcoming grazing season. CH<sub>4</sub> level of LMD for farm two, period one is high, compared to the other farms and periods (table 5). The reason for this high value could be miss calculations or cows muzzels to close to each other during measurements.

The non-significant relationship between CH<sub>4</sub> measurement by LMD and Visiolait estimations, CH<sub>4</sub> and NorFor estimations could depend on different factors. The mean value of the CH<sub>4</sub> measurements, LMD, done in this study represents a “snapshot” of CH<sub>4</sub> output for the cows in the test group during the specific activity of eating and should not be mistaken for a value of CH<sub>4</sub> production per day. Visiolait is however an estimation of CH<sub>4</sub> g per litres of milk and NorFor is an estimation expressed in gram per kg ECM. These both estimations are made on herdlevel. Therefore, the CH<sub>4</sub> estimations are not directly comparable. Although despite the differences in units the CH<sub>4</sub> values should have a similar variation between the periods due to the treatment feed. The lack of a significant correlation could be the result of a non-correct CH<sub>4</sub> measurements or CH<sub>4</sub> estimations or both.

## 5.2 Deviations from experimental design

### 5.2.1 Herds and experimental design

Originally 7 herds had accepted to participate in the study however two herds decided to leave the project at an early stage.

### 5.2.2 Number of cows

The number of cows in the test groups (figure 5) was reduced due to mastitis and other illnesses. Individual cows were also excluded from the group since EL- feed was stopped for individual cows before measurement, incomplete FA analysis with some exceptions under divergent data.

The goal was to achieve similar average stage of lactation among cows representing the different herds, thus cows with too high or too low DIM was extracted from the test group. This resulted in smaller test groups than the 20 that was originally planned for. The results of this study should thus be considered with caution.

### 5.2.3 Sampling

Due to the specific release pattern of CH<sub>4</sub> from cows it is likely that the LMD- technique, with its frequent sampling every 0.5 second, can capture the levels of CH<sub>4</sub> output from cows.

The LMD is not the most accurate way of measuring CH<sub>4</sub> but is comparably cheap and is a user-friendly way to measure many cows in a herd. Due to that CH<sub>4</sub> measured with LMD is done once per individual cow in each period, could be considered less accurate than the Greenfeed method and tracer gas method which makes several measurements per cow and day. Also due to that LMD measurement are done in open air close to other individual cows and not within an enclosed space as with the Greenfeed method, it is likely that CH<sub>4</sub> output from other individual could affect the measurement. The calculated accuracies for LMD differ in studies. The risk of report wrong CH<sub>4</sub> measurement the individual cow is likely to be increased due to that the LMD had no ear tag number detector, compared to the Greenfeed method.

A correlation of  $r= 0.93$  between LMD and respiration chamber was found of Ricci et al 2014 which is likely to give trustworthy results with LMD. Changunda 2013 calculated a correlation to  $r= 0.47$ . Ricci 2014 deleted background data which could be the reason for the difference between the two results. Another potential difference between the studies could be the animal used were steers and not dairy cows. In this study no background data was deleted, and dairy cows was used similar to the study of Changunda 2013. Therefore, the data and design of data of this study is more alike the study of Changunda 2013.

The fact that not all CH<sub>4</sub> measurements (Herd 1 Per 2) took place in high light can also have affected the result. According to the producer of the LDM measurements should preferable be made in strong sunshine and

outside (Crowcon production voucher 2013). None of the measurements was done outside and therefore this should be taken in consideration when reading the result.

Sampling of CH<sub>4</sub> and milk was supposed to be done within the duration of approximate one week according to CIP project. In this project, time varied between milk recording and LMD measurements from 0 to 40 days. This extra time gap could have been accompanied with a differentiated environment like changed feed ratio.

Deviation from the experimental design in form of cows not eating during the LDM measurements was a fact in this experiment. Although the number of cows that was eating was predominant, the number of cows not eating varied between periods and between herds. This is a factor of uncertainty according to Changunda et al. (2009).

CH<sub>4</sub> measurements was supposed to be 4 minutes long but deviations from this happened and some measurements were only 3 minutes long. However, a measurement of 3 minutes is in accordance with other scientific CH<sub>4</sub> measurements (Changunda et al.,2009; Changunda 2013). Therefore, this deviation did probably not affect the reliability of the data.

#### 5.2.4 Feed

The level of feed EL for individuals in the herds could be affected by the feeding strategy of the farm. Herd 1, 2, 5 fed EL in feeding stations, the given amount of EL can be understood as consumed of the correct individual. However, no protection gates were integrated on the feeding stations and thus cows could have been disturbed and or backed out from the feeding station after given ration. Due to the human interference when EL feed are scooped into the mixer wagon it is likely that feeding stations are more accurate when distributing the EL feed/cow. Also, for herds with TMR (table 2) it is possible that some cows consumed more EL-feed compared to others due to hierarchy between cows at the feeding table.

Forage differed between the test periods at the farms. There effect of a changed forage on CH<sub>4</sub> and FA profile in this study is unknown.

#### 5.3 Future

Future questions to be answered is:

If and how much EL decreases the NDF degradation in rumen.

If and how a payment for reduced CH<sub>4</sub> output in relation to milk yield could be introduced, and what effect it could have on a lowered CH<sub>4</sub> output from herds. Today the payment for milk is based on milk yield, milk fat and protein (Arla, 2018).

If EL have a long-term effect of CH<sub>4</sub> output.

## 6. Conclusion

The results suggested that EL increases unsaturated fats in the milk and decreases the saturated fat. Results further suggest that EL does not increase the milk yield. The results do not show that EL decreases methane output. However, the method to measure methane emissions might have a limited accuracy.

## 7. Acknowledgements

Thank you to my supervisors for all support and help during this time. Thank You to Mr. Konrad Lichtin for the idea of this project. Thank You Mr. Mathieu Guillevic and Mr. Nicolas Forzy who did the CH<sub>4</sub> measurements on the farms.

## 8. References

- Agrosom 2016. Available at: <http://www.agrosom.de/>
- Arla (2018). Ägarbladet. [Broschyr]. Vecka 18–20. Maj 2018. Gunnar Gidefeldt. [2018-06-03].
- Bauman, D. & Griinari, J. (2003). Nutritional regulation of milk fat synthesis. *Annual Review of Nutrition*, vol. 23, pp. 203-227. doi: 10.1146/annurev.nutr.23.011702.073408
- Beauchemin, K.A., Kreuzer, M., O'Mara F. & McAllister, T.A. (2008). Nutritional management for enteric methane abatement: a review. *Australian Journal of Experimental Agriculture*, vol. 48, pp. 21-27.
- Beauchemin, K.A., McGinn, S., Benchaar, C. & Holtshausen, L. (2009). Crushed sunflower, flax, or canola seeds in lactating dairy cow diets: Effects on methane production, rumen fermentation, and milk production. *Journal of Dairy Science*, vol. 92, pp. 2118-2127.
- Boadi, D., Benchaar, C., Chiquette, J. & Massé, D. (2004) Mitigation strategies to reduce enteric methane emissions from dairy cows: Update review. *Canadian journal of animal science*. no. 830, pp. 319-335.
- Boadi, D.A. & Wittenberg, K.M. Methane production from dairy and beef heifers fed forages differing in nutrient density using the sulphur hexafluoride (SF6) tracer gas technique. *Canadian journal of animal science*, vol. 88, pp. 201-209.
- Bousquet, P., Ciais, P., Miller, J., Dlugokencky, E.J., Hauglustaine, D.A., Prigent, C., Van der Werf, G.R., Peylin, P., Brunke, E.-G., Carouge, C., Langenfelds, R. L., Lathière, J., Papa, F., Ramonet, M., Schmidt, M., Steele, L.P., Tyler S.C. & White J. (2006). Contribution of anthropogenic and natural sources to atmospheric methane variability. *Nature*, vol. 443 pp. 439-443. doi:10.1038/nature05132
- Bu, D.P., Wang, J.Q., Dhiman, T.R. & Liu, S.J. (2008). Effectiveness of Oils Rich in Linoleic and Linolenic Acids to Enhance Conjugated Linoleic Acid in Milk from Dairy Cows. *Journal of Dairy science*, vol. 90, pp. 998-1007.
- Changunda, M.G.G., Ross, D. & Roberts, D.J. (2009). On the use of a laser methane detector in cows. *Computers and Electronics in Agriculture*, vol. 68, pp 157-160.
- Changunda, M.G.G., Ross, D., Rooke, J., Yan, T., Douglas, J.-L., Poret, L., McEwan, N.R., Teeranavattanakul, P. & Roberts, D.J. (2013). Measurement of enteric methane from ruminants using a hand-held laser methane detector. *Animal Science*, vol. 63, pp 68-75. DOI: 10.1080/09064702.2013.797487

- Chesneau, G. & Weill, P. (2008). Method for evaluating the quantity of methane produced by a ruminant used for meat production. Valorex. FR2933191.
- Chilliard, Y., Glasser, F., Ferlay, A., Bernard L., Rouel J. & Doreau, M. (2007). Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *Europe Journal of Lipid Science Technology*, vol. 109, pp. 828-855. DOI 10.1002/ejlt.200700080
- Chilliard, Y., Martin, C., Rouel, J. & Doreau, M. (2009). Milk fatty acids in dairy cows fed whole crude linseed, extruded linseed, or linseed oil, and their relationship with methane output. *Journal of Dairy Science*, vol. 92, pp. 5199-5211.
- CIP (2013). Rolling out of an innovative enteric methane emissions measurement method for cattle. Eco-Innovation. <https://ec.europa.eu/environment/ecoinnovation/projects/en/projects/eco-methane>
- Crowcon production voucher (2013). *Laser Methane Mini- Methane Detection at a Distance*. [Leaflet] Issue 1 December 2013 GB. doi: MKG03114
- Czerkawski, J.W., Blaxter, K.L. & Wainman, F.W. (1966). The metabolism of oleic, linoleic and linolenic acids by sheep with reference to their effects on methane production. *Journal of Nutrition*, vol. 20, pp349-362.
- Danielsson, R. (2016) *Methane Production in Dairy Cows. Impact of Feed and Rumen Microbiota*. Diss. Uppsala: Swedish University of Agricultural Sciences.
- Danielsson, R., Ramin, M., Bertilsson, J., Lund, P. & Huhtanen, P. (2017). Evaluation of a gas in vitro system for predicting methane production in vivo. *Journal of Dairy science*, vol. 100, pp. 8881-8894.
- Danielsson, R., Schnürer, A., Arthurson, V., & Bertilsson, J. (2012). Methanogenic Population and CH<sub>4</sub> Production in Swedish Dairy Cows Fed Different Levels of Forage. *Applied and Environmental Microbiology*, vol. 78, pp. 6172–6179.
- Denniston, J.D., Topping, J.J., Caret, R.L., (2010) *General, Organic and Biochemistry*. 7th edition. New York. McGraw-Hill.
- Dewhurst, R.J., Fisher, W.J., Tweed, J.K.S. & Wilkins, R.J. (2003). Comparison of Grass and Legume Silages for Milk Production. 1. Production Responses with Different Levels of Concentrate. *Journal of dairy Science*, vol. 86, pp. 2598-2611.

- Doreau, M. & Ferlay, A. (1994). Digestion and utilisation of fatty acids by ruminants. *Animal Feed Science and Technology*, vol. 45, pp. 379-396.
- Eurofins Steins (2015)  
<https://www.eurofins.se/tjaenster/mjoelk/leverantoersmjoelk-mjoelkvalitet-och-sammansattning/>
- FAO, (2006), Livestock's Long Shadow—Environmental Issues and Options. Food and Agriculture Organization of the United Nations, Rome
- FAO (2010) Gerber, P., Vellinga, T., Opio, C., Henderson, B. & Steinfeld, H.  
 Greenhouse Gas Emissions from the Dairy Sector A Life Cycle Assessment. A report prepared by: Food and agriculture organization of the United Nations. Animal Production and Health Division.
- Gonthier, C., Mustafa, A.F., Ouellet, D.R., Chouinard, P.Y., Berthiaume, R. & Petit H.V. (2005). Feeding Micronized and Extruded Flaxseed to Dairy Cows: Effects on Blood Parameters and Milk Fatty Acid Composition. *Journal of Dairy Science*, vol. 88, pp. 784-756.
- Griinari, J.M., Dwyer, D.A., McGuire, M.A., Bauman, D.E., Palmquist, D.L. & NURMELA, K.V.V. (1997). Trans -Octadecenoic Acids and Milk Fat Depression in Lactating Dairy Cows. *Journal of Dairy Science*, vol. 81, pp. 1251-1261.
- Harfoot, C.G., Crouchman, M.L., Noble C.R. & Moore J.H. (1974). Competition between Food Particles and Rumen Bacteria in the Uptake of Long-chain Fatty Acids and Triglycerides. *Journal of applied Microbiology*, vol. 37, pp. 633-641. doi: 10.1111/j.1365-2672.1974.tb00487.
- Harvey, R. & Ferrier, D. (2011). *Biochemistry* 5<sup>th</sup> edition. Lippincott Williams &Wilkins. Baltimore.
- Hristov, A.N., Oh, J., Giallongo, F., Frederick, T., Weeks, H., Zimmerman P.R., Harper, M.T., Hristova, R.A., Zimmerman R.S. & Branco, A.F. (2015). The Use of an Automated System (GreenFeed) to Monitor Enteric Methane and Carbon Dioxide Emissions from Ruminant Animals. *Journal of Visualized Experiments*, vol. 103, pp,1-8 doi:10.3791/52904
- IPCC, (2007): Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, Pachauri, R.K and Reisinger, A. (eds.)]. IPCC, Geneva, Switzerland, pp.1-104

- IPCC, (2013): *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- Johnson, K.A. & Johnson, D.E. (1995). Methane Emissions from Cattle. *Journal of Animal science*, vol. 73, pp. 2483-2492.
- Knapp, J.R., Laur, G.L., Vadas, P.A., Weiss, W.P. & Tricarico J.M. (2013) Invited review: Enteric methane in dairy cattle production: Quantifying the opportunities and impact of reducing emissions. *Journal of Dairy Science*, vol. 97, pp. 3231-3255.
- Lang, K., Schuldes, J., Klingl, A., Poehlein, A., Daniel, R. & Brune, A. (2015). New mode of energy metabolism in the seventh order of methanogens as revealed by comparative genome analysis of "Candidatus Methanoplasma termitum". *Applied and environmental microbiology*, vol. 81, pp. 1338-1352.
- Larios, A.D., Brar, S.K., Ramí´rez, A.A., Godbout, S., Sandoval-Salas, F., Palacios, J.H., (2016). Challenges in the measurement of emissions of nitrous oxide and methane from livestock sector. *Reviewing Environmental Science of Biotechnology*, vol. 15, pp. 285-297. DOI:10.1007/s11157-016-9394-x
- Lindmark Månsson, H. (2012). *Den svenska mejerimjölkens sammansättning 2009* Svensk mjölk (Raport, 7094).
- Lindmark Månsson, H., Svensson, E. & Swensson, C. (2006). *Vallfodrets inverkan på mjölkens sammansättning och teknologiska kvalitet*. Swedish Dairy Association. Rapport nr 7066.
- Liu, Y. & Whitman, W.B. (2008). Metabolic, phylogenetic, and ecological diversity of the methanogenic archaea. *Annals of the New York Academy of Sciences*, vol. 1125, pp. 171-189. DOI: 10.1196/annals.1419.019
- Lovett, D.K., Stack, L.J., Lovell, S., Callan, J., Flynn, B., Hawkins, M., & O'Mara, F.P. (2005). Manipulating Enteric Methane Emissions and Animal Performance of Late-Lactation Dairy Cows Through Concentrate Supplementation at Pasture. *Journal of Dairy Science*, vol. 88 pp. 2836–2842.
- Machmüller, A., Soliva, C.R. & Kreuzer, M. (2003). Methane-suppressing effect of myristic acid in sheep as affected by dietary calcium and forage proportion. *British Journal of Nutrition*, vol. 90, pp. 529-540.
- Martin, C., Rouel, J., Jouany, J.P., Doreau, M. & Chilliard Y. (2008). Methane output and diet digestibility in response to feeding dairy

- cows crude linseed, extruded linseed, or linseed oil. *Journal of Animal Science*, vol. 86, pp. 2642-2650.
- McAllister, T.A. & Newbold, C.J. (2008). Redirecting rumen fermentation to reduce methanogenesis. *Australian journal of Experimental Agriculture*, vol. 48, pp. 7- 13.
- McDonald, P., Edwards, R.A., Greenhalgh, J.F.D., Morgan, C.A., Sinclair, L.A., Wilkinson, R.G. (2011). *Animal Nutrition*. 7<sup>th</sup> edition. Gosport. Ashford Colour Press Ltd.
- McGinn, S.M., Beauchemin, K.A., Coates, T. & Colombatto, D. (2004). Methane emissions from beef cattle: Effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *Journal of animal science*, vol. 82, pp. 3346-3356.
- Moss, A.R., Givens D.I. & Garnsworthy, P.C. (1995). The effect of supplementing grass silage with barley on digestibility, in sacco degradability, rumen fermentation and methane production in sheep at two levels of intake. *Animal Feed Science and Technology*, vol. 55, pp. 9-33.
- Moss, A., Jouany, J.-P. & Newbold, J. (2000). Methane production by ruminants: its contribution to global warming. *Annales de zootechnie*. vol. 49, pp 231-253.
- Nationalencyklopedin*, metan. Available at: <http://www.ne.se/uppslagsverk/encyklopedi/lang/metan> . [2016-02-06].
- Nielsen, N.I., Volden, H., Åkerlind, M., Brask, B., Hellwing, A.L.F., Storlien, T. & Bertilsson, J. (2013) A prediction equation for enteric methane emission from dairy cows for use in NorFor. *Acta Agriculturae Scand Section A*, vol. 63, pp.126–130. <http://dx.doi.org/10.1080/09064702.2013.851275>
- Norfor. *Search for feedstuff*. Available at: <http://feedstuffs.norfor.info/>. [2018-11-10]
- Patra, A.K. (2013). The effect of dietary fats on methane emissions, and its other effects on digestibility, rumen fermentation and lactation performance in cattle: A meta-analysis. *Livestock Science*, vol. 155, Pp. 244-254.
- Penner, G.P. Taniguchi, M., Guan, L.L., Beauchemin, K.A. & Oba, M. (2009). Effect of dietary forage to concentrate ratio on volatile fatty acid absorption and the expression of genes related to volatile fatty acid absorption and metabolism in ruminal tissue. *Journal of Dairy science*, vol. 92, pp. 2767-2781.

- Ramin, M. & Huhtanen, P. (2012). Development of an in vitro method for determination of methaen production kinetics using a fully automated in vivo gas system-A modelling approach. *Animal Feed Science and Technology*, vol. 174, pp. 190-200.
- Ramin, M. & Huhtanen, P. (2013). Development of equations for predicting methane from ruminants. *Journal of Dairy Science*, vol. 96, pp. 2476-2493.
- Ramin, M. & Huhtanen, P. (2014) Hur kan man mäta metan från mjölkkor? *NYTT*. Swedish University of Agricultural Sciences. No. 6.
- Ricci, P., Chagunda, M.G.G., Rooke, J., Houdijk, J.G.M., Duthie, C.-A., Hyslop, J., Roehe, R. & Waterhouse, A. (2014). Evaluation of the laser methane detector to estimate methane emissions from ewes and steers. *Journal of Animal Science*, vol. 92, pp. 5239-5250. doi:10.2527/jas2014-7676
- Sjaastad, Ö.V., Sand, O. & Hove, K. 2010. *Physiology of Domestic Animals*. 2nd edition. Oslo: Scandinavia Veterinary Press. 804 pp.
- Toprak, N.N (2015). Do fats reduce methane emission by ruminants? - a review. (2015) *Animal Science Papers and Reports*, vol, 33, pp. 305-321.
- Volden, H., 2011. NorFor. The Nordic feed evaluation system. EAAP publication No. 130. Wageningen Academic Publishers, The Netherlands.
- Van Zijderveld, S.M., Dijkstra, J., Perdok, H.B., Newbold, J.R. & Gerrits, W.J.J. (2011). Dietary inclusion of diallyl disulfide, yucca powder, calcium fumarate, an extruded linseed product, or medium-chain fatty acids does not affect methane production in lactating dairy cows. *Journal of Dairy Science*, vol. 94, pp. 3094-3104.
- Zachut, M., Arieli, A., Lehrer, H., Livshitz, L., Yakoby, S. & Moallem, U. (2010). Effects of increased supplementation of n-3 fatty acids to transition dairy cows on performance and fatty acid profile in plasma, adipose tissue, and milk fat. *Journal of Dairy Science*, vol. 93, pp. 5877-5889.
- Zimmerman, P. R., Zimmerman, R. S. (2012) Method and system for monitoring and reducing ruminant methane production. US 8,307,785 B2.
- Personal Communication: Nils-Erik Larsson, Växa Sverige, 2015.