



# **Aquaculture: Animal welfare, the Environment, and Ethical implications**

*Fiskodling: Djurvälstånd, Miljö och Etiska implikationer*

**Jenny Bergqvist**

**Etologi och djurskyddsprogrammet**

---

**Sveriges lantbruksuniversitet  
Institutionen för husdjurens miljö och hälsa  
Etologi och djurskyddsprogrammet**

**Skara 2010**

**Studentarbete 313**

***Swedish University of Agricultural Sciences  
Department of Animal Environment and Health  
Ethology and Animal Welfare programme***

***Student report 313***

**ISSN 1652-280X**



## **Aquaculture: Animal welfare, the Environment, and Ethical implications**

*Fiskodling: Djurvälstånd, Miljö och Etiska implikationer*

**Jenny Bergqvist**

Studentarbete 313, Skara 2010

**Grund C, 15 hp, Etologi och djurskyddsprogrammet, självständigt arbete i biologi, kurskod EX0520**

**Handledare:** Stefan Gunnarsson,  
Box 234, 532 23 Skara

**Examinator:** Harry Blokhuis,  
Box 7068, 750 07 Uppsala

**Nyckelord:** aquaculture, ethics, animal welfare, environmental impact

**Sveriges lantbruksuniversitet**

Fakulteten för veterinärmedicin och husdjursvetenskap

Institutionen för husdjurens miljö och hälsa

Avdelningen för etologi och djurskydd

Box 234, 532 23 SKARA

**E-post:** [hmh@slu.se](mailto:hmh@slu.se), **Hemsida:** [www.hmh.slu.se](http://www.hmh.slu.se)

---

I denna serie publiceras olika typer av studentarbeten, bl.a. examensarbeten, vanligtvis omfattande 7,5-30 hp. Studentarbeten ingår som en obligatorisk del i olika program och syftar till att under handledning ge den studerande träning i att självständigt och på ett vetenskapligt sätt lösa en uppgift. Arbetenas innehåll, resultat och slutsatser bör således bedömas mot denna bakgrund.

# CONTENTS

<b>1. ABSTRACT .....</b>	<b>4</b>
<b>2. SAMMANFATTNING.....</b>	<b>4</b>
<b>3. INTRODUCTION .....</b>	<b>5</b>
<b>4. ASPECTS OF ANIMAL WELFARE.....</b>	<b>6</b>
<b>4.1. Sentience .....</b>	<b>6</b>
<b>4.2. Breeding.....</b>	<b>8</b>
<b>4.3. Growth period .....</b>	<b>9</b>
4.3.1. Disease .....	10
4.3.2. Aggression and abnormal behaviours .....	10
4.3.3. Stocking density .....	11
<b>4.4. Capturing and transportation .....</b>	<b>12</b>
<b>4.5 Slaughter.....</b>	<b>13</b>
4.5.1. Asphyxia .....	14
4.5.2. Stunning by ice water immersion.....	15
4.5.3. Exsanguination.....	15
4.5.4. Electrocutation .....	15
4.5.5. Carbon dioxide stunning .....	16
4.5.6. Percussive stunning.....	16
4.5.7. Salt bath and evisceration.....	17
<b>5. ETHICS IN CONCERN OF ANIMAL WELFARE .....</b>	<b>17</b>
<b>6. ASPECTS OF ENVIRONMENTAL IMPACT .....</b>	<b>19</b>
<b>6.1. Decline of wild fish populations.....</b>	<b>19</b>
<b>6.2. Waste and chemical discharge.....</b>	<b>20</b>
<b>6.3 Loss of habitat .....</b>	<b>21</b>
<b>6.4 Spreading of diseases to wild fish populations .....</b>	<b>21</b>
<b>6.5 Invasion of exotic organisms .....</b>	<b>22</b>
<b>7. ETHICS ON CONCERN OF THE ENVIRONMENT .....</b>	<b>23</b>
<b>8. CONCLUSION.....</b>	<b>24</b>
<b>9. ACKNOWLEDGEMENT .....</b>	<b>25</b>
<b>10. REFERENCES .....</b>	<b>25</b>

## 1. ABSTRACT

The aim of this review is to assess the ethical implications of aquaculture, regarding fish welfare and environmental aspects. The aquaculture industry has grown substantially the last decades, both as a result of the over-fishing of wild fish populations, and because of the increasing consumer demand for fish meat. As the industry is growing, a significant amount of research on the subject is being conducted, monitoring the effects of aquaculture on the environment and on animal welfare. The areas of concern when it comes to animal welfare have here been divided into four different stages: breeding period; growth period; capturing and handling; and slaughter. Besides these stages, this report includes a chapter on the current evidence of fish sentience, since this issue is still being debated among biologists. Although the existence of fish sentience is still debated, most biologists are at present acknowledging the probability of fish being sentient creatures of welfare concern. Current aquaculture practices are affecting fish welfare during all four of these stages, both on physical and mental levels, as well as on the ability of fish to carry out natural behaviours. The effect fish farming has on the environment is here separated into five different categories: the decline of wild fish populations; waste and chemical discharge; loss of habitat; spreading of diseases; and invasion of exotic organisms. There is evidence of severe negative effects on the environment when looking at these five categories, even when considering the difficulty of studying environmental effects, due to the closely interacting variables.

## 2. SAMMANFATTNING

Syftet med denna litteraturstudie är att bedöma de etiska konsekvenserna av fiskodlingar, gällande fiskvälfärd och miljöaspekter. Fiskindustrin har vuxit kraftigt de senaste decennierna, både till följd av överfiske av vilda fiskpopulationer, och på grund av konsumenters ökade efterfrågan på fiskprodukter. Eftersom industrin ökar bedrivs en betydande mängd forskning på området, för att undersöka effekterna av fiskodling på miljö och djurskydd. De problemområden som rör djurskydd har här delats in i fyra olika stadier: avelsperiod, tillväxtperiod, infångning och hantering, samt slakt. Förutom dessa stadier innehåller denna rapport ett kapitel om aktuell forskning gällande medvetenhet hos fiskar, eftersom de vetenskapliga bevisen för medvetande hos fiskar fortfarande diskuteras bland biologer. Även om förekomsten av fiskmedvetenhet fortfarande är debatterad, är de flesta biologer numer överens om att sannolikheten för att fisk är kännande varelser är tillräckligt stor för att deras välfärd ska tas i beaktande. Nuvarande fiskodlingar påverkar fiskvälfärden under alla dessa fyra stadier, både gällande fysiska och psykiska aspekter, liksom fiskens förmåga att utföra naturliga beteenden. Effekten fiskodling har på miljön är här uppdelad i fem olika kategorier: minskning av vilda fiskbestånd, avfall och kemikalieutsläpp, förlust av naturliga habitat, spridning av sjukdomar, och invasion av exotiska organismer. Det finns tecken på allvarliga negativa effekter på miljön vid studerande av dessa fem kategorier, även då det är komplicerat att studera fiskodlingars miljöpåverkan på grund av att variabler interagerar med varandra.

### 3. INTRODUCTION

The amount of fish killed for food each year is steadily increasing, as the consumer demand for fish, as well as the world's human population, is growing rapidly. Through unsustainable fishing of wild populations of fish, the number of fish reared in farms has increased greatly the last decades. From previously answering for 9% of the total of fish killed for human consumption in the middle of the 1980:s, aquaculture systems produced 43% of the consumed fish in 2006 (Food and Agriculture Organization of the United Nations, 2006), and will soon be answering for the majority of fish killed for meat worldwide (Cottee & Petersan, 2009). Between 1987 and 1997 the production of fish and shellfish in aquaculture more than doubled (Naylor et al., 2000). The rising consumer demand for high quality fish meat, and at the same time sustainable fish farming concerning animal welfare and the environment, has led to an increased interest in aspects regarding improvements of fish farm practices (Lambooij et al., 2004). In addition, the increasing public concern regarding fish welfare in aquaculture (Huntingford et al., 2006), has resulted in a substantial amount of research the last decade.

The term welfare is used in a broad sense, and can be defined in various ways. However, there are three wide definitions often used: the feeling-based, the function-based, and the nature-based definition (Fraser et al., 1997). In short they mean that the welfare of the animal can be assessed either in regards of its own subjective feelings, emotions and mental state; its physical functioning, health and capacities; or by the ability the animal has to express its natural behaviour and lead a natural life (Fraser et al., 1997). In order to assess fish welfare, it is possible to either use one of these definitions, or to combine them for a more complete assessment. However, in many cases of fish welfare, all three definitions are related and they all lead to the same conclusion, since poor welfare defined by one approach often leads to poor welfare according to the other two (Huntingford et al., 2006). In previous research on fish welfare, the function-based definition of welfare is the most commonly used. This is noticeable since several studies are focusing on the physiological responses of fish, alone or in combination with others (such as behavioural responses), to potentially stressful conditions and situations.

Considering that the various species of fish reared in farms have evolved differently to adapt to their natural habitats, in aspect of salinity, temperature, depths etc., they naturally have various differentialities regarding behavioural and physical needs. Previous research regarding fish welfare has almost exclusively handled teleost fish, i.e. finfish (Huntingford et al., 2006), and although the term "fish" includes other species than the teleosts, this paper will in general focus on these species.

In contrast to other species of animals in agriculture, there has been very little research on the capacity to feel pain and psychological stress on fish species farmed for meat (Chandoo et al., 2004). This is so though the number of fish that are being killed for meat each year probably exceeds the number of all other animals reared for meat put together. The actual number of fish being killed each year is not possible to say, because in the statistics fish are being measured in weight instead of in individuals like other livestock animals, which might be considered as an indication of the status fish have in today's society. Also, farmed fish is often regarded and spoken of, not as animals to care for, but as crops to harvest (Lund et al., 2007).

The aim of this investigation is to review and discuss the potential risks of aquaculture practices, particularly in regards of fish welfare, but also concerning the environmental impacts of these practices. In addition, the ethical implications for animal welfare and environmental issues will be examined, since the cases of animal and environmental protection are the primary subjects of ethical discussion concerning aquaculture.

#### **4. ASPECTS OF ANIMAL WELFARE**

The potential risks for impaired welfare in farmed fish caused by humans occur under different phases of the production. Here these will be divided into four different stages: breeding; growth period; capturing and transportation; and slaughter. Before the issues of fish welfare regarding these stages are discussed, the question of fish sentience and potential to feel pain is briefed. This is to investigate whether the term welfare is at all applicable to fish.

##### **4.1. Sentience**

The existence of pain and suffering in fish has been widely studied and debated in recent years, with some conflicting results. These studies are important since they may determine whether animal welfare should be considered at all concerning fish and fish production. Science has not yet, with certainty, come to an agreement whether fish can feel pain or experience suffering, though in recent years most scientists do acknowledge that fish can experience at least something similar to the mammalian experience of pain.

If considerations should be made in aspect of fish welfare, they need to be regarded as sentient creatures, i.e. creatures with physiological and psychological characteristics that enable them to feel for example pain, fear and psychological stress. Their reactions to stimuli would therefore not only be a physiological response but also cause an associated psychological experience that requires cognitive capabilities (Chandroo et al., 2004).

Although finfish have nociceptors (receptors with ability to perceive painful stimuli), hence should be able to at least acknowledge pain, there still is a debate concerning the ability of fish to have an actual cognitive experience of pain (reviewed in Håstein et al., 2005). The neuro-anatomical structure of fish has both similarities and dissimilarities to tetrapods, i.e. birds, reptiles, amphibians and mammals (Chandroo et al., 2004). In aspects of sentience, there are more than one brain structures that are involved. For example, the limbic system is associated with emotions, behaviour, memory and learning, although the functioning of this part of the brain is not fully understood and therefore does not explain the whole concept of emotions (reviewed in Chandroo et al., 2004). It seems clear however, that the relatively primitive brain structures of the limbic system are to some extent involved in emotional and cognitive responses (Chandroo et al., 2004), and the fact that fish neuro-anatomy includes the limbic system is an indicator that it is at least a possibility that fish are capable to feel aversion to noxious stimuli. The fish brain though, is far smaller related to body size than the human brain, and does not possess all of the structures that the human brain does (reviewed in Huntingford et al., 2006; Lund et al., 2007). In question of ability to feel pain, the neocortex in the telencephalon, or forebrain, is of special interest, since it is partly this structure that answers for the subjective feeling of pain in humans and other mammals, and a structure that the fish brain is lacking (reviewed in Huntingford et al., 2006; Lund et al., 2007). The question seems to be whether fish can experience subjective

feelings similar to the human feeling of pain, or if they merely react reflexively to noxious stimuli.

The issue of similarity to human or mammalian brain may be over-rated regarding ability to experience pain, fear and suffering. It is known that different parts of the brain in different groups of animals can have similar functioning (e.g. Huntingford et al., 2006; Lund et al., 2007). For instance, cephalopods are able to carry out complex behaviour with a very different brain structure than humans (Håstein et al., 2005; Huntingford et al., 2006), and the bird brain is, like the fish brain, lacking a neocortex and, nevertheless, birds are considered to be able to experience pain (reviewed in Lund et al., 2007). In regard of this, Huntingford and co-workers (2006) states that it should be possible for fish to experience the subjective mental state of pain or suffering without the neocortex brain structure.

In a study of fear in rainbow trout, Yue and co-workers (2004) used an avoidance learning task to determine whether rainbow trout, one of the most commonly farmed fish, could have the subjective experience of fear, instead of a merely reflexive response to fearful stimuli. The authors reason that if fish can show behavioural adjustments in their response to such stimuli, they are not just reacting with an innate reflex but are learning strategies to handle fearful situations and possible danger. In this study, the fish were subjected to a stimulus in form of a plunging net that they could escape from by swimming through a doorway into another part of the water tank. Ten seconds before the net plunged into the tank a light went off, testing if the fish could learn to avoid the net by escaping it before hand. Most fish learned to avoid the net when the light went off, and even showed that they have the capacity of long term memory, since they kept responding to the light stimuli even after seven days of no trials. The fish that did not learn to swim through the doorway at the illumination of the light mostly remained motionless and sank to the bottom of the tank, which was interpreted as a possible sign of severe stress. Yue and co-workers (2004) made the conclusion that rainbow trout not only have the ability to experience fear, but are also capable of long term remembrance of fearful experiences, as well as, performing behavioural modification to deal with stressful situations. This conclusion is in accordance with previous studies of fear in fish (reviewed in Yue et al., 2004). Yue and co-workers suggest, based on the findings of this study, that the emotional welfare of farmed fish needs to be taken into serious consideration, something that is not currently being done.

Behavioural coping strategies and neuro-endocrine responses to stress present in mammals have also been found in fish. These physiological responses to chronic stress include growth and immunosuppression, loss of appetite and impaired reproductive success, indicative of compromised health and, hence, welfare (Huntingford et al., 2006). Although occurrence of stress responses in fish should not be automatically viewed as indicators of poor welfare or suffering; prolonged or repetitive responses can be indicative of exposure to stressors that the animals are unable to handle aptly, and would therefore be of welfare concern (Huntingford et al., 2006).

Chandroo and co-workers (2004), state that the studies made on fish anatomy and behaviour suggests that fish are more likely to experience pain and suffering than not. Although the fish brain does not include the neocortex like the mammalian do, they react similarly to painful stimuli and analgesics as mammals do (reviewed in Chandroo et al., 2004; Huntingford et al., 2006). Based on the anatomy, behaviour and seemed cognitive capabilities of various fish species, Huntingford and co-workers (2006) state that the possibility of fish consciousness and ability to experience pain is likely, and therefore there

should be concerns for their welfare. It has also been suggested that the different tactic behaviour of individual fish, during situations such as reproduction, is complex enough to require memory and learning skills (St-Cyr & Aubin-Horth, 2009), hence fish behaviour in at least some cases, is beyond instinctual. Lund and co-workers (2007) also state that due to the physiological processes, as well as the complex behavioural patterns of fish indicating cognitive ability, by all likelihood mean that fish are capable of the conscious experience of pain.

Generally, the contributors to this discussion have drawn the conclusion that since the similarities between fish and mammals do not only include some aspects of neuro-anatomy, but also other anatomical and physiological structures and responses, as well as behavioural similarities, fish are creatures that should be taken into consideration regarding animal welfare and humane treatment (e.g. van de Vis et al., 2003; Lambooij et al., 2004; Huntingford et al., 2006; Lund et al., 2007). It can be concluded that fish have the sense organs and neural system acquired to react and respond to painful stimuli, and most likely to experience a subjective feeling of pain, as well as other mental states such as suffering. Thus, for the remainder of this paper, fish will be regarded as sentient creatures.

## **4.2. Breeding**

The natural spawning can be intervened in aquaculture by procedures that enhance the gonad maturation process. This can either be done by altering the photoperiod, or by hormonal injections (Donaldson, 1996), which can have impact on fish welfare. Especially the hormonal injections appear to have negative effects of the animals' wellbeing (Grigorakis, 2009). Aspects besides handling procedures in the spawning process may have negative effects on fish. For example, breeding fish for productive features, such as fast growth and high feed conversion efficiency, has had some unexpected welfare consequences. Due to genetic and/or environmental factors, conditions such as deformities of heart, swim bladder, spine, jaw, and lips are not uncommon in farmed fish, and impair welfare through poor health and impaired swimming ability, as well as capacity to compete for food (Imslund et al., 2006; Ashley, 2007). Affected fish show decreased stress resilience and have increased mortality rates compared to others (Ashley, 2007), indicating impaired welfare. In addition to the hindering of competitive and swimming abilities, it is likely that such malformations are painful for the individual fish, although research on this subject is lacking.

There are a number of different malformations of the spine that are common in farmed fish, they have been observed in most farmed fish species, and they are considered to be effects of both hereditary factors, e.g. inbreeding; and environmental factors, e.g. nutrition imbalance, especially vitamin deficiency, or inappropriate temperatures (reviewed in Afonso et al., 2000; Imslund et al., 2006). A study of Atlantic cod (*Gadus morhua*) by Imslund and co-workers (2006) found evidence implying that the feed regime during the larval stage had major implications for anatomical development and occurrence of skeletal deformations. In intensive aquaculture, feed for larvae compose mainly on rotifers, but when compared to zooplankton, led to three times the incidence of skeletal malformations, with an average of 14.2% versus 4.1% (Imslund et al., 2006). An introduction of zooplankton as feed in commercial fish farms might be difficult however, since the plankton are only accessible under a limited part of the year (Imslund et al., 2006). Yet this study points out an urgent matter for intensive aquaculture, namely that current breeding



and feed practices may be efficient enough regarding fish production, but mostly not concerning fish health and welfare.

### **4.3. Growth period**

The growth period is the longest stage for most fish in aquaculture, hence possibly the most important one from a welfare perspective. Some of the potential risks for fish welfare during this stage include crowding, handling, diseases, aggressive behaviour causing injuries or food deprivation, poor water quality, and starvation before transportation or during disease treatment. The intense rearing conditions commonly seen in commercial fish farms are often causing poor water quality, leading to increased aggression and disease susceptibility; reduced growth, stress and distress; and mortality (reviewed in Håstein et al., 2005). This is certainly a concern for fish welfare and closely interacts with other issues of concern in subsequent sections.

#### 4.3.1. Disease

To prevent and control spreading of disease among fish in aquaculture, knowledge of the interaction between pathogen, fish, and environment is required (Toranzo et al., 2005).

Physiological responses to chronic stress in fish include immunosuppression, hence repeated occurrences of diseases and high mortality rates can be an indication of a poor environment for, and impaired welfare of, farmed fish (Huntingford et al., 2006). Causes for chronic stress associated with disease susceptibility can for example be fin abrasions, aggressive social interactions, and crowding (Ashley, 2007). Obviously, also without stress and poor living conditions, disease itself can lead to suffering (Huntingford et al., 2006), hence it should be treated properly and prevented if possible. There are numerous diseases farmed fish can be exposed to, thus a thorough assessment is beyond the scope of this paper. However, some examples of diseases causing especial nuisance in aquaculture will be briefly presented.

One parasitic infestation that is common in aquaculture and of serious concern regarding fish welfare is sea lice (Ashley, 2007). It is associated with tissue damage as the lice eat through the skin (Ashley, 2007), and can cause such severe head wounds that exposure of the skull bones occurs (Håstein et al., 2005). This infection is therefore of serious welfare concern, and attempts to prevent it, as well as sufficient treatments of it, are critical for fish health. Besides parasitic infestation, viral diseases are also currently a serious threat to welfare for fish in aquaculture, with only moderately successful vaccines available (Ashley, 2007). Bacterial diseases are also of great impact in aquaculture, with a few common types of pathogens, such as vibriosis, furunculosis, streptococcosis, and mycobacteriosis, having vast influences on fish health (Toranzo et al., 2005). Some bacterial diseases, earlier associated with mainly fresh water farms, have now spread to marine ones (Toranzo et al., 2005). These pathogens generally cause high mortality rates in aquaculture, although the same mortality rates for wild fish are low, demonstrating the impact intense aquaculture's stressful conditions have on fish immune system (Toranzo et al., 2005). Vaccination for preventing diseases has increased and a reduction in infectious diseases has been observed in concerned farms (Håstein et al., 2005). The vaccinations themselves though, pose a threat for fish welfare through increased handling, inflammation of the body part injected, and occasional deformations of the spine (Håstein et al., 2005).

Fin rot is a condition resulting from fin necrosis, which means that eroded fins become infected with a variety of potential bacteria, causing ulcers and degrading the fish's swimming ability, as well as causing even mortality (Latremouille, 2003). The erosion of fins can be inflicted by malnutrition, aggressive interaction, overcrowding, or inappropriate pen design (Latremouille, 2003), making it not just a risk to welfare itself, but also a symptom of risked welfare through improper rearing conditions.

#### 4.3.2. Aggression and abnormal behaviours

Fish species with hierarchal social orders can, particularly in case of improper housing conditions such as high densities and deficient feeding methods, express aggressive behaviour that leads to social stress, especially for the subordinates (Ashley, 2007). For low ranking individuals, this may lead to immunosuppression, food competition with decreased growth, decreased swimming activation, and food deprivation, alongside fin and eye injuries with infections and increased mortality as results (reviewed in Ashley, 2007). Furthermore, social traits such as increased aggression can be partially caused by unintentional selection, since behaviour is influenced by genetic, as well as environmental factors (St-Cyr & Aubin-Horth, 2009).

Rearing of Atlantic cod (*Gadus morhua*) in fish farms is problematic, since highly aggressive behaviour and cannibalism is common among juveniles (Höglund et al., 2005). Though Höglund and co-workers (2005) acknowledge that hierarchal and aggressive behaviours are important parts of the juvenile Atlantic cods behavioural repertoire, the authors suggest a use of feed supplement (L-tryptophan, or TRP) that is aggression suppressing, since “[...] by changing the behaviour of the fish instead of altering the social and/or physical environment, TRP-supplemented feed could be more cost-effective and less time-consuming than other manipulations aimed at decreasing cannibalism”. Other manipulations of the environment for farmed cods named by the authors are feeding strategies and usage of optimal stocking densities. The freedom to express natural behaviour would hereby be compromised, not only by rearing fish in improper conditions, but also by feeding them supplements that essentially alter their behaviour. The reason for this given is that altering fish behaviour is less costly and time-consuming than changing the inadequate rearing system that originated the problem to begin with. Although it is reassuring that aggressive behaviour of farmed fish is regarded as a concern for scientific research, it may be viewed as incautious or unethical to manipulate the behaviours of fish to solve issues of improper production practices.

Besides aggression, abnormal behaviour in farmed fish, such as atypical swimming, is as well a potential threat to welfare. These behaviours can be results of suppression of behavioural needs, indicative of exposure to stressors and potentially poor welfare, as the animal makes unsuccessful efforts to alter its situation (Ashley, 2007). In other cases, fish abnormal behaviour is consequences of injury, disease, or other health conditions (Ashley, 2007). Whichever, the abnormal behaviour of fish should be taken into concern, evaluated, and handled.

The lacking comprehension of behavioural needs for farmed fish, both social and individual such, represents a serious risk for fish welfare (Håstein et al., 2005). Regarding fish as sentient creatures with individual needs besides physical health is a critical step towards treating fish morally acceptable. Further scientific evidence is needed to fully realize the welfare implications fish farm conditions have on fish behaviour and welfare.

#### 4.3.3. Stocking density

The impact of stocking density on fish welfare is difficult to assess, due to the interaction between environmental factors, such as water carrying capacity, as well as the diverse physiological and behavioural requirements of different species (Håstein et al., 2005; Ashley, 2007). For instance, solitary living fish may be stressed of, and turn aggressive in high densities, while for fish that naturally live closely together with others in large numbers, too low a density may be stressful (Ashley, 2007).

Rainbow trout is the main fish species being brought up for meat in fish farms in Europe and North America (Ellis et al., 2002). The stocking density varies among farms, which are mainly guided by the farmers' own experience, since there rarely is any legislation controlling fish density (Ellis et al., 2002). Usually the density varies between 15 and 40 kg per m<sup>3</sup> in Europe, North America and Australia, but both higher and lower densities are in practice (reviewed in Ellis et al., 2002). Animal welfare in fish farms has recently become a central issue, because it has been shown that the meat quality and fish health are closely linked to the welfare of the animals (Ellis et al., 2002). Although there are conflicting results, studies have shown that an increase in density leads to an increase in gill and fin injuries, as well as mortality of farmed rainbow trout (reviewed in Ellis et al., 2002), and that the overall welfare is impaired (Ashley, 2007). Others studies have shown that a low density can lead to increased fish mortality, since the aggression of rainbow trout can be triggered by low densities, and this can raise the susceptibility of disease and infection due to the damage inflicted (reviewed in Ellis et al., 2002). In high density farms as well, an associated problem seems to be an increased disease occurrence, which could be explained by poorer water quality, easier spread of pathogens, and that the immune system of the fish could be impaired by the stress crowding inflicts (Haya et al., 2001; Ellis et al., 2002). Studies have shown that feed intake decreases with increased density in fish farms, and studies have also shown that feed conversion efficiency decreases with increased densities (reviewed in Ellis et al., 2002), which clearly demonstrates that animal welfare issues are important not only for the reared fish and concerned consumers, but also have an economic importance to the farmers.

To assess the fish welfare implications of high fish density in rainbow trout, Ellis and co-workers (2002) state that more research is needed. The studies cited in their review have conflicting results and do not clearly show that high density is a stressor that will impair fish welfare. Ellis and co-workers (2002) do conclude though, that there are signs suggesting that the welfare of the animals is at risk with high densities, as well as with too low densities. The increase in gill and fin damage, as well as the increased mortality in at least part of the high density farms studied are, according to Ellis and co-workers (2002), suggesting that high fish density in rainbow trout farms are likely resulting in a poorer welfare for the animals. The authors further discuss the welfare implications of decreased food intake and growth rate in high density farms, and argue that to be sure of impaired welfare for these fishes, it has to be shown that the fish are suffering from hunger, stress or other discomforts associated with decreased food intake or growth rate, since these effects do not mean impaired welfare per se. However additional studies (reviewed in Håstein et al., 2005), have shown that increased stress levels and mortality, inhibited swimming behaviour, and decreased ability to digest food are results of high densities for farmed fish. Other aspects of high density in fish farms that could impair welfare are that the behavioural needs for the fish might be made impossible to meet, and that the water quality is often poorer in high densities, due to more intense use of the water for metabolic

reactions and that increased fish movement will hinder the particles in the water to settle (Ellis et al., 2002). This may lead to stress, aggressive behaviour and damages to the gills and fins due to collisions (Ellis et al., 2002; Håstein et al., 2005). These evidences illustrate that high density is undoubtedly a matter of welfare concern.

The implications of high stocking density for farmed fish vary among different species, and are, as mentioned, interconnected with other rearing conditions, such as water quality and food availability (Turnbull et al., 2005; Ashley, 2007). The difficulty to assess welfare, due to lack of research, with the wide range of different species reared in aquaculture further complicating the matter, lead to difficulties for development of appropriate densities recommendations (Turnbull et al., 2005; Ashley, 2007). However, in a study of Atlantic salmon (*Salmo salar*), Turnbull and co-workers (2005) found that high density impair welfare if above a threshold density of 22 kg m<sup>3</sup>, when body and fin condition, as well as plasma concentration of cortisol and glucose was measured. The water conditions regarding temperature and oxygen levels during the study was for the more part within the recommended ranges, and should therefore not have influenced the results. The authors conclude though, that a recommended threshold density for farmed fish does not guarantee fish welfare, since there are a number of factors connected to density that affects the animals' well-being. This means that, for some farms, a density of 22 kg m<sup>3</sup> would be appropriate, but for others a decreased density would be required due to differences in farm practices and conditions. Ashley (2006) suggests behavioural observations of on farm fish as an important measure animal keepers can take to see indications of poor welfare due to improper stocking density.

#### **4.4. Capturing and transportation**

Fish are mainly captured and transported when transferred to pens for growth or in connection to slaughter. Although the duration of these stages are comparably short to the growth period, they are vital stages in regards of welfare, since they can cause numerous potential stressors for the animals. Examples of such are crowding, handling during capture, poor water quality with low oxygen levels, food deprivation, exhaustion, injuries, as well as increased aggression and spreading of diseases.

When moved to the new environment of the growth pen, acclimatisation to this environment regarding water composition, temperature, and light levels, needs to be considered and assured (Ashley, 2007). The handling and transport itself to the growth site is stressful for the juvenile fish, thus their ability to cope with the transfer can be subsided, with increased mortality as a result (Ashley, 2007).

Before slaughter, when captured and transported, the fish are subjected to a number of stressors, such as crowding with skin abrasions, as well as low oxygen and high ammonia levels in the water as results of the crowding (Poli et al., 2005; Gatica et al., 2008). Any handling of fish before slaughter, such as capturing or transfer from transport vehicle to holding pens before slaughter, is a stressful experience for fish (Håstein, et al., 2005), due to the crowding and struggling the fish are facing, as well as the duration of the process and the poor water quality the crowding leads to (Poli et al., 2005; Gatica et al., 2008). This exhausts the fish, and as they are most often not given the opportunity to recover before the slaughter (Poli et al., 2005), it also affects their ability to handle the actual stress of being stunned and slaughtered. Since the struggle decreases the pH in the fish musculature, and the animals are not allowed to recover, this is shown in the meat quality after slaughter

(Poli et al., 2005; Gatica et al., 2008). To avoid this low meat quality, fish are often starved several days or even weeks before slaughter, for sake of water quality during transport and so that the animals' energy reserves are emptied and they do not have the energy to struggle during capturing (Poli et al., 2005; Gatica et al., 2008). Hence the pH does not drop in the muscles, though the procedure is sure to be further stressful to the fish (Poli et al., 2005; Gatica et al., 2008). However some evidence shows that the food deprivation itself is not necessarily a source of critically impaired welfare, thought mainly to depend on the fact that fish are ectothermic (Huntingford et al., 2006). However, the starvation has an impact, on their energy reserves and therefore on the overall condition of the fish, and also results in stress and possible changes of their social behaviour such as increased aggression due to competition over food, or result in abnormal behaviours such as "eye snapping" (Håstein et al., 2005; Ashley, 2007), which may lower their ability to cope with other sources of stress during this stage.

When the fish are captured they usually get loaded onto the boat by pumps and then again transferred by pipes from their resting cages or directly from the boat to slaughter (Gatica et al., 2008). The pumps and pipes transferring the fish stress the animals and can cause injuries and skin abrasions, therefore they should be well constructed and the duration of transfer in pipes should be as short as possible (Ashley, 2007; Gatica et al., 2008). Overcrowding occurs in the resting cages, in purpose of optimising productivity, but this is also stressful for the animals and leads to aggressive behaviour and injuries among the fish, as well as an increased risk of diseases spreading (Ashley, 2007; Gatica et al., 2008). The stress caused by capturing, handling and transport should be minimized by, ideally, slaughtering the fish directly after capturing, and if that is not possible, by improving the practices used in handling and transport to slaughter (Gatica et al., 2008).

Handling and other disturbances in the pre-slaughter process are also associated with stress for the fish (Håstein et al., 2005; Poli et al., 2005; Ashley, 2007; Gatica et al., 2008). Being taken out of water is a severely stressful situation for fish; thus setting off an emergency response leading to high plasma cortisol levels and behaviour indicating acute stress, such as excessive struggling and attempts to escape (Håstein et al., 2005). Manual handling can cause skin abrasions, along with scale and mucous coat removal (Ashley, 2007). The mucous coat serves as a protection from infections, and if damaged the fish is much more susceptible to diseases, as well as weakened in locomotion and osmoregulation abilities (Ashley, 2007). Thus, handlers should always take precautions when handling fish, such as using appropriate nets, gentle and wet hands, and making the duration for fish out of water as short as possible (Håstein et al., 2005; Ashley, 2007).

Continuous or multiple stressful situations for fish are additive and exhaust the animals, making them less capable to handle further stressors (Ashley, 2007; Gatica et al., 2008). Therefore, in aspect of animal welfare, the fish should not be starved longer than enough for the stomach to empty, so that the water quality during transport and pre-slaughter is not further lowered by waste, and the fish in that same aspect need to be rested and free from skin abrasions, muscle bruises or other injuries (Poli et al., 2005; Ashley, 2007; Gatica et al., 2008).

#### **4.5. Slaughter**

The moments leading up to death for fish in aquaculture may be short in terms of life span, but as discussed below, is often associated with severe stress and pain for the animals.

There has been a substantial amount of research in slaughter of farmed fish recently, perhaps due to the poor meat quality of stressed and exhausted fish, but probably also because of the increased public concerns regarding current slaughter procedures. Slaughter techniques are often adjusted to assure efficiency, not fish welfare (Ashley, 2007), and there are various different methods used in commercial fish farms worldwide, most of them described below. Methodology used to assess sensibility, consciousness, and reactions to stimuli in fish is also briefly described.

Ability to confirm unconsciousness, and therefore insensibility, in fish depends on sufficient methodology, since behavioural observations alone may be misleading due to the immobility that can occur without loss of consciousness (van de Vis et al., 2003; Lambooij et al., 2004). Other ways to assess the stress response and sensibility of fish, such as electroencephalogram, or EEG; electrocardiogram, or ECG; plasma hormonal levels; somatosensory evoked responses; and visual evoked responses, are available as a complement to the behavioural parameters (van de Vis et al., 2003; Lambooij et al., 2004; Ashley, 2007). The somatosensory evoked responses is brain responses to painful stimuli (such as a scratching with a nail), and visual evoked responses is brain responses to stimuli in form of a flashing light directed at the eye. Insensibility is concluded when the brain responses do not occur, this due to brain disruption or brain failure (reviewed in van de Vis et al., 2003; Lambooij et al., 2004). The occurrence of visual or somatosensory responses does not make it possible to automatically assume conscious awareness of light or pain stimuli, though it is a sign of intact pathways to higher brain centres and implies remained sensibility (Lambooij et al., 2004).

There are different methods of stunning and killing farmed fish in praxis, both forceful methods that quickly render fish unconscious, and methods that slowly decrease the level of consciousness in the animals. With certainty, handling, stunning and slaughter are highly stressing moments for farmed fish whichever method being used, and the more stressful it is, the lower meat quality, which makes it measurable (reviewed in Poli et al., 2005). Poli and co-workers (2005) view the relative fast working methods of some types of stunning (e.g. knocking and spiking) generally as less stressing for the fish than other methods, though they point out that these methods are not always practical, for example with large numbers of smaller sized fish.

The following methods of slaughter are currently being used for farmed fish in Europe: asphyxia, carbon dioxide narcosis, ice stunning, spiking (though used mostly on an experimental level), knocking, exsanguination through gill cutting or decapitation, salt bath in combination with evisceration (i.e. gutting), and electrical stunning (e.g. van de Vis et al., 2003; Lambooij et al., 2004; Poli et al., 2005). The following section deals with these different slaughter methods.

#### 4.5.1. Asphyxia

The oldest method of fish slaughter is asphyxia, which essentially means suffocation through taking fish out of water, and has been shown (for example in plasma cortisol levels and meat quality) to be associated with high occurrence of stress reaction in the animals (Poli et al., 2005; Ashley 2007). The duration until fish lose consciousness varies widely among different species, depending on how resistant the species is to hypoxia, and ranges from about five minutes to over an hour (Poli et al., 2005). Behavioural studies of fish being slaughtered in this manner also indicates that this experience is highly stressful, based on observations of attempted escapes and other avoidance behaviour, and this

method is thought to mean prolonged suffering before unconsciousness occurs (Ashley 2007; and reviewed in van de Vis et al., 2003; Poli et al., 2005).

#### 4.5.2. Stunning by ice water immersion

Cause of death when fish is being immersed in ice water is anoxia and the time until stunning varies among species also in regard of this method (Poli et al., 2005; Ashley 2007), from 12 minutes (in eel) to up to 198 minutes (in trout) (reviewed in Poli et al., 2005). This method has been concluded to be inhumane due to behavioural observations of vigorous movement, and because of the length of time from immersion to unconsciousness (van de Vis et al., 2003; Poli et al., 2005; Ashley, 2007), although in some species of fish that are adapted to warm water, the level of stress seems to be relatively low, in for example plasma glucose and lactate levels, as well as the behavioural reaction of the fish with little or no violent reactions (reviewed in Poli et al., 2005). In comparison with asphyxia, this seems to be a less stressful experience for the warm water adapted species of fish (Poli et al., 2005), but is still considered inhumane due to the duration until stunning (Ashley, 2007).

#### 4.5.3. Exsanguination

Death by gill cutting, without previous stunning is not considered to be a humane method of slaughter. Studies have found that the behaviour of Atlantic salmon, such as vigorous movements, indicates occurrence of sensibility at least for the more part of the time from cutting until death (Robb et al., 2000). van de Vis and co-workers (2003) found that the visual evoked responses to light stimuli were not lost directly after the cut, further indicating the method's welfare implications for fish. Lambooij and co-workers (2004) studied ECG and somatosensory evoked responses in African catfish following gill cutting without stunning, and found that although the ECG patterns depressed after 2-5 minutes, the somatosensory evoked responses lasted 15 minutes in all fish tested, at which time the tests were aborted and the fish killed. Gill cutting as a method of slaughter is yet in practice in countries such as United Kingdom and Norway (Robb et al., 2000). In comparison, decapitation is a way of slaughtering eels in the Netherlands, although rarely used except by retailers, since it is not practical when slaughtering eels in great numbers (van de Vis et al., 2003). This is by certain not a humane treatment of eels: tests have shown that the animals do not lose their visual evoked responses for an average of 13 minutes after the cut (van de Vis et al., 2003) and eels can stay alive for up to eight hours after decapitation (Verheijen & Flight, 1997). Therefore this method is detrimental from an animal welfare perspective.

#### 4.5.4. Electrocutation

Electrocutation can be an effective, and more humane, method of stunning considering that it can lead to an immediate loss of consciousness, as well as reduced handling, transportation, and taking fish out of water, when carried out in water (van de Vis et al., 2003; Ashley, 2007). Electricity can either be applied over the whole body, or across the head only, if done on individual fish (van de Vis et al., 2003). Without high enough electrical current used though, immobilization without lost sensibility may result, hence becoming a threat to welfare (Ashley, 2007).

Lambooij and co-workers (2004) studied head-only stunning with electricity and subsequent gill cutting on African catfish. Two out of seven fish responded to noxious stimuli up to five minutes after 1.2 seconds of electricity applied and gill cut, even though a relatively high current was being used. According to van de Vis and co-workers (2003) one

second of application across the head only lead to sufficient stunning in one out of ten sea bream tested, and the fish with brain disruption regained consciousness after only 37 seconds, when visual evoked responses were observed. Ten seconds of electricity applied across the head was more efficient with six out of ten fish stunned for at least ten minutes and the fish were assumed to be dead, though one of the ten sea breams tested did not lose visual evoked responses at all, and three of the stunned fish recovered within 16 seconds. For these results, the current used varied and the fish who was sufficiently stunned, received relatively high currents (over 400 mA). The low currents often being used in commercial farms is explained by the violent muscle contractions the electrocution leads to, which can leave blood spots in the muscles, as well as broken bones, and in that sense give decreased meat quality (Poli et al., 2005; Ashley, 2007). However, with an optimised current and insurance that death will occur before regained consciousness, this method can be quicker and less stressful to fish than most other methods currently used in commercial fish farms (van de Vis et al., 2003; Lambooij et al., 2004; Poli et al., 2005; Ashley, 2007).

#### 4.5.5. Carbon dioxide stunning

A widely used method of stunning of farmed fish is carbon dioxide stunning. The fish are stunned by immersion in water with added carbon dioxide, CO<sub>2</sub>, which decreases the pH-level in the blood and intoxicates the brain, leading to narcosis (Poli et al., 2005; Ashley, 2007). When carbon dioxide is added to water, it dissolves to carbonic acid, H<sub>2</sub>CO<sub>3</sub>, which is highly aversive to mucous membranes, and causes intense avoidance behaviour in fish, such as frantic swimming and attempts to escape the water (Rollin (1993) in Håstein et al., 2005). The duration until stunning varies, but although immobility is reached within 30 seconds to 4 minutes, sensibility remains up to 5 minutes, often up to being bled by gill cut (reviewed in Håstein et al., 2005; Poli et al., 2005). It has been concluded by a number of authors that this method is not humane (e.g. van der Vis et al., 2003; Håstein et al., 2005; Ashley, 2007), yet still it is an internationally common practice in aquaculture.

#### 4.5.6. Percussive stunning

For larger fish, insensibility by a blow to the head using a club can be used as a stunning method, leading to concussion, a disruption of the brain and immediate unconsciousness or death (van de Vis et al., 2003; Poli et al., 2005). If done correctly this is therefore one other most humane methods (van de Vis et al., 2003; Poli et al., 2005), though to hit this relatively small target in the correct way it requires trained, focused, and rested personnel (Poli et al., 2005; Ashley, 2007). In practice, the first hit is often not accurate enough to disrupt the brain, leaving the fish with remained sensibility, and are therefore hit more than once before stunning (reviewed in van de Vis et al., 2003; Poli et al., 2005). The repeated hitting leads to suffering for fish, since consciousness is not lost immediately. Although, this method is one of the least stressful, if quick and proper stunning can be assured.

Another fast method associated with less stress reactions than others is spiking, used mainly on tuna and salmon (Poli et al., 2005). A spike is inserted into the brain and unconsciousness is immediate, if done correctly (Poli et al., 2005). The spike must hit a specific point to ensure stunning, and if it misses the small target inside the skull, it is painful, leaving the fish with full consciousness and tissue damage caused by the spike (Poli et al., 2005). Automatic stunning equipment may thus represent an improvement of this method, since it is more accurate and minimal handling is possible (Ashley, 2007), although care must be taken to use the appropriate pressure and size adjustments to ensure successful stunning before exsanguination (Håstein et al., 2005).



#### 4.5.7. Salt bath and evisceration

For eels, one method of slaughter being used, mainly in the Netherlands, is salt bath in combination with evisceration (van de Vis et al., 2003). The eels are exposed to NaCl in a dry tank for desliming for about 20 minutes, and then eviscerated and bled to death (van de Vis et al., 2003). The whole procedure takes approximately an hour, and the eels might have maintained consciousness under the entire process (Ashley, 2007), and eels have even shown signs of life for up to 18 hours after initiation of the procedure (Verheijen & Flight, 1997). This is not considered to be a humane method, also because of the behavioural signs of severe stress, and the attempts of eels to escape the tank for at least three minutes after addition of salt to the tank (van de Vis et al., 2003).

The stress levels of fish before slaughter can be so severe, due to pre-slaughter procedures, that the positive effects of a more humane method of stunning no longer shows any significant differences in physiological measuring (Poli et al., 2005). Therefore, although slaughter methods are important for fish welfare purposes, cautious handling in the capturing and pre-slaughter procedure cannot be over-emphasized.

## **5. ETHICS IN CONCERN OF ANIMAL WELFARE**

It is but recent that fish ethics has become a real issue of discussion. Other animals in human care have been of interest for research and public concern for decades, and it would seem that fishes are the last to be included in the moral debate. This is likely due to more than one reason, perhaps most worth mentioning the lack of human-fish interaction. Humans have had relationships with other animals under a long time, building strong emotional bonds. With fish however, bonds are hard to make, depending on the obvious water-air boundaries, as well as difficulty for humans to communicate with fish; we are not capable of interpreting fish body language and facial expressions, and they make no human receivable sound to indicate their mental states (Lund et al., 2007). Although fish are unsuccessful of evoking feelings of empathy in most humans, does this mean that it is right to exclude them from moral concern? Lund and co-workers (2007) argues that 1) sentient animals should be of moral concern, 2) fish are most likely sentient, thus 3) fish should have the right to be morally treated. Hence, the level of empathy humans have for different species, should not be determining the treatment received.

Although the moral significance of animals is essentially an ethical issue, the base for such ethical reasoning should be based on scientific research of animals' ability to experience suffering or pleasure, as well as on which conditions or situations that inflict such experiences (Lund et al., 2007), to enable sound ethical conclusions. Mainly the animal's cognitive abilities has been the focus of animal ethics, making fish a group of animals largely neglected in sense of moral concern of the individual animal (Lund et al., 2007), since the cognitive abilities of fish are questioned and debated. However, the substantial amount of new research in this subject has led to an increased interest in welfare of the individual fish from scientists, as well as the public (Lund et al., 2007). Hopefully, this will lead to a shift in human attitudes towards fish, making them regarded as animals of moral concern, and no longer as crops to be harvested.

In the debate regarding fish sentience and the actual experiences fish have of pain, fear and suffering, authors (e.g. Ashley, 2007; Lund et al., 2007) advocates to give fish the benefit of the doubt. Since it is not possible for science to fully comprehend the subjective emotional states fish are capable of (nor of any other animal), to say that welfare issues

should be including fish seem to be the morally just approach. Lund and co-workers (2007) state that the lack of scientific evidence of fish sentience should not lead to the presumption that sentience is absent in fish, and make moral concern of fish insignificant. In addition the authors suggest that a risk assessment would speak in favour of giving fish the benefit of the doubt, since the risk of treating fish as sentient if not in case sentient is, morally, significantly smaller than treating fish as non-sentient if in fact sentient, considering the vast amount of fish reared in farms yearly and the severe implications for welfare of aquaculture practices.

In the wild, fish are exposed to a number of different stressors that can lead to suffering, including starvation, injuries as a result of predation, difficult environmental conditions, and diseases. In regard of this, it can be presumed that fish are accustomed, hence adapted, to handle stressful situations. According to Huntingford and co-workers (2006), this does not give humans the right to expose fish under human care to such situations, since there is a moral distinction between the impaired welfare caused by natural incidents or conditions, and that caused by humans, especially in human imposed confinement. The authors continue to define the difference by suggesting that the stressors fish are exposed to in natural conditions are often avoidable and of short duration, while human imposed stressors are often constant and unavoidable. This is objected to by Arlinghaus and co-workers (2007), through the view that humans are a part of nature that should interact with other species, and through the argument that for the individual fish, if stressors are imposed by humans or otherwise is of no importance. A possible response to these arguments could be that, like Huntingford and co-workers (2006) state, there is a difference in stressors that could be, at least, attempted to escape from, and stressors that are unavoidable because of confinement. Another argument could be that there is a difference for us humans if it is we who are exposing others of possible suffering, to if it is induced by natural elements. Furthermore, the stressors caused by human interaction with fish are often not instead of, but in addition to, nature invoked stressors, hence they should be of importance for the individual fish.

Arlinghaus and co-workers (2007) are further challenging the “emotion-centred feeling-based” view adopted by Huntingford and co-workers (2006), in terms of assessing fish welfare. They argue that since the science-based evidence of the mental states of fish is inconclusive, concepts such as suffering should not be applied to fish. Instead, they are advocating a function-based assessment of fish welfare, meaning that good welfare is when the fish is “in good health, with its biological system functioning properly and not being forced to respond beyond its capacity”. This would mean that good fish welfare could be acquired without allowing fish to perform their natural behaviours, and without possibility for fish to be in good mental states. Arlinghaus and co-workers (2007) continue by saying that since the individual experiences of fish is impossible to fully comprehend and measure, biologists should focus on signs of poor or good welfare; not the existence or non-existence of fish sentience, or ethical discussions of moral attitudes towards fish, because the focus of biologists should be on that what is objectively measurable. A different view is possible though, namely that of the significance of science to be a part of ethical discussions. Science should take its place in society, not as all-knowing, but as a voice in the on-going discourse that is essential for evolvment. Taking a stand and openness for possibilities should be encouraged in science, not disregarded as insincerity. Kaiser and Stead (2002) also acknowledge a need for scientists to be a voice in the debate by themselves, and are stating that participation in relevant discussions should be adopted

more widely. Although, or perhaps just because, research often is inconclusive, with more investigations required to prove hypothesis to be fact, a discussion of appropriate measures that should be taken on the base of currently available research is necessary.

Balcombe (2009) welcomes the approach of science to animal sentience research, in contrast to the conclusions of Arlinghaus and co-workers (2007). According to Balcombe (2009), research on the individual experiences of animals deserves serious regard, both in science and in animal ethics, since ethical deliberations require scientific bases. The author points out however, a lacking in current animal sentience related literature, i.e. the absence of the concept of animal pleasure. Instead, the focus on animal sentience research seems to be on negative states; such as pain, distress and suffering. Although Balcombe (2009) grants that the negative states are of higher moral significance, it does not mean that pleasure is a negligible aspect of animal welfare. This since there are numerous indicators in different groups of animals, that animals are willing to work for rewards (e.g. play, tasteful food, comfort, and touching or sexual behaviours), and not only to avoid pain- or stressful situations (reviewed in Balcombe, 2009). One example related to fish is the interaction between cleaner fish and their “clients”. The clients let the cleaner fish clean them from parasites, loose skin, and fungal infections; not for the reason of benefits in terms of health and survival, but because of the pleasure the clients get from the actual touch (reviewed in Balcombe, 2009). Fish, as well as other animals, kept in human imposed captivity, can be deprived of pleasure in two ways: by keeping them enclosed in impoverished conditions, unsatisfying their behavioural, mental, or physical needs; and by the killing of healthy animals, hence denial of the future pleasures a full life could give (Balcombe, 2009). In that sense, the concept of fish pleasure has moral significance and therefore should be treated as a vital aspect of animal welfare research and animal ethics discussions. Balcombe (2009) concludes that when viewing animals as motivated not merely by pain avoidance, but also by pleasure seeking, the human inflicted deprivation of freedom and taking of lives, for gains such as human entertainment or gastronomic matters, are in essence immoral practices.

Whether you are of the view that ethical treatment of animals means that animals have the right to their own lives and freedom, or of the view that it is morally just to kill animals for meat, but that they should be treated humane until and during slaughter, the review of current literature in the previous section clearly implies that the treatment of fish in aquaculture is at present not to be regarded as ethical.

## **6. ASPECTS OF ENVIRONMENTAL IMPACT**

The environmental impact that aquaculture means includes: habitat destruction; invasion of exotic species and pathogens; waste and chemical discharge; nutrient pollution; spreading of diseases; and further decline of wild fish populations, partly through the large amount of wild caught fish used as feed (e.g. Naylor et al., 2000; Costa-Pierce, 2003; Dempster & Holmer, 2009). In the following section, these issues will be briefly addressed.

### **6.1. Decline of wild fish populations**

Wild fish populations have declined rapidly, due to the over-fishing of fisheries, leading to the substantial growth of fish reared in aquaculture (Naylor et al., 2000). Although these fishes are reared and not a part of the wild fish populations, aquaculture is still partly responsible for the decline of wild fish in the world’s oceans (Naylor et al., 2000). This

actual negative influence on the status of fisheries' stocks (most are currently over or fully fished) is in contrast with the presumed effect of the rapid growth of aquaculture production, which was thought to be a decline in catches of wild fish (Naylor et al., 2000). To explain this, Naylor and co-workers (2000), point out that wild fish are being used as feed, in form of fish meal and fish oil, in aquaculture as well as other forms of meat production, and fish from intensive systems usually have been fed fish protein that equals 2-5 times the fish protein produced on-farm. When it comes to the ten most farmed omnivorous species, for 1 kg of farmed fish, it takes about 1.9 kg of wild caught fish, and the equivalent for the carnivorous species is on average 2.5-5 kg (Naylor et al., 2000). Carnivorous fish species are commonly farmed and are in the western society more demanded than herbivorous or omnivorous species, making the decline of wild fish populations accelerate (Naylor & Burke, 2005).

Besides using wild fish as feed for producing farmed fish, other factors associated with aquaculture have impacts on wild fish populations. These are in form of spreading of diseases and parasites, as well as genetic and ecological factors (Hansen & Windsor, 2006), which will be discussed below, both as threats to wild fish populations, and as threats to ecosystems in whole. For the preservation of threatened wild fish, the aquaculture industry needs to consider its environmental impact and improve its practices, which has not been the case hitherto (Hansen & Windsor, 2006). The drastic decline of wild fish in the world's oceans is not just a moral issue in itself, but also a critical environmental issue, since fish play an vital protective role of their ecosystems (Holmlund & Hammer, 1999). The services to ecosystems made by fish include: predation on other organisms; food web maintenance; regulation of sediment processes and flow of carbon and nutrients; and acting as feed for other aquatic, terrestrial, or aerial species, hence in addition providing services to other ecosystems than their own (Holmlund & Hammer, 1999).

## **6.2. Waste and chemical discharge**

There are various chemicals used in aquaculture practices that can affect the environment if discharged to surrounding water bodies, including different types of pesticides, fertilizers, disinfectants, antibiotics, and oxidants (Boyd & Massaut, 1999; Naylor & Burke, 2005). These can be harmful to natural waters through eutrophication from nutrients or pollution from heavy metals, as well as influencing through medicinal or toxic impact on wild aquatic animals (Boyd & Massaut, 1999; Naylor & Burke, 2005). The environmental impact of effluents is hard to assess however, considering the variety of different practices, species and ecosystems utilized in aquaculture, as well as the interaction between these variables (Fernandes et al., 2001).

Nitrogen, in form of nitrite and ammonia, that leaks out into nearby waters can be toxic to fish if in high levels (Naylor et al., 2000). Waste water with faeces and excess feed can lead to pollution, mostly in shallow waters and nearby coastal fish farms, in areas with restricted water exchange, or in areas with high occurrence of intensive aquaculture (Ervik et al., 1997; Naylor et al., 2000; Fernandes et al., 2001). Especially benthic environments under and around fish farms seem to be affected, through sedimentation of feed and faeces, which can lead to anaerobic conditions, methane or hydrogen sulphide exposure, pathogen spreading, and increased risk of bacteria resistance in case of antibiotics or other anti-bacterial agents additives in the feed (Ervik et al., 1997). The use of antibiotics has been

widespread in aquaculture, both legal and illegal such, and is likely to increase as farms grow more intensive and stocking densities increases (Garrett et al., 1997).

The poor water quality and crowding conditions in intense aquaculture may impair the immune system of fish, leaving them more receptive to diseases, thus more medical treatments are required (Haya et al., 2001). There are a number of different chemicals used in aquaculture, and there is little knowledge of the quantity of these chemicals used (Haya et al., 2001). Since the industry has grown rapidly the last decades, the usage of medicines and other chemicals (for example feed additives) are probably in many regions influencing the environment, the indigenous species and their habitats (Haya et al., 2001). One example of the consequences common fish farm medicines may have on the oceanic environment is the anti-parasitic drug used on Atlantic salmon in case of sea lice infestation. The pesticide is toxic, even lethal, to lobsters and shrimps that are subjected to it in high enough concentrations (Haya et al., 2001). Mortality of shrimps and lobsters were observed by Haya and co-workers (2001); half of the lobsters tested did not survive if exposed to the pesticide an hour in the concentrations being used in current bath treatment practices of salmon, and spawning failure was observed in both shrimps and lobsters tested. However, it is not certain how wild lobsters and shrimps are affected by the sea lice treatment of salmon, since the concentration of the pesticide will decrease when released into the surrounding waters, although mortality and reproductive failures are likely to increase (Haya et al., 2001).

### **6.3. Loss of habitat**

The usage of natural habitats for wild fish as sites for aquaculture can have detrimental effects for wild fish, as well as the environment. It is mainly coastal habitats that are being exploited. One example of this is the cultivation of mangrove forests for fish farming, leading to loss of nursing grounds and shelter for fish, and leading to sediment transport to downstream coral reefs (Naylor et al., 2000). This threatens the survival of a number of aquatic species, and has been reported to cause climate change through decline of rainfall (Garrett et al., 1997). Another effect aquaculture can have on habitats is that of clearing of land to make saltwater ponds. This increases the salinity of the ground, making it an irreversible alteration of the natural environment (Diana, 2009).

Along the coastline of the Pacific outside of Chile, aquaculture production of salmon has led to a loss of benthic biodiversity of an average of 50% on farm sites, due to sediment changes, generated by organic matter discharge, decreased oxygen levels, and copper deposition (reviewed in Buschmann et al., 2006). Hazardous algal blooms have also increased, as has the pollution of nitrogen from intensive salmon farms, giving raised mortality rates of aquatic species (Buschmann et al., 2006).

### **6.4. Spreading of diseases to wild fish populations**

Diseases spreading between wild and farmed fish are problematic from an environmental point of view in the sense of threats to fish populations. The spread of pathogens often originates from wild fish, though due to crowding and intensity, it can often turn into epidemics when spreading to fish farms (Naylor & Burke, 2005). In regions where aquaculture is common and concentrated, it is thought that diseases and parasitic infections that is relatively uncommon in natural waters, can spread more easily and become a severe hazard to wild fish (Naylor & Burke, 2005). Another source of spread seems to be

contaminated feed, in form of wild fish products that is imported from different parts of the world (Naylor & Burke, 2005), as well as interaction between infected escaped farmed fish with wild fish populations (Naylor et al., 2005). Interbreeding of wild and farmed fish may also pose a risk regarding disease spread, through weakening of the immune system of wild fish (see subsequent section), hence enabling spread and prevalence of pathogens and diseases (Naylor et al., 2005). For example, there now is evidence suggesting that current high levels of sea lice outbreak in wild fish populations is closely linked to the high concentration of aquaculture (reviewed in Krkošek et al., 2005; Naylor et al., 2005). Regarding salmon farms, spreading of parasites from farmed to wild fish mainly takes place during the juvenile stage of the wild fish, when they are most vulnerable due to their small size and recent transition to salt water conditions, and since they are, under natural conditions, spared from parasites associated with adult salmon (Krkošek et al., 2005). Moreover, the influence of a single salmon farm may result in a four to five fold increase in infection pressure, compared to natural levels, and if near a wild salmon migration route, can cause severe damage to wild populations through high mortality rates on juveniles (Krkošek et al., 2005; Krkošek et al., 2006), in addition to the welfare implications of sea lice infestation for the individual fish (Ashley, 2007). The effect seen on juvenile salmon is most likely not restricted to salmon; other species of fish might be at risk of similar parasitic development, with especial high risk for fish species with migratory behaviour, as aquaculture is expanding internationally (Krkošek et al., 2006).

Diseases spreading between farmed and wild populations of fish often occur, though it is problematic to assess these spreads due to their complexity (Naylor et al., 2000). To reduce disease spread from farmed to wild fish however, it is recommended to make serious attempts to decrease stress that can suppress the immune system in farmed fish, as well as careful filtering of used water before discharge (Naylor and Burke, 2005).

### **6.5. Invasion of exotic organisms**

It is not unusual that fish in aquaculture production escapes, becoming a part of the ecosystem (Naylor et al., 2000), either through frequent low scale escapes, or occasionally through mass escapes due to mostly weather conditions such as storms (Naylor et al., 2005). Effects of the ecosystem and wild fish populations include competition for food or mates with fish of the same or another species; predation; introduction of pathogens; and impaired fitness through interbreeding (Naylor & Burke, 2005; Streftaris et al., 2005). More than once, exotic species of fish have competed out the indigenous fish, resulting in exotic fish displacing other species, leading to a decrease in biodiversity (Diana, 2009). The most common type of farmed salmon, the Atlantic salmon, frequently escapes and constitutes about 40 % of the wild fish caught of this species in the North Atlantic Ocean (Naylor et al., 2000), and outnumbers Atlantic salmon with wild origin returning to streams by far (Naylor et al., 2005). Competition for food or mates between wild and farmed fish may have serious implications, especially since farmed Atlantic salmon are selected for fast growth, and unintentionally have become more aggressive and risk-taking than their wild counterparts, which means that wild salmon may be stressed by the competition (Naylor et al., 2005).

Wild Atlantic salmon have greater genetic variability than farmed ones, since much of the farmed salmon come from a small number of breeders (reviewed in Naylor et al., 2005). Interbreeding could alter the genetics of the native fish, making them less adapted to their

specific habitats and to environmental changes, and therefore further risk depletion (Naylor et al., 2000). Survival of hybrids between wild and feral salmon have lower survival rates than wild offspring, indicative of the effect interbreeding might have on wild populations (Naylor et al., 2005). Considering the amount of feral Atlantic salmon, the decline of wild populations, as well as the feral salmon's ability to compete, it is not unlikely that all wild salmon soon will have descended from farmed fish (Tufto & Hindar, 2003). This would mean a huge loss of genetic diversity that potentially will make Atlantic salmon unable to handle challenges such as changes in their environment (Naylor et al., 2005). Attempted breeding between different species is mainly unsuccessful to produce fertile offspring, thus not an issue in regards of genetic diversity, but may pose a risk to populations of fish that are declining in the wild through failed reproduction (Naylor et al., 2005). Introduction of transgenic fish in aquaculture would have an even larger impact of the concerned ecosystems, since transgenic fish have very dissimilar genetic make-up compared to their wild counterparts, making the genetic influence of escapees greater in case of successful interbreeding (Naylor & Burke, 2005).

The environmental, biological, and biodiversity aspects of introduction of exotic aquatic animals require legal framework to protect oceans from the negative effects of such introduction, especially since the scope of these effects are largely unknown to date (Streftaris et al., 2005). Although the uncertain effects, Diana (2009) acknowledges the introduction of exotic species or altered genotypes of indigenous species as, most likely, the largest aquaculture impact on aquatic biodiversity.

## **7. ETHICS IN CONCERN OF THE ENVIRONMENT**

One claim of supporters of aquaculture is that the growing industry results in more employment opportunities for local communities, which has to be taken into account when assessing the environmental implications of fish farm practices (Kaiser & Stead, 2002). However, this claim can be challenged, considering the advance in technology leading to more automated production, reducing the personnel required (Kaiser & Stead, 2002). In addition to this, there are uncertainties, according to Kaiser and Stead (2002), regarding the extent to which aquaculture can continue to expand, the questions being of environmental sustainability and public consent. It is not implausible that the same practices that are the threats to aquatic environment and wildlife may also essentially be the end of the industry itself.

The significant growth of aquaculture production to date, and the continuing increase of demand for fish products worldwide, indicate what a tremendous economic role the industry has and will continue to have, not least in developing countries (Diana, 2009). The developing countries produce a vast majority of the world's fish products, and the export of these products constitutes a substantial part of the economy of these countries (Diana, 2009). This makes aquaculture an industry of immense economic importance for countries that are struggling with poverty, which must be taken into account when assessing the ethics of that industry's practices. However, with wild fish populations declining, the world's climate changing, and sites of fish farms becoming damaged, the question is whether an industry that is dependent of captured wild fish as feed and a healthy environment for production, should be counted on, and invested in, for future economic gain. According to Diana (2009), as a result of the decline of wild fish populations, aquaculture will soon need to depend on plant protein for feed. Considering this, to assure

ability to feed the world's fast growing human population, by meanwhile considering the environmental impacts of food production, it could be claimed that investments in more energy and water efficient food production is the appropriate measure to be taken. In short, why rear fish and feed them plant protein that could be feeding humans directly, without the energy losses associated with rearing fish, as well as other animals, for meat?

It has been suggested that advances in technology and the international implementations of those is the answer to sustainable food production. This is challenged by Khare and Khare (2006), who are of the view that technology by itself cannot be the solution, and that ethical concerns, values and attitudes constitute the most important base for environmental protection. The recent decades' substantial increase in public interest in environmental ethics, is indicating that there is a shift in views of quality of life, which for many humans has started to mean a clean and secure environment, instead of a increased economical income per capita through usage of natural resources (Khare & Khare, 2006). Although science can provide the knowledge required to develop environmentally sound food production practices, ethics is necessary for giving the public as well as the legislators the urge to support these practices (Khare & Khare, 2006).

Although the public's interest in environmentally sustainable aquaculture is increasing, there is a gap between ethical attitudes and consumer behaviour, meaning that concern for the environment does not seem to lead to environmentally sound consumer choices, and that ethical consumption is applied only by a small part of the public (Verbeke et al., 2007). The reasons for this could be the lack of information and availability of environmentally sensible products, as well as marketing and familiarity of other products (Verbeke et al., 2007). So the question is: how to close this gap? The public is showing an increasing concern for food production, but is not generally implementing this in their consumption. There is clearly a difference between ethical concern in theory and in practice, though not meaning a lacking in importance of, and sincerity in, moral concerns. What this does indicate however, is that the moral views of the public should be a matter of policies and legislation, not of the individual choices of the consumer. Besides legislation regarding the production of food, comprehensible consumer information should be available, making it easier to implement individual moral standards.

Environmentally sound eating basically means eating low on the food chain, since vast energy and water losses are associated with rearing animals on feed such as grains or legumes, that are suitable for humans (Goodland, 1997). Rearing fish for human consumption with current practices also means, as discussed above, impact on natural waters through chemical and waste discharge, as well as damage to habitats. Goodland (1997) expresses the view that products from environmentally unsustainable agriculture should be highly taxed, hence not preferred by consumers, leading to a shift towards more sound food consumption.

## **8. CONCLUSION**

The research reviewed in this paper illustrates the current issues of ethical concern in aquaculture practices. They consist of animal welfare issues such as: 1) physical issues of pain, disease, immunosuppression, and skeletal malformations due to breeding, 2) behavioural issues of overcrowding, aggression, abnormal behaviour, and repression of natural swimming and social behaviour, and 3) stress issues of fear, exhaustion, food deprivation and starvation. Furthermore, they consist of environmental issues such as: 1)



decline of wild fish population due to using wild caught fish for feed, introduction of non-indigenous species, spreading of diseases, and habitat destruction, and 2) threats to the environment through chemical and waste discharge, and destruction of ecosystems.

Considering the challenges aquaculture would be facing, if attempting to resolve the issues of ethical concern, resolution seems essentially unattainable. The economical and nutritional gains of fish farms are not comparable with the vast environmental and animal welfare losses. In addition, the aquaculture industry has not been able to solve the issue of decline of wild fish populations, but is instead further exacerbating the decline. The ethical arguments and scientific evidences here reviewed have not all come to the same conclusions. What can be concluded, however, is that in spite of dissimilar views on animal and environmental ethics, or different results of empirical studies, the general agreement is that current aquaculture practices are neither meeting the needs of fish nor environment. Thus, it does not seem unreasonable to state the obvious environment and animal friendly aspects of a fish-free diet.

## 9. ACKNOWLEDGEMENT

I would like to thank Stefan Gunnarsson, Head of Department of Animal Environment and Health, Swedish University of Agricultural Sciences, for valuable advice, guidance and encouragement. I also thank Frida Lundmark, Guest teacher at Swedish University of Agricultural Sciences, for raising the important issues of fish welfare and ethics, and thus for the motivation behind this review.

## 10. REFERENCES

- Afonso, J. M., Montero, D., Robaina, L., Astorga, N., Izquierdo M.S., Ginés, R. (2000). *Association of a lordosis-scoliosis-kyphosis deformity in gilthead seabream (Sparus aurata) with family structure*. *Fish Physiology and Biochemistry* 22, 159–163.
- Arlinghaus, R., Cooke, S. J., Schwab, A., Cowx, I. G. (2007). *Fish welfare: a challenge to the feeling-based approach, with implications for recreational fishing*. *Fish and Fisheries* 8, 57–71.
- Ashley, P. J. (2007). *Fish welfare: Current issues in aquaculture*. *Applied Animal Behaviour Science* 104, 199–235.
- Balcombe, J. (2009). *Animal pleasure and its moral significance*. *Applied Animal Behaviour Science* 118, 208–216.
- Boyd, C. E., Massaut, L. (1999). *Risks associated with the use of chemicals in pond aquaculture*. *Aquacultural Engineering* 20, 113-132.
- Buschmann, A. H., Riquelme, V. A., Hernández-González, M. C., Varela, D., Jiménez, J. E., Henríquez, L. A., Vergara, P. A., Guíñez, R., Filún, L. (2006). *A review of the impacts of salmonid farming on marine coastal ecosystems in the southeast Pacific*. *Journal of Marine Science* 63, 1338-1345.
- Chandoo K. P., Duncan, I. J. H., Mocciaet, R. D. (2004). *Can fish suffer?: perspectives on sentience, pain, fear and stress*. *Applied Animal Behaviour Science* 86, 225–250.
- Costa-Pierce, B. A. (2003). *The 'Blue Revolution' - Aquaculture Must Go Green*. *World Aquaculture (in press)*, 1-5.
- Cottee, S. Y., Petersan, P. (2009). *Animal Welfare and Organic Aquaculture in Open Systems*. *Journal of Agricultural and Environmental Ethics* 22, 437–461.
- Dempster, T., Holmer, M. (2009). *Introducing the new Multidisciplinary journal*

- Aquaculture Environment Interactions*. *Aquaculture Environment Interactions* 1, 1-2.
- Diana, J. S. (2009). *Aquaculture Production and Biodiversity Conservation*. *BioScience* 59, 27-38.
- Donaldson, E. M. (1996). *Manipulation of reproduction in farmed fish*. *Animal Reproduction Science* 42, 381–392 .
- Ellis, T., North, B., Scott, A. P., Bromage, N. R., Porter M., Gadd, D. (2002). *The relationship between stocking density and welfare in farmed rainbow trout*. *Journal of Fish Biology* 61, 493–531.
- Ervik, A., Hansen, P. K., Aure, J., Stigebrandt, A., Johannessen, P., Jahnsen, T. (1997). *Regulating the local environmental impact of intensive marine fish farming : II. The concept of the MOM system (Modelling-Ongrowing fish farms-Monitoring)*. *Aquaculture* 158, 85-94.
- Fernandes, T. F., Eleftheriou, A., Ackefors, H., Eleftheriou, M., Ervik, A., Sanchez-Mata, A., Scanlon, T., White, P., Cochrane, S., Pearson, T. H., Read, P. A. (2001). *The scientific principles underlying the monitoring of the environmental impacts of aquaculture*. *Journal of Applied Ichthyology* 17, 181-193.
- Food and Agriculture Organization of the United Nations (2006). *Nearly half of all fish eaten today farmed, not caught*. FAO Newsroom, <http://www.fao.org/newsroom/en/news/2006/1000383/index.html>, Accessed 9 April 2010.
- Fraser, D., Weary, D. M., Pajor, E. A., Miligan, B. N. (1997). *A scientific conception of animal welfare that reflects ethical concerns*. *Animal Welfare* 6, 174–186.
- Gatica, M. C., Monti, G., Gallo, C., Knowles, T. G., Warriss, P. D. (2008). *Effects of well-boat transportation on the muscle pH and onset of rigor mortis in Atlantic salmon*. *Veterinary Record* 163, 111-116.
- Garrett, E. S., dos Santos, C. L., Jahncke, M. L. (1997). *Public, Animal, and Environmental Health Implications of Aquaculture*. *Emerging Infectious Diseases* 3, 453-457.
- Goodland, R. (1997). *Environmental sustainability in agriculture: diet matters*. *Ecological Economics* 23, 189-200.
- Grigorakis, K. (2009). *Ethical Issues in Aquaculture Production*. *Journal of Agricultural and Environmental Ethics* (Online DOI 10.1007/s10806-009-9210-5).
- Hansen, L. P., Windsor, M. L. (2006). *Interactions between Aquaculture and Wild Stocks of Atlantic Salmon and other Diadromous Fish Species: Science and Management, Challenges and Solutions*. *Journal of Marine Science* 63, 1159-1161.
- Haya, K., Burridge, L. E., Chang, B. D. (2001). *Environmental impact of chemical wastes produced by the salmon aquaculture industry*. *Journal of Marine Science* 58, 492–496.
- Holmlund, C. M., Hammer, M. (1999). *Ecosystem services generated by fish populations*. *Ecological Economics* 29, 253–268.
- Huntingford, F. A., Adams, C., Braithwaite, V. A., Kadri, S., Pottinger, T. G., Sandøe, P., Turnbull, J. F. (2006). *Current issues in fish welfare*. *Journal of Fish Biology* 68, 332–372.
- Håstein, T., Scarfe, A.D., Lund, V.L. (2005). *Science-based assessment of welfare: aquatic animals, Review of Science and Technology*. *Office International des Epizooties* 24, 529-547.
- Höglund, E., Jørgensen Bakke, M., Øverli, Ø., Winberg, S., Nilsson, G. E. (2005). *Suppression of aggressive behaviour in juvenile Atlantic cod (Gadus morhua) by l-tryptophan supplementation*. *Aquaculture* 249, 525– 531.
- Imslund, A. K., Foss, A., Koedijk, R., Folkvord, A., Stefansson S. O., Jonassen, T. M.

- (2006). *Short- and long-term differences in growth, feed conversion efficiency and deformities in juvenile Atlantic cod (Gadus morhua) started on rotifers or zooplankton*. *Aquaculture Research* 37, 1015-1027.
- Kaiser, M., Stead, S. M. (2002). *Uncertainties and values in European aquaculture: communication, management and policy issues in times of “changing public perceptions”*. *Aquaculture International* 10, 469–490.
- Khare, A., Khare, K. K. (2006). *Towards a Global Environmental Ethics - A Review Discussion*. *Asian Journal of Experimental Science* 20, 101-106.
- Krkošek, M., Lewis, M. A, Volpe, J. P. (2005). *Transmission dynamics of parasitic sea lice from farm to wild salmon*. *Proceedings of the Royal Society B* 272, 689-696.
- Krkošek, M., Lewis, M. A., Morton, A. Frazer, L. N., Volpe, J. P. (2006). *Epizootics of wild fish induced by farm fish*. *Proceedings of the National Academy of Sciences* 103, 15506–15510.
- Lambooj, E., Kloosterboer, R. J., Gerritzen, M. A., van de Vis, J. W. (2004). *Head-only electrical stunning and bleeding of African catfish (Clarias gariepinus): assessment of loss of consciousness*. *Animal Welfare* 13, 71-76.
- Latremouille, D. N. (2003). *Fin Erosion in Aquaculture and Natural Environments*. *Reviews in Fisheries Science* 11, 315-335.
- Lund, V., Mejdell, C. M., Röcklinsberg, H., Anthony, R., Håstein. T. (2007). *Expanding the moral circle: farmed fish as objects of moral concern*. *Diseases of Aquatic Organisms* 75, 109–118.
- Naylor, R., Hindar, K., Fleming, I. A., Goldburg, R., Williams, S., Volpe, J., Whoriskey, F., Eagle, J., Kelso, D., Mangel, M. (2005). *Fugitive Salmon: Assessing the Risks of Escaped Fish from Net-Pen Aquaculture*. *BioScience* 55, 427-437.
- Naylor, R., Burke, M. (2005). *Aquaculture and Ocean Resources: Raising Tigers of the Sea*. *Annual Review of Environment and Resources* 30, 185-218.
- Naylor, R. L., Goldburg, R. J., Primavera, J. H., Kautsky, N., Beveridge, M. C. M., Clay, J., Folke, C., Lubchenco, J., Mooney, H., Troell, M. (2000). *Effect of aquaculture on world fish supplies*. *Nature* 405, 1017-1024.
- Poli, B. M., Parisi, G., Scappini F., and Zampacavallo, G. (2005). *Fish welfare and quality as affected by pre-slaughter and slaughter management*. *Aquaculture International* 13, 29–49.
- Robb, D. H. F., Wotton, S. B., McKinstry, J. L., Sorensen, N. K., Kestin, S. C., Sorensen, N. K. (2000). *Commercial slaughter methods used on Atlantic salmon: determination of the onset of brain failure by electroencephalography*. *The Veterinary Record* 147, 298-303.
- St-Cyr, S., Aubin-Horth, N. (2009). *Integrative and genomics approaches to uncover the mechanistic base of fish behaviour and its diversity*. *Comparative Biochemistry and Physiology* 152, 9–21.
- Streftaris, N., Zenetos, A, Papatthanassiou, E. (2005). *Globalisation in Marine Ecosystems: the Story of Non-indigenous Marine Species Across European Seas*. *Oceanography and Marine Biology: An Annual Review* 43 , 419-453.
- Toranzo, A. E., Magariños, B., Romalde, J. L. (2005). *A review of the main bacterial fish diseases in mariculture systems*. *Aquaculture* 246, 37– 61.
- Tufto, J., Hindar, K. (2003). *Effective size in management and conservation of subdivided populations*. *Journal of Theoretical Biology* 222, 273–281.
- Turnbull, J., Bell, A., Adams, C., Bron, J., Huntingford, F. (2005). *Stocking density and welfare of cage farmed Atlantic salmon: application of a multivariate analysis*. *Aquaculture* 243, 121-132.

van der Vis, H., Kestin, S., Robb, D., Oehlenschläger, J., Lambooij, B., Münkner, W., Kuhlmann, H., Kloosterboer, K., Tejada, M., Huidobro, A., Otterå, H., Roth, B., Sørensen, N. K., Akse, L., Byrne, H., Nesvadba, P. (2003). *Is humane slaughter of fish possible for industry?*, *Aquaculture research* 34, 211-220.

Verbeke, W., Vanhonacker, F., Sioen, I., Van Camp, J., De Henauw, S. (2007). *Perceived importance of sustainability and ethics related to fish: a consumer behavior perspective*. *Ambio* 36, 580-585.

Verheijen, F. I., Flight, W. F. G. (1997). *Decapitation and brining: experimental tests show that after these commercial methods for slaughtering eel *Anguilla anguilla* (L.), death is not instantaneous*. *Aquaculture Research* 28, 361-366.

Yue, S., Moccia, R. D., Duncan I. J. H. (2004). *Investigating fear in domestic rainbow trout, *Oncorhynchus mykiss*, using an avoidance learning task*. *Applied Animal Behaviour Science* 87, 343–354.