

Sveriges lantbruksuniversitet Swedish University of Agricultural Sciences

Department of Wildlife, Fish, and Environmental Studies

Using camera traps to identify the influence of seasonal climate variations on the passage rates of a multi-species ungulate community in Öster Malma, Sweden

Nicholas Varley



Master's thesis • 30 credits Examensarbete/Master's thesis, 2019:18 Umeå 2019

Using camera traps to identify the influence of seasonal climate variations on the passage rates of a multi-species ungulate community in Öster Malma, Sweden

Nicholas Varley

Supervisor:	Tim Hofmee Department	Tim Hofmeester, Swedish University of Agricultural Sciences, Department of Wildlife, Fish, and Environmental Studies			
Assistant supervisor:	Joris Croms of Wildlife, I	Joris Cromsigt, Swedish University of Agricultural Sciences, Departmen of Wildlife, Fish, and Environmental Studies			
Assistant supervisor:	Sabine Pfef of Wildlife, I	ffer, Swedish University of Agricultural Sciences, Department Fish, and Environmental Studies			
Examiner:	Wiebke Nei Department	umann, Swedish University of Agricultural Sciences, t of Wildlife, Fish, and Environmental Studies			
Credits:		30 credits			
Level:		Second cycle, A2E			
Course title:		Master Thesis in Environmental science, A2E – Management of Fish and Wildlife Populations – Master's Programme			
Course code:		EX0936			
Course coordinating de	partment:	Department of Wildlife, Fish, and Environmental Studies			
Place of publication:		Umeå			
Year of publication:		2019			
Cover picture:		From camera traps			
Title of series:		Examensarbete/Master's thesis			
Part number:		2019:18			
Online publication:		https://stud.epsilon.slu.se			
Keywords:		ungulates, passage rates, temperature, snow depth, camera trap			

Swedish University of Agricultural Sciences Faculty of Forest Sciences Department of Wildlife, Fish, and Environmental Studies

Table of contents

1	Introduction	7
1.1	Hypotheses	9
2	Methods	10
2.1	Study Area	10
2.2	Site Characteristics	11
2.3	Camera Traps	12
2.4	Image Classification	13
2.5	Climate Data	14
2.6	Data Analysis	15
3	Results	17
3.1	Climate Variations	17
3.2	Monthly Passage Rates per Species	17
3.3	Monthly Passage Rates per Species & Temperature	18
3.4	Monthly Passage Rates per Species & Snow Depth	20
3.5	Effect of Temperature on Seasonal Passage Rates	22
4	Discussion	25
4.1	Effect of Climate Variations on Ungulate Passage Rates	25
4.2	Limitations	27
5	Conclusion	29
Refe	rences	30
Ackn	owledgements	35

1 Introduction

Climate change projections have consistently shown that globally, surface temperatures are expected to continue to rise with most observed warming trends being seasonal, the largest increases occurring in winter and spring (Jones et al., 2005). Europe is heading towards a future with much milder winters, hotter summers with less precipitation and an increase in the frequency and intensity of extreme weather events (Kelemen et al., 2009; IPCC, 2014). Flood risk is also predicted to increase due to snowmelt floods shifting from spring to winter as a result of a reduced snow season and less accumulation during winter months (Behrens et al., 2010).

The climate changes outlined in the latest IPCC report (IPCC, 2018) are set to have serious effects on the behaviour and distribution of a range of both flora and fauna. Studies into the relationship between climate change and ecology have shown that plant and animal species are facing changes to their current range due to new climate conditions, with many expected to shift poleward and to higher altitudes, in keeping with expanding climate boundaries (Walther et al., 2002). Native plant and tree species may also be threatened by competition with the range expanding non-native species that may be better adapted to the novel climatic conditions (Katona et al., 2013). Such changes in plant community may affect food availability for animals as climate change is causing an advance in phenology, whether that be the flowering of a plant or breeding of an animal (Visser and Both, 2005). The shift in phenology of a plant species, for example, could cause the whole ecological system to become mistimed, posing a serious threat to the fitness of other species within the system which may depend on it for their own survival and reproduction (Visser and Both, 2005).

Ungulate species are expanding throughout Europe due to a combination of factors. These include a greater focus on conservation, displacement or removal of competitors and predators, and hunting establishments introducing native and non-native species that may be more desirable to hunt (Apollonio et al., 2010). Land-use change is another important factor that has influenced ungulate populations in Europe as changes to forestry practices has increased habitat size and quality, as well as more intense agriculture increasing high-quality crops (Kuiters et al., 1996; Cutini et al., 2015). Supplementary feeding of ungulates has also been used as a management tool to improve winter survival rates, reduce forest and crop damage and attempt to reduce human-animal conflict (Felton et al., 2017). With an increase in warmer winters in northern Europe, it is possible that ungulate species will expand further north as higher latitudes become more hospitable for more species (Büntgen et al., 2017).

The expansion of multiple species would create a situation where native and nonnative species would occupy the same habitat areas forming novel species-rich ungulate communities. Where once there may have only been one or two ungulate species within a habitat range, one can increasingly find four to six species, increasing pressure on the system supporting these species (Putman et al., 2011; Pfeffer et al., 2018). Such an increase in community size can have positive and negative impacts on the wider ecosystem. While in some cases it could lead to overexploitation of food sources, damage to the plant community and greater competition (Stewart et al., 2011; Bowyer et al., 2016; Lecomte et al., 2016) in others, ungulates can act as 'ecosystem engineers', altering their habitat through changes to plant, soil and water conditions as a result of grazing, trampling, urination and excretion (Baruzzi and Krofel., 2017).

The main threat to Sweden in terms of climate change might occur in the form of warmer summers, milder winters and an increase in extreme events. Scandinavia has seen the strongest warming in Europe, particularly in winter, since the 1980s and climate projections have shown temperatures could rise by up to 3 °C by 2100 (IPCC, 2014; Eklund et al., 2015). Currently, there is a gap in the knowledge of how changing climatic conditions such as variations in temperature and snow depth, affect the behaviour and movement of northern ungulates. Previous studies have focused to understand the relationship between a changing climate and topics such as supplementary feeding of ungulates, the influence of hunting on habitat selection, and moose-vehicle collisions (Seiler, 2004; Felton et al., 2017; Niemi et al., 2017).

Understanding the impact climate has on ungulate behaviour is important for their survival as it allows identification of any ways in which ungulates might be affected by future climate change.

In this study, I have investigated the relationship between ungulate patch use and climate conditions over monthly and seasonal timescales in southern Sweden for a three year period from September 2015 to September 2018 using camera traps.

1.1 Hypotheses

I hypothesised that (I) as average monthly snow depth increases, monthly ungulate passage rates will decrease as snow limits movement of ungulates and covers ground forage. As well as this, I expect that (II) passage rates will decrease during hot months as ungulates choose to stay in shaded areas to maintain a lower body temperature than if they were moving through patches.

2 Methods

2.1 Study Area

The study area 'Öster Malma' is located in the province of Södermanland, South-East Sweden, within the surrounding area of the city of Nyköping (Figure 1). The area has a temperate climate with average summer temperatures between 15.4 °C and 17.2 °C and mean winter temperatures between -0.6 °C and -2.6 °C (Climatedata.org, 2015). Precipitation varies (including rain and snow) with a minimum of 27 mm of precipitation in March and a maximum of 66 mm in August (Climatedata.org, 2015). There are five ungulate species present in the study area, which include moose, fallow deer, roe deer, red deer and wild boar (*Sus scrofa*).



Figure 1: Location of the study area 'Öster Malma', Sweden with its ten study sites marked in red (Ånöstam, 2017).

2.2 Site Characteristics

The study area consists of ten individual sites (Table 1). Nine of the ten sites were clear cut in 2014 and replanted with Norway spruce (*Picea abies*) or Scots pine (*Pinus sylvestris*) in 2015. Horn was the only site that was clear cut in 2013 and naturally regenerated with pine. The experiment (see description below) started in autumn 2015 and is still ongoing. Today some sites are dominated by one or two tree species, mainly spruce and pine, while others have much more species diversity, including silver birch (*Betula pendula*), downy birch (*Betula pubescens*), aspen (*Populus tremula*), willow (*Salix*) and rowan (*Sorbus*).

Site	Latitude	Longitude
Elghammar	59.06579	-17.0917
Grundsdal	59.00261	-17.2265
Horn	58.73905	-17.138
Jakobsberg	59.00117	-17.1395
Klippan	58.95803	-17.1809
Kristineholm	58.8511	-16.9143
Marö	58.93267	-17.3262
Nygård	58.96657	-17.1577
Trollesund	58.91737	-17.163
Vibyholm	58.97424	-16.9005

Table 1: Latitudes and longitudes of the ten sites.

Each of the ten sites consists of four exclosures that measure 14x14 m. All exclosures were set up in 2015 before the experiment began. The exclosures have four different treatment conditions: control, closed, summer, and winter. The control exclosures are open all year and are therefore constantly accessible by all animals. The closed exclosures are closed and fenced off all year to prevent animals from entering and accessing the vegetation within them. Summer exclosures are closed and fenced off during the vegetation period (April to October) only and are then opened during the months October to April. The winter exclosures are the opposite and are therefore closed during the non-vegetation period (October to April) and opened during the vegetation period.

2.3 Camera Traps

Each control exclosure is permanently equipped with a camera trap (Reconyx Hyperfire HC500) in the south-east corner all year round. A second camera trap is shared between the summer and winter exclosures, being switched interchangeably during opening months. The cameras are set up to monitor the wildlife using the exclosures with a focus on the four ungulate species; moose, red deer, fallow deer, and roe deer. As the closed exclosure is permanently free from animals, there is no camera covering that area. The cameras have been in place and continuously taking photographs since they were first installed at the end of September 2015 and are checked regularly to ensure that they are still operating correctly and have sufficient battery power.

A passive infrared sensor triggers the cameras, when the sensor registers a difference in thermal infrared between moving objects and their background in the detection zone, such as the temperature of the background vegetation or soil and a moving animal (Welbourne et al., 2016). Once triggered, the camera takes a series of three images at one-second intervals to capture the event detected (see for an example, Figure 2). Each camera also records a time-lapse image per day at 13:00 to check the camera is still functioning.



Figure 2: Example of a trigger event from a camera on-site showing a female fallow deer foraging on a grass species.

2.4 Image Classification

Images captured by the camera traps were classified to create a data set of the animals seen in the area. I classified images taken from September 2017 to September 2018 from eight out of the ten sites (Nygård & Vibyholm were excluded due to time constraints). Images taken between September 2015 and September 2017 were previously classified and therefore included in my analysis (excluding Nygård and Vibyholm). The classification process involved identifying a number of variables including the species present, the number of animals present per species and whether individuals were foraging. Foraging was identified where an animal had its head to the ground or could clearly be seen eating leaves from plants or trees. Foraging behaviour was not separated by species, therefore if there was an image containing two species where one was foraging but the other wasn't, it was classified as a foraging event. Anything else was marked as 'other behaviour'. Species identification could be done due to obvious differences in the appearance of the four ungulate species in question, such as body size and shape. In some instances it was not possible to identify the species present due to blurred images or very dark night images, therefore the animal present was recorded as 'unknown species' and excluded from the analyses. Other details such as gender, age and male antler points were also recorded. However, these characteristics were not relevant in this study and thus have not been included here for further analysis.

For my analysis, I used passage events as a measure of patch use. A single passage event represents a sequence of continuous pictures triggered by an individual or group of animals. To determine the end of one sequence and the start of a new sequence, I implemented a time gap of 5 mins between two triggers. If the second trigger was taken >5 mins after the initial sequence of images, it was deemed to be a separate passage event.

2.5 Climate Data

I extracted data for daily temperature and snow depth from the webpage of the Swedish Meteorological Hydrological Institute (SMHI, and https://www.smhi.se/data/meteorologi/ladda-ner-meteorologiska-observationer#param=airtemperatureInstant,stations=all). Since there has been no weather station located directly in the study area, I extracted daily temperature data from three weather stations surrounding Öster Malma; Floda A, Södertälje and Oxelösund (Figure 3). Based on the data of these three stations, I calculated an average daily temperature for the whole study area. These average daily temperatures were used to extract average values per month and season. Snow depth data was only available from one weather station around Öster Malma, Södertälje (Figure 3). Therefore, I used the snow depth data from this individual station only to calculate monthly and seasonal averages for the study area. To define a season I used each seasons' start and end date (SMHI, https://www.smhi.se/en/weather/swedenweather/season-map/, see also Table 2) and rounded them to receive full months. Rounding to months was done due to a delay in finding the exact start and end dates of each season.

Season	Months
Spring	March/April/May
Summer	June/July/August
Autumn	September/October/November
Winter	December/January/February

Table 2: Definition of which months form each season.



Figure 3: Location of the three SMHI weather stations. Weather stations used for temperature data are marked with red spots while snow depth data is marked with a blue spot (Google Maps., 2019).

2.6 Data Analysis

I used the data on passage events to calculate a species and site-specific daily index (passage rate per day) as the sum of passages per species, per site, per day. Similarly, I calculated a monthly and seasonal passage rate per species by taking a sum of all passage events divided by the number of days covering those specific time periods, for example, the fallow deer passage rate in June 2018 would be the sum of fallow deer passage events divided by 30 days. I repeated this step for every month between September 2015 and September 2018 and similarly each season from autumn 2015 to autumn 2018. This allowed for a monthly and seasonal passage rate to be calculated for each species for the whole study period as seen below:

Passage rate = number of passages/number of days camera was active (per month or season)

I first used this dataset to run linear regression tests to identify any relationship between the passage rates of each species separately and the climate variables. My dependent variables were monthly and seasonal passage rates of the four species, where I ran separate models for the monthly and seasonal variables, with two covariates, temperature and snow depth. I then ran general linear mixed effect models in which the passage rate was always the dependent variable. In the initial models, I used the passage rates of all four ungulate species together. I then ran separate models for each species individually to identify species-specific responses to the climate variables, using the passage rate of each individual species as the dependent variable. I included ungulate species as a fixed effect, temperature, temperature² and snow cover as covariates, and site and year as random factors. Again, I ran separate models for monthly and seasonal response variables. I performed all statistical analyses in SPSS Statistics 25.

3 Results

3.1 Climate Variations

Winter 2016/17 was on average approximately 3 °C warmer than the winters of 2015/16 and 2017/18 (Figure 4). The summer of 2018 was on average approximately 3 °C warmer than the two previous summers (Figure 4). Snow depth in the study area varied across the three year study period with winter 2016/17 having the lowest snow depth and winter 2017/18 having the highest snow depth (Figure 4). Summer months always had a snow depth of 0 m, while during spring months (usually in March) there was a very small amount of snow present before it melted to 0 m (Figure 4).



Figure 4: Average monthly temperature variations (in red) and average monthly snow depth variations (in blue) from September 2015 to September 2018 with standard error.

3.2 Monthly Passage Rates per Species

During the study period, fallow deer and roe deer strongly dominate the visits to the sites, followed by red deer and then moose. The passage rates for all four species, and in all years, dropped strongly during mid-winter. Overall, there appears to be a

general trend towards passage rates of fallow deer and roe deer declining slightly over the years. There also seems to be a general trend towards moose and red deer passage rates increasing slightly over the same time period.





3.3 Monthly Passage Rates per Species and Temperature

Temperature and temperature² had a strong effect on the dependent variable, monthly passage rates (p = <.001) (Table 3). Ungulate species also had a significant effect on the dependent variable (p = .002). However year and site, the two random factors did not have a significant effect on monthly passage rates (p = .169; p = .732) (Table 3). Among the four species, roe deer had the highest R² value (0.3817) (Table 4). When looking at the influence of temperatures on the monthly passage rates of the four species individually, it can be seen that temperature had a positive effect on the monthly passage rates of roe deer (estimate = 0.005; p = <.001) (Table 4). Moose, red deer and fallow deer however, were not significantly influenced by temperature (estimate = <.001, p = .124; estimate = .001, p = .235; estimate = .001, p = .616) (Table 4).



Figure 6: Monthly passage rate of moose (blue), red deer (orange), fallow deer (green), and roe deer (yellow) in relation to temperature. Dotted lines indicate polynomial regressions.

Table 3: Univariate statis	tical analysis for t	the effect of tempera	<i>uture, temperature²,</i>
species, year and site, on t	the dependent vari	able, monthly ungu	late passage rates.

Explanatory Variables		df	f	р
Temperature	Hypothesis	1	32.310	.000
	Error	1010		
Temperature ²	Hypothesis	1	12.687	.000
	Error	1010		
Ungulate Species	Hypothesis	3	6.538	.002
	Error	22.236		
Year	Hypothesis	3	2.033	.169
	Error	10.755		
Site	Hypothesis	7	.623	.732
	Error	22.374		

Species	Estimates	Standard	t	р	R ²
		Error			
Moose	.000	.000	1.542	.124	0.0908
Red Deer	.001	.001	1.190	.235	0.1215
Fallow Deer	.001	.002	.502	.616	0.1299
Roe Deer	.005	.001	3.846	.000	0.3817

Table 4: Regression coefficient estimates, standard error, significance (p) and R^2 for the effect of temperature on the monthly passage rates per species.

3.4 Monthly Passage Rates per Species and Snow Depth

Snow depth had a strong effect on the dependent variable, monthly passage rates (p = <.001) (Table 5). Ungulate species also had a significant effect on the dependent variable (p = .002). Year and site did not have a significant effect on monthly passage rates (p = .197; p = .732) (Table 5). Fallow deer had the highest R² value out of the four species (0.3214) (Table 6). The individual analysis of the effect of snow depth on the monthly passage rate of each species shows that snow depth had a strong effect on fallow deer monthly passages (estimate = -1.289, p = .001) (Table 6). Moose, red deer and roe deer were not significantly influenced by snow depth (estimate = .004, p = .952; estimate = -.202, p = .342; estimate = -.354, p = .249) (Table 6).



Figure 7: Monthly passage rate of moose (blue), red deer (orange), fallow deer (green), and roe deer (yellow) in relation to snow depth.

Table	5:	Univa	riate	statistica	l analy	sis for	the	effect	of sn	<i>9</i> W	depth,	ungulat	e spe-
cies, y	vear,	and	site or	1 the dep	endent	variab	le, n	10nthl	y ung	ula	te pass	age rate	zs.

Monthly Passag	jes	Df	f	р
Snow Depth	Hypothesis	1	68.754	.000
	Error	7.224		
Ungulate Species	Hypothesis	3	6.538	.002
	Error	22.236		
Year	Hypothesis	3	1.865	.197
	Error	10.409		
Site	Hypothesis	7	.622	.732
	Error	22.379		

Standard R² Species Estimates t р Error Moose .004 .060 .952 0.045 .072 Red Deer -.202 .212 -.951 .342 0.1332 -1.289 .392 -3.285 0.3214 Fallow Deer .001 Roe Deer -.354 .306 -1.156 .249 0.2394

Table 6: Regression coefficient estimates, standard error, significance (p) and R^2 for the effect of snow depth on the monthly passage rates per species.

3.5 Effect of Temperature on Seasonal Passage Rates

On a seasonal time scale temperature, temperature² and snow depth all have a strong effect on ungulate passage rates (p = .001; p = .015; p = <.001) (Tables 7 & 8). In the temperature model ungulate species is also significant (p = .003) (Table 6). Neither year nor site, the two random factors, were statistically significant (p = .553; p = .679) (Table 7). Similarly, in the snow depth model, ungulate species was statistically significant while year and site were not statistically significant (p = .003; p = .533; p = <.680) (Table 8).

Individually, none of the four species' seasonal passage rates are significantly influenced by temperature (Table 9). There is a strong effect of snow depth on fallow deer seasonal passage rates (estimate = -2.735, p = .019) (Table 10). There is no effect of snow depth on the seasonal passage rates of moose, red deer or roe deer (p = .334; p = .892; p = .373) (Table 10).

Seasonal Passag	ges	df	f	р
Temperature	Hypothesis	1	11.613	.001
	Error	274		
Temperature ²	Hypothesis	1	9.951	.015
	Error	274		
Ungulate Species	Hypothesis	3	6.437	.003
	Error	22.582		
Year	Hypothesis	3	.763	.533
	Error	14.125		
Site	Hypothesis	7	.691	.679
	Error	24.520		

Table 7: Univariate statistical analysis for the effect of temperature, temperature², ungulate species, year and site on the dependent variable, seasonal ungulate passage rates.

Table 8: Univariate statistical analysis for the effect of snow depth, ungulate species, year and site, on the dependent variable, seasonal ungulate passage rates.

Seasonal Passages		df	f	р
Snow Depth	Hypothesis	1	27.562	.000
	Error	275		
Ungulate Species	Hypothesis	3	6.437	.003
	Error	22.582		
Year	Hypothesis	3	.765	.533
	Error	13.313		
Site	Hypothesis	7	.690	.680
	Error	24.552		

Table 9: Regression coefficient estimates, standard error, significance (p) and R^2 for the effect of temperature on the seasonal passage rates per species.

Species	Estimates	Standard	t	р
		Error		
Moose	.001	.001	1.810	.073
Red Deer	.002	.002	.835	.406
Fallow Deer	003	.003	891	.375
Roe Deer	.004	.003	1.433	.155

~			<u> </u>	A	A
	Species	Estimates	Standard	t	р
			Error		
	Moose	.206	.213	.970	.334
	Red Deer	096	.708	136	.892
	Fallow Deer	-2.735	1.150	-2.378	.019
	Roe Deer	750	.839	894	.373

Table 10: Regression coefficient estimates, standard error, significance (p) and R^2 for the effect of snow depth on the seasonal passage rates per species.

4 Discussion

4.1 Effect of Climate Variations on Ungulate Passage Rates

I carried out this study to understand the influence climate has on the patch use of a multi-species ungulate community in the south of Sweden. As Sweden's ungulate communities are expanding in both density and range it is important to understand how the climate affects their behaviour. With climate projections showing a strong likelihood of future warming, up to 3 °C warmer by 2100 (IPCC, 2014), understand-ing the response of ungulates to these factors is important. I investigated the effect of climate on ungulate patch use by using data collected from camera traps to assess whether climate changes were having any influence on ungulate passage rates in southern Sweden. The outcome of this study is that temperature only had a strong effect on the monthly passage rates of roe deer, suggesting that as temperature increases, roe deer visitation increases. Snow depth only had a strong effect on the monthly and seasonal passage rates of fallow deer. Fallow deer passage rates were negatively affected over a monthly and seasonal scale in relation to snow depth, which suggests that their visitation rates decline as snow depth increases.

Hypothesis I, that as average monthly snow depth increases, monthly ungulate passage rates will decrease, is only supported by fallow deer as the estimates show that the effect of snow depth on them was found to be significant, their monthly and seasonal passage rates decreasing as snow depth increased (Tables 6 and 10). The relationship between increasing snow depth and decreasing fallow deer passage rates could be caused by the snow covering the field layer forage therefore covering a major food source for the fallow deer. Supplementary feeding sites could also influence passage rates as supplementary feeding has been found to maintain higher densities of fallow deer in areas around feeding sites than would normally occur (Felton et al., 2017), therefore suggesting that feeding sites are an important determinant in fallow deer habitat patch selection.

The reduction in passage rates as snow depth increases has also been reported in other studies, especially in larger ungulates as the movement in snow requires a greater expenditure of energy (Rivrud et al., 2010). Moose, in particular, have been previously found to have their movement limited by snow cover (van Beest et al., 2011) however my results do not support this. Again this could be due to the low density of moose in the area giving a poor representation of how patch use of moose in Sweden is influenced by climate variations.

Hypothesis II, that passage rates will decrease during hot months, was not supported by these results. Over the years moose and red deer summer passage rates increased while there was a general trend towards decreasing summer passage rates for fallow deer and roe deer (Figure 5). This differs to other studies into the effect of temperature no movement, which have found that high summer temperatures have a negative effect on the movement of cervids (Hayes and Krausman., 1993; Rivrud et al., 2010). A possible explanation for moose and red deer summer month passage rates increasing during the study period could be related to the set-up of the study area. When the ten sites were clear cut and replanted in 2014/2015, grasses and small trees were the dominant species. This would have suited the smaller species, fallow deer and roe deer, as their dietary preferences are grasses, herbs and small trees. However, during the following years, the trees have grown taller with bigger branches, therefore being more favourable for the larger species, moose and red deer. This would mean that there was higher forage availability for moose and red deer in summer 2018.

4.2 Limitations

A major limitation of this study is the short time period during which the camera traps were taking images at the ten sites. Three years of continuous data was used as this is what was available to me at the time. However, when trying to assess the influence of weather on another variable, in this case, ungulate passage rates, it would be preferential to have at least 30 years of data as climate often refers to average weather patterns during the previous 30 years (NASA, 2017).

The snow depth data is also a limiting factor to this study. Only one weather station around the study area, Södertälje, had snow depth data available for the time period covering this study. This data is collected manually by SMHI either with a ruler, or fixed measuring stick (SMHI, 2013). This measurement is also not always carried out daily. If this is the case then the snow depth from the nearest weather station also recording snow depth is taken. This leads to the possibility of erroneous readings through human error as well as having to rely on the readings of different weather stations that is potentially tens of kilometres away from the study area. The other issue with snow depth data is that snow depth may vary significantly across small scales as it is influenced by the topography of the area, for example, slope or canopy cover (Zheng et al., 2016). As this is not uniform across the study area there will be areas which are deeper or shallower than the SMHI measurements.

Regarding image classification, a limitation arises due to the fact that it has been carried out by at least three different people, including myself. This could lead to human errors such as the misidentification of a species or misinterpretation of behaviour such as foraging (Young et al., 2018). While this is not likely to lead to significant errors that would be detrimental to the study, it is something that future studies should take into consideration. The image classification should be done consistently by the same individual to remove the human error.

The camera traps work with a passive infrared sensor which detects heat differences between the background and a moving object (such as an animal). On extremely hot days the background, such as vegetation and rocks, may heat up to the point where there is no detectable difference between the background and an animal walking in front of the camera (van Berkel, 2014). This is more of an issue with smaller species as they emit less body heat than larger species. Detection distance is another issue related to body size as detection of smaller species could decrease compared to the larger species the further away from the camera they are (Apps and McNutt, 2018). This would, therefore, mean that some passage events could be missed on an extremely hot day and therefore affect the outcome of the study by giving a false representation of the hottest days. Detectability and passage rates can also be affected by differences in insulation of summer and winter fur. As ungulates grow a winter coat to better insulate themselves during the colder months, there will be a difference in their heat signature in summer and winter, thereby leading to the possibility of being less detectable by the sensor in the camera trap (Hofmeester et al., 2019). This could mean that during winter, the ungulates are more detectable than in summer as there will be a bigger difference in the background temperature and their body temperature which is increased due to better-insulated fur.

5 Conclusion

Understanding the effect of climate variations on ungulate patch use can help to develop management strategies and further our knowledge on how climate change could affect multi-species ungulate communities. Here, I have shown that temperature and snow depth may have an effect on ungulate passage rates but that effects varied widely among species. Temperature was found to strongly affect the monthly passage rates of roe deer while snow depth was found to strongly affect both the monthly and seasonal passage rates of fallow deer. While climate variations have been found to have an effect on ungulate passage rates, it is more likely that other factors had more of an influence on changes to ungulate passage rates, such the setup of the study site and forage availability. After being clear cut and replanted, the smaller species (fallow deer and roe deer) initially benefitted from the vegetation available, However as the tree species in the sites have grown they are now more desirable for the larger species, moose and red deer. While some strong effects of both temperature and snow depth were found in my study there is still a lot of uncertainty and further studies should be carried out to better understand how climate variations affect ungulate patch use.

References

Ånöstam, F., 2017. Timing of ungulate browsing and its effect on sapling height and the field layer vegetation. *Examensarbete i ämnet biologi / Sveriges lantbruksuniversitet, Institutionen för vilt, fisk och miljö*, 2017(4), pp.1-28.

Apollonio, M., Andersen, R. and Putman, R. eds., 2010. European ungulates and their management in the 21st century. Cambridge University Press.

Apps, P.J. and McNutt, J. W., 2018. How camera traps work and how to work them. *African Journal of Ecology*, 56, pp.702-709.

Baruzzi, C. and Krofel, M., 2017. Friends or foes? Importance of wild ungulates as ecosystem engineers for amphibian communities. *North-Western Journal of Zoology*, 13(2), pp.320-325.

Behrens, A., Georgiev, A. and Carraro, M., 2010. FUTURE IMPACTS OF CLI-MATE CHANGE ACROSS EUROPE. *CEPS WORKING DOCUMENT*, 324, pp.1-27.

Bowyer, R.T., Bleich, V.C., Stewart, K.M., Whiting, J.C. and Monteith, K.L., 2014. Density dependence in ungulates: a review of causes, and concepts with some clarifications. *California Fish and Game*, 100(3), pp.550-572.

Büntgen, U., Greuter, L., Bollmann, K., Jenny, H., Liebhold, A., Galván, J.D., Stenseth, N.C., Andrew, C. and Mysterud, A., 2017. Elevational range shifts in four mountain ungulate species from the Swiss Alps. *Ecosphere*, 8(4), p.1761.

Climate-Data.org. 2015. CLIMATE NYKÖPING. (Online) Available at: https://en.climate-data.org/europe/sweden/soedermanlands-laen/nykoeping-19/.

Cutini, A., Chianucci, F. and Apollonio, M., 2015. WILD UNGULATES AND FORESTS IN EUROPE: INSIGHTS FROM LONG TERM STUDIES IN CEN-

TRAL ITALY. Atti del II Congresso Internazionale di Selvicoltura. Progettare il futuro per il settore forestale, Firenze, 26-29 novembre 2014. Firenze: Accademia Italiana di Scienze Forestali, 1, pp.509-517.van

Eklund, A., Axén Mårtensson, J., Bergström, S., Björck, E., Dahné, J., Lindström, L., Olsson, J., Simonsson, L. and Sjökvist, E., 2015. Sveriges framtida klimat: Underlag till Dricksvattenutredningen. SMHI.

Felton, A.M., Felton, A., Cromsigt, J.P., Edenius, L., Malmsten, J. and Wam, H.K., 2017. Interactions between ungulates, forests, and supplementary feeding: the role of nutritional balancing in determining outcomes. *Mammal Research*, 62(1), pp.1-7.

Google Maps, 2019, Öster Malma, Nyköping, Sweden (Online). Available at: https://www.google.co.uk/maps/place/%C3%96ster+Malma/@58.9323978,16.990 4486,9.88z/data=!4m8!3m7!1s0x465f2496792779d9:0xcb88af3d4cd3fd6d!5m2!4 m1!1i2!8m2!3d58.950833!4d17.158055.

Hayes, C.L. and Krausman, P.R., 1993. Nocturnal Activity of Female Desert Mule Deer. *Journal of Wildlife Management*, 57, pp.897-904.

Hofmeester, T., Cromsigt, J.P.G.M., Odden, J., Andrén, H., Kindberg, J. and Linnell, J.D.C., 2019. Framing pictures: A conceptual framework to identify and correct for biases in detection probability of camera traps enabling multi-species comparison. *Ecology and Evolution*, 2019, pp.1-17.

IPCC, 2014: Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Core Writing Team, Pachauri, R.K. and Meyer, L.A. (eds.) (p. 151). IPCC, Geneva, Switzerland.

IPCC, 2018: Summary for Policymakers. In: Global warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty. Masson-Delmotte, V., Zhai, P., Pörtner, H.O., Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J.B.R., Chen, Y., Zhou, X., Gomis, M.I., Lonnoy, E., Maycock, T., Tignor, M. and Waterfield, T. (eds.) (p. 32). IPCC, Geneva, Switzerland.

Jones, G.V., White, M.A., Cooper, O.R. and Storchmann, K., 2005. CLIMATE CHANGE AND GLOBAL WINE QUALITY. *Climate Change*, 73, pp.319-343.

Katona, K., Kiss, M., Bleier, N., Székely, J., Nyeste, M., Kovács, V., Terhes, A., Fodor, Á., Olajos, T., Rasztovits, E. and Szemethy, L., 2013. Ungulate browsing

shapes climate change impacts on forest biodiversity in Hungary. *Biodiversity and conservation*, 22(5), pp.1167-1180.

Kelemen, A., Munch, W., Poelman, H., Gakova, Z., Dijkstra, L. and Torighelli, B., 2009. Regions 2020 The Climate Change Challenge for European Regions. *European Commission Directorate General for Regional Policy*, 1, pp.1-27.

Kuiters, A.T., Mohren, G.M.J. and van Wieren, S.E., 1996. Ungulates in temperate forest ecosystems. *Forest Ecology and Management*, 88, pp.1-5.

Lecomte, X., Fedriani, J.M., Caldeira, M.C., Clemente, A.S., Olmi, A. and Bugalho, M.N., 2016. Too Many Is Too Bad: Long-Term Net Negative Effects of High Density Ungulate Populations on a Dominant Mediterranean Shrub. *PLoS ONE*, 11(7), pp.1-14.F

NASA. 2017. NASA - What's the Difference Between Weather and Climate? (Online) Available at: <u>https://www.nasa.gov/mission_pages/noaa-n/climate/climate_weather.html</u>.

Niemi, M., Rolandsen, C.M., Neumann, W., Kukko, T., Tiilikainen, R., Pusenius, J., Solberg, E.J. and Ericsson, G., 2017. Temporal patterns of moose-vehicle collisions with and without personal injuries. *Accident Analysis & Prevention*, 98, pp.167-173.

Pfeffer, S.E., Spitzer, R., Allen, A.M., Hofmeester, T.R., Ericsson, G., Widemo, F., Singh, N.J. and Cromsigt, J.P., 2018. Pictures or pellets? Comparing camera trapping and dung counts as methods for estimating population densities of ungulates. *Remote Sensing in Ecology and Conservation*, 4(2), pp.173-183.

Putman, R., Apollonio, M. and Andersen, R. eds., 2011. Ungulate management in Europe: problems and practices. Cambridge University Press.

Rivrud, I., Loe, L.E. and Mysterud, A., 2010. How does local weather predict red deer home range size at different temporal scales? *Journal of Animal Ecology*, 79, pp.1280-1295.

Seiler, A., 2004. Trends and spatial patterns in ungulate-vehicle collisions in Sweden. *Wildlife Biology*, 10(4), pp.301-313.

SMHI. 2013. Hur mäts snödjup? (Online) Available at: <u>https://www.smhi.se/kun-skapsbanken/meteorologi/hur-mats-snodjup-1.27291</u>.

SMHI. 2019. Ladda ner meteorologiska observationer. (Online) Available at: https://www.smhi.se/data/meteorologi/ladda-ner-meteorologiska-observa-

tioner#param=airtemperatureInstant,stations=all.

SMHI. 2019. Season Map. (Online) Available at:

https://www.smhi.se/en/weather/sweden-weather/season-map/.

Stewart, K.M., Bowyer, R.T., Dick, B.L. and Kie, J.G., 2011. Effects of density dependence on diet composition of North American elk Cervus elaphus and mule deer Odocoileus hemionus: an experimental manipulation. *Wildlife Biology*, 17, pp.417-430.

van Beest, F.M., Rivrud, I.M., Loe, L.E., Milner, J.M. and Mysterud, A., 2011. What determines variation in home range size across spatiotemporal scales in a large browsing herbivore? *Journal of Animal Ecology*, 80(4), pp.771-785.

van Berkel, T., 2014. Camera Trapping for Wildlife Conservation. *Expedition Field Techniques*, 1, pp.1-122.

Visser, M.E. and Both, C., 2005. Shifts in phenology due to global climate change: the need for a yardstick. *Proceedings of the Royal Society B: Biological Sciences*, 272(1581), p.2561.

Walther, G.R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J., Fromentin, J.M., Hoegh-Guldberg, O. and Bairlein, F., 2002. Ecological responses to recent climate change. *Nature*, 416(6879), p.389.

Welbourne, D.J., Claridge, A.W., Paull, D.J. and Lambert, A., 2016. How do passive infrared triggered camera traps operate and why does it matter? Breaking down common misconceptions. *Remote Sensing in Ecology and Conservation*, 2(2), pp.77-83.

Young, S., Rode-Margono, J. and Amin, R., 2018. Software to facilitate and streamline camera trap data management: A review. *Ecology and Evolution*, 8(19), pp.9947-9957.

Zheng, Z., Kirchner, P.B. and Bales, R.C., 2016. Topographic and vegetation effects on snow accumulation in the southern Sierra Nevada: a statistical summary from lidar data. *The Cryosphere*, 10, pp.257-269.

Acknowledgements

I would like to thank my supervisors Joris Cromsigt, Tim Hofmeester, and Sabine Pfeffer for their help and support throughout the duration of my study and for providing me with the opportunity to take on this study overseas in Sweden. Also, thanks to Sabine for enabling me to take part in fieldwork. I would also like to extend my thanks to SLU and Beyond Moose for allowing me to use their facilities and data for this study. Special thanks to thank my family for the support they have given me over the course of my thesis and my two years in Utrecht. Finally, thanks to my friends, especially Joel, Rosie and Nathan for their continued support over the course of both of my degrees.

Latest issue number in our series Examensarbete/Master's thesis

2019:5	Romsugning i Alterälven för att utreda sikens lekhabitatpreferenser Author: Simon Sundberg
2019:6	How Does White Rhino Respond to Fires During Dry Season? Author: Tim Herkenrath
2019:7	The role of predation in mass mortality of wood lemmings (Myopus schisticolor) Predationens roll i massdöd av skogslämlar (Myopus schisticolor) Author: Björn Wallgren
2019:8	An investigation into whether poaching creates an ecological trap for white rhinoceros in Hluhluwe-iMfolozi Park, South Africa Author: Alice Michel
2019:9	Evaluating effects of preventive actions to reduce wild boar damage in the agricultural landscape Author: Matilda Söderqvist
2019:10	Fire effects in a landscape of fear - food availability and predation risk as determining factors in microhabitat utilization prey in a heterogeneous landscape Author: David Kymmell
2019:11	Effect of variant ovarian fluid on sperm performance and egg fertilization rates of Arctic charr (Salvelinus alpinus L.) Author: Lwabanya Mabo
2019:12	Pedigree reconstruction reveals large scale movement patterns and population dynamics of wolverines (Gulo gulo) across Fennoscandia Author: Stephanie Higgins
2019:13	Community structure of polyporous fungi after wildfire in boreal forest Author: Isak Vahlström
2019:14	Population ecology of golden eagles (Aquila chrysaetos) using remote cameras Author: Andressa L. A. Dahlén
2019:15	Variations in nutritional content of key ungulate browse species in Sweden Author: Leonardo Capoani
2019:16	Assessing trust in the Swedish survey system for large carnivores among stakeholders Author: Philip Öhrman
2019:17	Beavers and environmental flow – the contribution of beaver dams to flood and drought prevention Author: Wali Uz Zaman

The entire list of published numbers can be found at www.slu.se/viltfiskmiljo