

Forest structure of Dalby Söderskog National Park - patterns of carbon storage in living and dead trees

Kumetra Achuthan

Master's thesis • 30 credits Swedish University of Agricultural Sciences Southern Swedish Forest Research Centre Alnarp 2021

Forest structure of Dalby Söderskog National Park - patterns of carbon storage in living and dead trees

Kumetra Achuthan

Supervisor:	Jörg Brunet, SLU, Southern Swedish Forest Research Centre
Examiner:	Per-Ola Hedwall, SLU, Southern Swedish Forest Research Centre

Credits:	30 ECTS
Level:	Advanced Level A2E
Course title:	Master's thesis in Forest Science
Course code:	EX0984
Programme/education:	SUFONAMA Master's Programme
Course coordinating dept:	Southern Swedish Forest Research Centre
Place of publication:	Alnarp
Year of publication:	2021

Keywords: Ash dieback disease; Deadwood; Dutch elm disease; Living tree volume; Coarse woody debris; Carbon stock and storage; Pest invasions; Temperate mixed broadleaved forest

Swedish University of Agricultural Sciences

Faculty of Forest Sciences

Southern Swedish Forest Research Centre

Publishing and archiving

Approved students' theses at SLU are published electronically. As a student, you have the copyright to your own work and need to approve the electronic publishing. If you check the box for **YES**, the full text (pdf file) and metadata will be visible and searchable online. If you check the box for **NO**, only the metadata and the abstract will be visible and searchable online. Nevertheless, when the document is uploaded it will still be archived as a digital file.

If you are more than one author you all need to agree on a decision. You can find more information about publishing and archiving here: <u>https://www.slu.se/en/subweb/library/publish-and-analyse/register-and-publish/agreement-for-publishing/</u>

 \boxtimes YES, I/we hereby give permission to publish the present thesis in accordance with the SLU agreement regarding the transfer of the right to publish a work.

 \Box NO, I/we do not give permission to publish the present work. The work will still be archived and its metadata and abstract will be visible and searchable.

Abstract

Carbon sequestration and storage is an ecosystem service supplied by forests, and is of increasing importance in the context of mitigation of global climate change. Forest pest invasions, driven primarily by globalizations, represent a risk to the efficiency of carbon sequestration as they may greatly reduce the amount of the living biomass in forests.

This study provides a combined assessment of living trees and coarse woody debris in the temperate broadleaf forest of Dalby Söderskog National Park in southern Sweden. The specific objectives were: (1) to investigate the storage of biomass and carbon pools, in both live and dead wood; (2) to find out the relationship between the distribution of live and dead woody biomass and carbon in relation to stem diameter; (3) to analyse the relative share of live and dead wood carbon pools along the stem diameter gradient and (4) to study the effect of fungal tree diseases on biomass and carbon dynamics.

Length and diameter of coarse woody debris including dead standing trees, cut stumps, and dead downed trees and branches were measured in 50 circular 100 m^2 (5.64 m radius) sample plots. Data for living trees for the same sample plots were taken from a previous inventory.

The results show that the studied broadleaf forest stored a total aboveground carbon stock of 176 t C/ha. Most carbon stock is stored by living trees (107.7 t C/ha), followed by downed deadwood (logs) (52.5 t C/ha) and standing dead wood (15.8 t/ha). The distribution of carbon storage in relation to stem diameter was species specific, and partly influenced by effects of Dutch elm disease and ash dieback.

This study shows that old-growth forests store large carbon stocks in living and dead wood. In the specific case of Dalby Söderskog, effects of tree diseases in mean time have increased the relative share of dead wood compared to live tree volumes, in particular for smaller diameter trees. The mixed tree species composition of the forest has, however, buffered the effects of Dutch elm disease and ash dieback, and living tree biomass and carbon still remains considerably larger than the amounts stored in dead wood.

Keywords: Ash dieback; Dead wood; Dutch elm disease; Living tree volume; Coarse woody debris; Carbon stock and storage; Pest invasions; Temperate mixed broadleaved forest.

Table of contents

Li	st of tab	lles	7
Li	st of fig	ures	8
Α	bbreviat	ions	9
1.	Intro	duction	10
	1.1.	Carbon stock in forest ecosystems	10
	1.2.	Carbon pool in forest ecosystem	10
	1.3.	Forest biomass and carbon stock	11
	1.4.	Factors influencing forest biomass	11
	1.5.	Study aims	13
2.	Mate	rials and Method	14
	2.1.	Study site description	14
	2.2.	Previous inventories	14
	2.3.	Method for live trees measurement	15
	2.4.	Field inventory 2021	16
	2.5.	Method for coarse woody debris measurement	16
3.	Resu	ılts	19
	3.1.	Volume by species for living trees	19
	3.2.	The volume of living trees by diameter classes	20
	3.2.1	. Fagus sylvatica	20
	3.2.2	. Fraxinus excelsior	20
	3.2.3	Quercus robur	20
	3.2.4	. Ulmus glabra	20
	3.2.5	. Other species	20
	3.2.6	. Tukey pairwise comparison	21
	3.3.	Biomass and carbon	22
	3.3.1	. Living biomass and its carbon	22
	3.3.2	. Carbon stock and DBH class relationship in Fagus sylvatica	23
	3.3.3	. Carbon stock and DBH class relationship in <i>Fraxinus excelsior</i>	24
	3.3.4	. Carbon stock and DBH class relationship in <i>Ulmus glabra</i>	25
	3.3.5	. Carbon stock and DBH class relationship in Quercus robur	25
	3.3.6	. Carbon stock and DBH class relationship in other species	26

	3.3.7.	Volume of deadwood	
	3.3.8.	Deadwood biomass and its ca	rbon 27
	3.3.9.	Carbon stock and DBH class	relationship in downed deadwood (logs). 28
	3.3.10	Carbon stock and DBH class	elationship in standing dead wood 28
	3.3.11	Overall carbon stock	
	3.3.12	Relative share of carbon stock	in living and dead woody biomass 30
	3.3.13	Paired analysis of live tree and	d CWD volumes 30
	3.3.14	Regression live tree and dead	wood
4.	Discu	sion	
4.	.1. I	sturbance by tree diseases	
4.	2. I	adwood volume dynamics	
4.	.3. I	adwood carbon stock	
4.	.4. I	e tree volume dynamics	
4.	5. I	e tree carbon stock	
4.	.6. I	e tree versus dead tree carbon	
4.	7.	udy limitations	
4.	.8.	nclusion	
5.	Refer	ces	
Acki	nowled	ements	
Арр	endix.		

List of tables

Table 1. Grouping information using the Tukey method and 95% confidence21
Table 2. Aboveground biomass (AGB), and carbon stock distribution for the tree species in Dalby Söderskog.
Table 3. Deadwood aboveground biomass (AGB) and carbon stock distribution forthe different types of deadwood category
Table A4. Descriptive Statistics. Volume of live trees by species.
Table A5. Summary of statistics for CWD49
Table A6. Wood volumes in 50 sample plots in Dalby Söderskog.

List of figures

Figure 1. Transect lines and sample plots in Dalby Söderskog (from Brunet et al., 2014). Blue color indicates the 50 plots included in this study15
Figure 2. Volume of living trees by species in Dalby Söderskog19
Figure 3. Volume of living trees divided into diameter classes by species21
Figure 4. Distribution of mean living tree volume (m ³ /ha ⁾ among plots by tree species
Figure 5. Carbon stock (t/ha) of different tree species in Dalby Söderskog23
Figure 6. Amount of carbon stock in different DBH classes in Fagus sylvatica24
Figure 7. Amount of carbon stock in different DBH classes in Fraxinus excelsior.
Figure 8. Amount of carbon stock in different DBH classes in Ulmus glabra25
Figure 9. Amount of carbon stock in different DBH classes in Quercus robur26
Figure 10. Amount of carbon stock and DBH class in Other species
Figure 11. Volume of different types of deadwood27
Figure 12. Carbon stock and DBH class relationship in logs
Figure 13. Carbon stock and DBH class relationship in standing deadwood29
Figure 14. Carbon stock among three fractions of aboveground tree biomass29
Figure 15. Percentage of the carbon stock in live and deadwood in different diameter classes
Figure 16. Relationship between CWD volume and live tree volume
Figure A17. Distribution of downed dead wood (logs) in different diameter classes.
Figure A18. Distribution of snags in different diameter classes47
Figure A19. Distribution of stumps in different diameter classes

Abbreviations

AGB	Above Ground Biomass
ADB	Ash Dieback
BEF	Biomass Expansion Factor
CWD	Coarse Woody Debris
DED	Dutch Elm Disease
IPCC	Intergovernmental Panel on Climate Change
SLU	Swedish University of Agricultural Sciences

1. Introduction

1.1. Carbon stock in forest ecosystems

The forest ecosystem is one of the essential parts of terrestrial ecosystems, and it occupies a key role in maintaining the global carbon cycle of terrestrial ecosystems (Liu et al., 1997; Kuuluvainen and Gauthier, 2018). Houghton, Hall, and Goetz (2009) estimated that forests store 70-90 percent of total global terrestrial biomass, even though this biomass is asymmetrically distributed across biogeographical regions: more than half of the total global terrestrial biomass is stored in tropical forests, while a third is stored in boreal forests and only 14% is stored in temperate forests (Pan et al., 2011). Conifers generally contain slightly more carbon (47-55%) than deciduous trees (46-50%), mainly because of a higher average lignin content, approximately 30%, compared with 20% for deciduous trees (Lamlom and Savidge, 2003)

Carbon sequestration and storage is one of the most vital ecosystem services supplied by forests, and it provides a lot of potential for mitigating climate change (Birdsey and Pan, 2015). Forests are major carbon sinks because of their high carbon content (Houghton, 2007). The world's forest ecosystems store more than double the carbon content in the atmosphere (Canadell and Raupach 2008; IPCC 2001). Ekoungoulou et al. (2015) state that the forest acts as a reservoir of carbon as it can store carbon from the atmosphere. This statement is supported by Kumar et al. (2019) who further add that forests keep large amounts of carbon in trees, understory vegetation, forest floor, and soil. Forest ecosystems store more than 80% of all terrestrial aboveground carbon (IPCC, 2001), which render forest ecosystem pivotal to maintaining the global carbon balance and mitigating climate change (IPCC, 2001).

1.2. Carbon pool in forest ecosystem

According to the Intergovernmental Panel for Climate Change (2006), the carbon dioxide fixed by plants through photosynthesis is transferred across different carbon pools. The IPCC identified five carbon pools in the terrestrial ecosystem that influence biomass storage: aboveground biomass, below-ground biomass, litter, woody debris, and soil organic matter. The estimates of total carbon stock from

each pool are summed to obtain the total forest carbon stock. Each year, as forests develop and their biomass grows, they take up carbon from the surrounding atmosphere and store up as plant tissue; hence, the forest has numerous and major functions regulating the atmospheric concentration of carbon dioxide (Ghosh, 2018).

1.3. Forest biomass and carbon stock

The mass of living biological organisms, including plants, animals, and microorganisms, is referred to as biomass in a given region or at a certain period (Hess and Tasa, 2011). Biomass in forest is counted for both above and below-ground biomass. Forest ecosystems contribute to climate change mitigation by storing and sequestering massive amounts of carbon in their biomass. The most accurate method to measure plant biomass is to harvest, dry, and weigh it (Clark and Kellner, 2012). However, this method is costly, time-consuming, and not always feasible, for instance, when monitoring forest's carbon stock. As a result, a number of non-destructive methods for effectively estimating biomass have been developed (Dittmann et al., 2017). Techniques to estimate terrestrial carbon storage and flux for large areas include a combination of simulation modelling, satellite imagery such as remote sensing and ground-measurement inventory data analysis (Goodale et al., 2002)

An allometric equation is a method of determining quantitative correlation between measurable parameters of stem diameter at breast height or height and relating this to characteristics that are harder to quantify, like the Above Ground Biomass (AGB). These relationships are usually species and geographically specific (Muukkonen, 2007). Muukkonen, (2007) further found that the allometric methodology to estimate biomass is relatively accurate but requires time, thus making it best suited for implementing across small areas.

1.4. Factors influencing forest biomass

Many drivers contribute to the changes in biomass and carbon storage. Climate (Luyssaert et al., 2008), soil (Angst et al., 2018) and natural disturbance (Bond-Lamberty et al., 2007), as well as forest stand properties including tree species composition (Nord-Larsen et al., 2019), age and silviculture practices (Liski et al., 2006), substantially affect carbon sequestration in forests. Stand age is one of the key factors determining above ground biomass in forests (Bradford et al., 2008; Chatterjee, Vance and Tinker, 2009). Furthermore, Luyssaert et al., (2008) stated that the total ecosystem carbon pool is corresponding to stand age and strongly aligned with site productivity. Besides that, several studies (Vucetich et al., 2000; Seedre et al., 2015), documented that forest biomass storage, particularly AGB storage, has been found to decline with increasing altitude and latitude.

Environmental factors, particularly temperature and precipitation and land use history, have been linked to biomass storage and accumulation in primary forests, with forests with less or no land-use history retaining the most carbon (Keith, Mackey, and Lindenmayer, 2009). These authors further found that the highest biomass densities have been reported in temperate forests with less land use history, where the mean annual temperature is lower than 10°C and the mean annual precipitation is between 1 000 and 2 500 mm. On the other hand, a high amount of live tree biomass can be changed into deadwood following windthrow or fire events, while a high proportion of live biomass is removed from the site following clear-cut (Seedre et al., 2011). Besides that, invasive pests (Fei et al., 2019), due to climate change are also regarded as a factor resulting in biomass losses.

Primary forest carbon storage is receiving more attention (Badalamenti et al., 2019), and research on primary forests in various ecoregions has shown that they can store large amounts of carbon and that many are still accumulating carbon (Luyssaert et al., 2008, Keith, Mackey, and Lindenmayer, 2009; Pan et al., 2011; Pugh et al., 2019). However, there is a lack of research related to natural forest disturbance, although such disturbances are part of natural dynamics in a forest ecosystem.

Disturbances work to move carbon from living trees to dead wood, soil, and the atmosphere in the short term, but they also change long-term patterns of carbon storage as stand structure and species composition shift during recovery (Houghton, Hall and Goetz, 2009). Disturbance can be divided into two factors, biotic (anthropogenic activity, insect damage, disease-causing fungi) and abiotic (storm, wind, fire, and snow).

Fungal pathogens can do significant damage to trees and are a common problem in forests across the world (Potter and Urquhart, 2017). Tree mortality is a critical factor in forest structural dynamics and carbon cycling in terrestrial ecosystems. Disturbance that results in partial stand replacement has been proven to influence biomass accumulation (Keith, Mackey, and Lindenmayer, 2009).

One of the major environmental concerns of the modern era is the increase in carbon dioxide in the atmosphere and its potential effect on climate warming, but also indirect effects of warming on pathogen and disease spreading. Almost all effects of insect outbreaks are strongly dependent on climate (Pinkard et al., 2011). In southern Sweden, the spread of Dutch Elm Disease (DED) started in the 1980s and has since destroyed many elm populations. Since the early 2000s and on a continental scale, the ascomycete *Hymenoscyphus fraxineus* which was first reported in Poland in Europe (McKinney et al., 2014) is causing harm to the widely spread native European ash species, *Fraxinus excelsior*. As a result of a spread across Europe from east to west over the last two decades, populations of ash trees (*Fraxinus excelsior*) were decimated (Pautasso et al., 2013). Ash dieback (ADB) is one of several tree diseases and insect pests that have recently arrived and swept across European forests (Boyd et al., 2013). Besides that, Waller (2013), stated that the global climate change may worsen the long-term effects of Dutch elm disease and ash dieback.

In southern Sweden, a quarter of *Fraxinus excelsior* trees were reported to be dead or critically harmed (Stenlid et al., 2011). Both Ash (*Fraxinus excelsior*) and Elm (all three species; *Ulmus glabra*, *U. minor*, and *U. laevis*) populations have declined severely in Sweden because of fungal infections and have been categorized as vulnerable on the Swedish red list since 2010 (Pihlgren et al., 2010).

In this thesis, we set out to bridge the gap in knowledge about the effects of disturbance severity on the long-term carbon dynamics in broadleaf tree species. For this purpose, we explore extensive tree volume and coarse woody debris data from an unmanaged deciduous forest located in southern Sweden, Dalby Söderskog. This forest is chosen as it is severely affected by Ash dieback and Dutch elm disease. However, since not all trees are killed, structural growth following perturbations occurs in various ways with different live and dead tree biomass components.

Hence, the aim is to determine how disturbance of DED and ADB affected carbon stocks on live tree boles and coarse woody debris (CWD). Although forest structure in the study area has repeatedly been documented in the past century (Brunet et al., 2014), this is the first study on woody biomass and carbon content to date.

1.5. Study aims

The objectives of the study were:

- 1) To investigate the storage of biomass and carbon pools, in both live and dead wood in Dalby Söderskog.
- 2) To find out the relationship between the distribution of live and dead woody biomass and carbon in relation to stem diameter.
- 3) To analyse the relative share of live and dead wood carbon pools along the stem diameter gradient.
- 4) To study the effect of fungal tree diseases on biomass and carbon dynamics.

2. Materials and Method

2.1. Study site description

Dalby Söderskog has size of 37 ha and is located in the County of Skåne, which is in southern Sweden, 10 km east of Lund (55°41′N, 13°20′E, 65m a.s.l.). It is a mixed broadleaved deciduous forest that is dominated by Pedunculate oak (*Quercus robur* L.), European ash (*Fraxinus excelsior L.*), European beech (*Fagus sylvatica L.*), and wych Elm (*Ulmus glabra* Huds). The mineral soil, which is derived from Baltic moraine (from the Weichsel glacial period), is a calcareous, nutrient-rich clay with a high calcium content. The soil is a eutric cambisol, and humus is of the mull type. The climate is temperate suboceanic, with an annual mean temperature of 7.5 degrees Celsius and annual mean precipitation of approximately 650 millimeters (Oheimb and Brunet, 2007). The latest significant cuttings were done in 1914 – 1916, whereby 1,600 m³ wood were taken out of a total of 8,000 m³. The forest status changed into a protected forest in the year of 1918. Since then, this forest has no major management except removal of trees that can be dangerous for forest visitors. Since 1988 dead elms were cut down due to safety reasons around hiking paths (Brunet et al. 2014).

2.2. Previous inventories

Several inventories were carried out previously in Dalby Söderskog, mainly about stand structure and vegetation (Lindquist 1938; Malmer et al., 1978; Brunet et al., 2014). Lindquist (1938) studied forest structure and vegetation by establishing transect lines and sample plots. This method used a straight path through the forest as a baseline and distributed 74 sample plots along the perpendicular lines. The line-to-line distance is 50 m, while the plot-to-plot distance along lines in most cases is 100 meters (Lindquist 1938). From a first survey in 1909, species wise total stems number with more than 20 cm diameter breast height (DBH) are known, while species wise stem numbers divided in dbh-classes are available for surveys in 1916, 1935, 1970 and 2012 (Brunet et al. 2014).

In the early spring of 2010, 74 sample plots were reconstructed for a survey of herb layer vegetation by using an aerial photograph with Lindquist's map as a digital overlay to extract GPS coordinates (Fig. 1, Brunet et al. 2014). In addition, the plot centre of each plot was marked with a plastic stick, and a short iron rod for future relocation of plots in case of loss of plastic stick. In February 2012, Bukina (2012),

carried out a tree inventory in circular sample plots. Trees with dbh of 10 cm or more were identified and measured in plots of 100 m^2 (5.7 meter radius); meanwhile trees with dbh of 20 cm or more were inventoried in 314 m² (10 meter radius plot). Trees below 10 cm in dbh are not included in the inventory. All the plots were taken from the center points of the 74 plots which were established previously.

In addition, vitality of trees was assessed due to the Dutch Elm Disease. All stems were classified as living or dead.



Figure 1. Transect lines and sample plots in Dalby Söderskog (from Brunet et al., 2014). Blue color indicates the 50 plots included in this study.

2.3. Method for live trees measurement

The most recent inventory was conducted by Ruks (2020), who followed the same method as Bukina (2012) and registered all the trees with dbh class 10 - 19 cm in a plot size of 100 m^2 (5.64 m radius), and dbh of 20 cm or more in a plot size of 314 m² (10 m radius), sharing the same plot centre. All the trees were identified to the species, and stem diameter was measured at 13 m height. Diameter (dbh) was registered for all individuals by using the cross-calliper method. For trees that were larger than 50 cm in dbh, measuring tape was used.

For the calculation of live tree carbon contents in this study, the available data set measured by Ruks (2020) was used. The carbon content of live trees, bole and bark, was assumed to be 50% of bole and bark mass for all species. Tree height and DBH are necessary to estimate volume, used to calculate biomass accumulation and carbon stock. The DBH of trees were directly obtained from Ruks (2020) data set, but live tree height data to calculate volume were lacking. Instead, data from Lindgren's (1971) volume table chart were used to assume the volume of whole living tree (Table A7). Lindgren's data summarize the results of the forest inventory of Dalby Söderskog in 1970 and provide average volumes in 5 cm diameter classes and specified for the four main tree species and a fifth group containing other species.

2.4. Field inventory 2021

Data were collected in January and February 2021. The inventory was conducted on the existing permanent plots used by Ruks (2020). The focus of this inventory was on the measurement of the above ground deadwood accumulation in the forest, partly due to the past DED and the ADB. The sample plots consisted of 50 of the 74 permanent plots (Fig. 1). Due to time constraints, only 50 randomly distributed plots throughout the forest were chosen as sample plots. To locate the plots, handheld GPS and a map were used.

2.5. Method for coarse woody debris measurement

Three principal forms of aboveground coarse woody debris were sampled in each plot: logs, stumps and snags. Logs were defined as downed tree boles at least 10 cm in diameter at mid-length. Notably, only sections of logs that fall inside the plots were measured. Stumps were classified as standing cut tree boles with at least 10 cm in stem diameter. Meanwhile, snags were defined as standing, uncut dead trees at least 10 cm in dbh. Finally, each log, stump and snag were assigned having tree bark or not.

Log lengths were measured with measuring tape. Log mid-diameters were measured using a calliper, and a diameter tape was used for diameters more than 50 cm. Stump diameter was measured just below the cut by calliper, meanwhile, stump height by using measuring tape. Snag diameter was measured at 1.3 m height by using a calliper. For snags the height was estimated visually "by eye" calibrating against a 2 m folding rule (Harmon and Sexton, 1996).

The volume of cylinder method calculated log and snag stump volume as below

V=h $\times \pi \times d2/4$ Equation 1

Where V is volume (m^3) , h is height (m), and d is the diameter (cm)

Snag volume is estimated by the formula for an ellipsoid cone method (Brunet and Isacsson, 2009). The formula is as follows:

 $V=\pi \times d2 \times h/6$ Equation 2

Where V is volume, d is diameter, and h is snag height in meters.

CWD biomass was calculated by multiplying Biomass Expansion Factor (BEF) and total volume of CWD

W=BEF×V..... Equation 3

where W denotes the aboveground tree biomass (dry weight, Mg), BEF denotes the biomass expansion factor (Mg m^{-3}), and V denotes the stem volume (m^{3}).

The BEFs were used to calculate the total aboveground biomass of trees based on the stem volume estimates

In this study, the constant BEF of 0.64 for broadleaved stands was used to calculate the total aboveground biomass of trees based on the stem volume estimates. The constant BEF value (0.64) was obtained from the Swedish National Inventory Report (Feldhusen et al., 2004). This is consistent with the most recent greenhouse gas reporting from Sweden. Still, they consider aboveground biomass as a product of volume and 64 percent for all broadleaved tree species in their calculations (Feldhusen et al., 2004).

Lastly, in accordance with IPCC guidelines, biomass was converted to carbon by assuming a carbon content of 47% (IPCC, 2006):

C=Wx0.47.....Equation 4

where W is the aboveground tree biomass (dry weight, Mg), 0.47 carbon content.

Meanwhile for live tree:

C=Wx0.5.....Equation 5

where W is the aboveground tree biomass (dry weight, Mg), 0.50 carbon content.

Assume 50% of biomass is made up of carbon (Houghton 2007; Thomas and Martin, 2012).

Total biomass carbon is the sum of all individual carbon pools. We will refer to the sum of live and dead biomass carbon as total biomass carbon. We used paired sample t-test analysis and analyse of posthoc Tukey HSD test for pairwise comparisons to determine differences between volume of living and deadwood trees after disturbances. All statistical analyses were done in Microsoft Excel 2016 and Minitab 19 (Minitab Inc.) with statistical significance established at $\alpha = 5\%$ level.

3. Results

3.1. Volume by species for living trees

Dalby Söderskog is mainly dominated by broadleaf species of European Ash (*Fraxinus excelsior*), Beech (*Fagus sylvatica*), Pedunculate oak (*Quercus robur*), and Wych Elm (*Ulmus glabra*). In contrast, other species are less abundant (consist of black alder (*Alnus glutinosa*), Norway maple (*Acer platanoides*) and crab apple (*Malus sylvestris*). The results obtained for the old-growth broadleaf stand show that the mean standing volume for the live trees is $336.7\pm 29.6 \text{ m}^3$ /ha (Mean \pm S.E., Tables A4, A6). Figure 2 shows that *Fraxinus* and *Fagus* were the most abundant tree species accounting for 35.4% and 32.6% of the total wood volume found in the plots. *Quercus* has the third highest volume (24.2%), followed by *Ulmus*, which has the second lowest volume (4.8%). Other species were least plentiful, constituting only 3.1% of the total volume.



Figure 2. Volume of living trees by species in Dalby Söderskog.

3.2. The volume of living trees by diameter classes

3.2.1. Fagus sylvatica

For beech, the highest volume was recorded in the diameter class 30 to 39 cm and the lowest in the dbh class of 110 to 119 cm (Fig. 3). Compared to the other tree species, the distribution of volume over diameter classes was more even.

3.2.2. Fraxinus excelsior

For European ash, the diameter class 40 cm to 49 cm held the largest volume (Fig. 3). Besides, the volume of ash occurring in diameter classes greater than 50 cm is relatively high except for diameter classes ranging from 90 to 99 cm (no trees available). Meanwhile, the lowest volume can be seen in the diameter class of 10 to 19 cm, which indicates a low distribution of small trees.

3.2.3. Quercus robur

Pendunculate oak ranks third in terms of living tree volume. The highest value is reported for the diameter class of 80 to 89 cm, meanwhile, only a small volume is recorded for the diameter class of 10 to 19 cm, and no trees with dbh 20 to 39 cm were recorded (Fig. 3).

3.2.4. Ulmus glabra

For wych elm, the volume distribution is very different from other species and dominated by small trees. The volume of trees is highest on the small-sized tree of the diameter class interval 10 to 19 cm, and the smallest volume can be seen in the diameter class of 30 to 39 cm, while no larger trees were found (Fig. 3).

3.2.5. Other species

Lastly, for the other tree species, small volumes range from diameter classes from 10 to 19 cm to the diameter class of 70 to 79 cm (Fig. 3).



Figure 3. Volume of living trees divided into diameter classes by species.

3.2.6. Tukey pairwise comparison

According to Table 1, relationships between species and volume revealed that species within groups do not differ significantly in their means, but among groups they do. *Fraxinus, Fagus* and *Quercus* categorized under one group, meanwhile *Ulmus* and Others in another group which is differ from Group A species. The group A has significantly higher means than group B

Factor (volume)	Ν	Mean	Grouping	
Fraxinus	50	119,0	А	
Fagus	50	109,9	А	
Quercus	50	81,4	А	
Ulmus	50	16,0	В	
Other species	50	10,3	В	

Table 1. Grouping information using the Tukey method and 95% confidence.

Accordingly, Figure 4 shows the means and confidence intervals for the different species with confidence intervals overlapping for *Fraxinus, Fagus* and *Quercus* on the one hand, and *Ulmus* and Others on the other hand.



Figure 4. Distribution of mean living tree volume (m^3/ha) among plots by tree species.

3.3. Biomass and carbon

3.3.1. Living biomass and its carbon

Dalby Söderskog forest stocked, a total of 215.5 t/ha living aboveground biomass (Table 2). *Fagus sylvatica* and *Fraxinus excelsior* accounted for about 70 % of total biomass stored in the old broadleaf forest (Table 2, Fig. 5). The quantity of carbon stored by the different tree species was hugely different, with *Ulmus glabra* and the group of other species carbon stocking least carbon. The mean carbon stored by the broadleaf forest's living biomass was 107.7 t C/ha.

T	AGB)	Carbon stock	Percentage
Tree species	(t/ha)	(t C/ha)	%
Fagus sylvatica	70.3	35.2	32.6
Fraxinus excelsior	76.2	38.1	35.4
Ulmus glabra	10.3	5.1	4.8
Quercus robur	52.1	26.0	24.2
Others	6.6	3.3	3.1
Total	215.5	107.7	100.0

Table 2. Aboveground biomass (AGB), and carbon stock distribution for the tree species in Dalby Söderskog.



Figure 5. Carbon stock (t/ha) of different tree species in Dalby Söderskog.

3.3.2. Carbon stock and DBH class relationship in Fagus sylvatica

Results obtained show that *Fagus sylvatica* had the second-highest carbon storage (32.6 %) among the tree species. Carbon stock was relatively evenly distributed across DBH-classes (Fig. 6). The diameter class of 30 to 39 cm had the greatest carbon stock, with 5.1 t C/ha, and the largest diameter range from 110 to 119 had the lowest carbon stock, 1.6 t C/ha.



Figure 6. Amount of carbon stock in different DBH classes in Fagus sylvatica.

3.3.3. Carbon stock and DBH class relationship in *Fraxinus* excelsior

Fraxinus excelsior species has recorded the highest carbon stock of 35.4% total carbon content among the tree species for the diameter class of 40 to 49 cm with 8.5 t C/ha. Meanwhile, there were no trees recorded in the 90 to 99 cm class, and the small diameter classes of 10 to 19 cm (0.56 t C/ha) and 20 to 29 cm (0.54 t C/ha) only contained low amounts (Fig. 7).



Figure 7. Amount of carbon stock in different DBH classes in Fraxinus excelsior.

3.3.4. Carbon stock and DBH class relationship in Ulmus glabra

Ulmus glabra has only three diameter classes of living trees (Fig. 8). There are no trees present for the diameter classes of more than 40 cm, and this species is classified as having young stand distribution across all the area. The amount of stored carbon is gradually decreasing in between diameter classes. The maximum carbon content of 3.6 t C/ha was observed in the 10 to 19 cm diameter class, followed by (1.12 t C/ha) in the 20 to 29 cm, and finally, the lowest value (0.43 t C/ha) in the 30 to 39 cm class.



Figure 8. Amount of carbon stock in different DBH classes in Ulmus glabra.

3.3.5. Carbon stock and DBH class relationship in Quercus robur

Quercus robur accumulated the third-highest amount of carbon storage among the live tree species. The most significant carbon sequestration (2.84 t C/ha) is seen in the 80 to 89 cm diameter class (Fig. 9). Besides, approximately 52% of the carbon sequestration is represented in the diameter class range between 70 to 79 cm and 80 to 89 cm. Meanwhile, a very low carbon stock is seen in the diameter classes below 50 cm.



Figure 9. Amount of carbon stock in different DBH classes in Quercus robur.

3.3.6. Carbon stock and DBH class relationship in other species

The group of other species has been shown to have the lowest carbon storage of all the species studied. The highest carbon content (1.40 t C/ha) was stored between 30 to 39 cm diameter. Meanwhile, no trees and thus no carbon stock was found in the diameter range of 50 to 69 cm (Fig. 10).



Figure 10. Amount of carbon stock and DBH class in Other species

3.3.7. Volume of deadwood

According to the findings, the mean deadwood volume in Dalby Söderskog is 227.8 \pm SD 196.1 m³/ha (Tables A5, A6). The volume of deadwood is divided into two types: logs had the highest amount (174.4 \pm SD 156.4 m³/ha), and the snags had (53.5 \pm SD 117.9 m³/ha, Fig. 11).



Figure 11. Volume of different types of deadwood

3.3.8. Deadwood biomass and its carbon

The mean deadwood aboveground biomass in Dalby Söderskog was estimated to be $145.3\pm$ SD 125.5 t/ha (Table 3). The details show large variation around the means, which is expected given the relatively small size of our sample plots.

A higher biomass is stored in downed logs (111.6 t/ha), than in standing dead wood (33.7 t/ha). Downed deadwood comprised the greatest proportion of total carbon stock, followed by standing deadwood 77%, and 23%, respectively (Table 3). The total carbon stored in deadwood was $68.3\pm$ SD 59.0 t C/ha on average.

	71 5 0 7		
Type of deadwood	AGB)	Carbon stock	Percentage
	(t/ha)	(t C/ha)	%
Downed logs	111.6	52.5	77
Standing deadwood	33.7	15.8	23
Total	145.3±SE17.7	68.3±SE8.3	100

 Table 3. Deadwood aboveground biomass (AGB) and carbon stock distribution for the different types of deadwood category.

3.3.9. Carbon stock and DBH class relationship in downed deadwood (logs)

The overall amounts of carbon stored in logs were higher in smaller than larger diameter classes (Fig. 12), due to a much higher number of logs and snags in these classes (Appendix, Fig. A17, A18).



Figure 12. Carbon stock and DBH class relationship in logs.

3.3.10. Carbon stock and DBH class relationship in standing dead wood

Results show that the carbon content among standing deadwood was largely stored in smaller diameter classes (< 60 cm) (Fig. 13).



Figure 13. Carbon stock and DBH class relationship in standing deadwood.

3.3.11. Overall carbon stock

The results obtained indicate that in Dalby Söderskog, the total aboveground carbon stock is 176 t C/ha. The overall result for carbon storage among various aboveground biomass shows that carbon storage decreases from live trees over logs to standing dead wood (Fig. 14). The highest biomass is stored in living trees 107.7 t C/ha, followed by downed deadwood (logs) 52.5 t C/ha, and then standing dead wood with 15.8 t C/ha.



Figure 14. Carbon stock among three fractions of aboveground tree biomass.

3.3.12. Relative share of carbon stock in living and dead woody biomass

According to Figure 15, the relative share of carbon stored in live trees increases with stem diameter. Up to a stem diameter of 39 cm, it is lower than 50%, which means that more carbon is stored in dead than living trees. For the large diameter classes, living trees generally store more carbon than dead wood, ranging from a share of 59 to 100% of total carbon stored in wood.



Figure 15. Percentage of the carbon stock in live and deadwood in different diameter classes.

3.3.13. Paired analysis of live tree and CWD volumes

A paired t-test analysis indicated that the mean living tree volume $(336.8 \pm SE 29.6)$ was significantly higher than the CWD volume $(227.9 \pm SE 27.7)$ with (t=-2.32, P=0.025). The variation between plots is high, but the p-value is still statistically significant.

3.3.14. Regression live tree and deadwood

A linear regression analysis showed that there is a weak but significant negative relationship between live tree volume and CWD (p=0.015). The model is implying that the amount of CWD increases with decreasing live tree volume (Fig. 16).



Figure 16. Relationship between CWD volume and live tree volume (V=335, P<0.015<, R^2 11.7%. The regression equation is: CWD vol $m^3/ha = 335.5 - 0.3196$ (y), y = live tree volume.

4. Discussion

4.1. Disturbance by tree diseases

Insect and disease invasions in southern Sweden have resulted in considerable increases in regional tree death rates and the conversion of live carbon to dead matter. The findings of this study show that CWD has a substantial carbon storage capacity; however, it is lower than live tree biomass. The Dutch elm disease, which wiped out millions of elms in the 20th century (Martin et al., 2010), has been affecting Dalby Söderskog since 1990. It appears to be a significant influence in modifying forest succession processes. A comparable investigation carried out by Brunet et al., (2014), reported pests and diseases can shape ecosystems and influence succession. Meanwhile, Ash dieback was also affecting the trees in Dalby Söderskog. In Sweden, the first evidence of ash dieback was documented in 2002 (Stener, 2013).

The regression model between living and deadwood volume was found to be inversely related. The model predicts that CWD increases with decreasing live tree volume. When living trees in a stand with low live tree stocking have been killed by disturbance, the highest deadwood stocking will be obtained. This study's findings are comparable to those of other studies which indicated that forest disturbance influences pattern of deadwood accumulation (Woodall and Westfall, 2009) as the amount of deadwood in the forest is determined by both accretion variables (disturbance, self-thinning, and senescence) and depletion variables (harvesting and decay). Forest debris is formed because of forest biomass mortality (Sollins, 1982; Spies et al., 1988; Sturtevant et al., 1997; Bond-Lamberty and Gower, 2008), hence correlation between live and dead forest biomass should be apparent at some level. Such mortality might be expected to impact carbon dynamics by reducing forests' carbon sequestration capacity and by converting live materials to dead carbon source (Fei et al., 2019).

4.2. Deadwood volume dynamics

The total mean volume of deadwood found in the study area is 227.8 m³/ha. Deadwood was present in all the surveyed plots as lying logs and snags (standing and stump). However, lying logs accounted for a greater proportion of the total dead volume than snags. Logs are the most important deadwood components with 77 % of total volume.

This finding is supported by prior research by Krankina et al., (2002). Meanwhile, several studies have found similar results, reporting 2–3 times more lying (approximately 60%–70%) than standing deadwood regardless of the dominant species (Siitonen, 2001; Nilsson et al., 2002; Karjalainen and Kuuluvainen, 2002). The most recent study conducted in Romanian deciduous forests (Öder et al., 2021) reported that lying deadwood objects accounting more than 70% of the total deadwood compares to standing deadwood (30%).

The fact that there were fewer snags in most plots surveyed could be due to the transitory nature of this component. This finding means that after increasing tree mortality, the number of snags can increase briefly, but they eventually fall over and become felled logs (Herrero et al., 2014). Therefore, the distribution of downed deadwood on the forest floor is higher. Additionally, Passovoy and Fule, (2006) found that standing dead trees with a smaller diameter collapse over more quickly, resulting in top breaking, so this could be why the volume of downed deadwood is higher.

The present study highlights that in the broadleaf forest of Dalby Söderskog, deadwood pieces are mainly concentrated in the smaller diameter classes (diameter class between 10 cm and 30 cm). Data from Dalby Söderskog significantly correlate to the finding of Keßler et al. (2012) that the *Hymenoscyphus fraxineus* pathogen attacks all size classes, yet, symptoms and eventual mortality progress more rapidly in smaller individuals classes. When a tree is dead and standing, it requires a certain amount of structural support to keep its vertical structure; failing in doing it will cause more destruction (Woodall and Westfall, 2009). Alternatively, CWD is highest in young stands, decreases in developing stands, and rises slightly in mature stands as tree mortality increases after disturbance (Harmon et al., 1986; Spies et al., 1988).

The CWD volume found in Dalby Söderskog is comparable with the findings of Bujoczek et al. (2018) who found a mean volume of deadwood in montane beechfir forests of 223.9 m³/ha. Meanwhile, according to Christensen et al. (2005), the mean deadwood amount in unmanaged beech forests was 130 m³/ha, but the variation among reserves was high, ranging from almost nil to 550 m³/ha. Also, in the Pacific Northwest, Spies et al. (1988) identified tremendous values for Douglas fir of 423 m³/ha mean CWD volume for young stands (age 65 years) and 250 m³/ha for mature forests (age 121 years). Also, according to Burrascano et al. (2013), significant levels of deadwood (over 200 m³/ha) are typical in old growth stands in central European forest. These findings can be correlated with our study whereby the old stand has almost similar values.

4.3. Deadwood carbon stock

The large disturbance initiated by Dutch elm disease three decades ago greatly influenced the amount and composition of carbon stock. Notably, CWD stored 68 t C/ha, 39% of the total carbon stock of woody material in the entire surveyed area. Modelled deadwood carbon stocks for broadleaf species are higher than most results from other literature. This could be due to the decay rates reduction factors were not used in this investigation.

According to Christensen et al., (2005), the carbon stock of deadwood in beech forests in central and eastern Europe ranges from 1.4 to 82.4 t C/ha⁻ Similar species composition in the semi-natural forest reserve Suserup Skov in Denmark accounted for the carbon stocks in deadwood about 35 t C/ha from the overall ecosystem carbon stock (Nord-Larsen et al., 2019). Dalby Söderskog stored two times higher carbon stock compared to Suserup Skov. This could be because of disease related supply of large diameter logs from ash and elm, not only beech and oak on the forest floor, versus Suserup Skov, which is only dominated by beech and oak deadwood.

Apart from it, a study by Wirth and Lichtstein (2009), show at the age of 200 years, deadwood stocks in temperate broadleaved forest (among others *Nothofagus* and *Fagus*) lies between 10 and 90 t C/ha. This finding falls within the range of our results. Hadden and Grelle (2017), demonstrated when relatively slight disruptions resulted in more deadwood in the primary boreal forest. Therefore, net emissions from deadwood rose, changing overall ecosystem respiration and shifting the forest from a carbon sink to a source of carbon dioxide (Hadden and Grelle, 2017).

4.4. Live tree volume dynamics

The mean volume of live trees is 338 m³/ha, 100 m³ more than the deadwood volume. The live tree volume remains high after forest disturbance, except there are changes in species composition per volume. Remarkably, the present composition and structure of Dalby Söderskog vary drastically from their historical range of variability. According to Malmer et al. (1978), the order of species abundance in Dalby Söderskog in the 1970's was as follow *Ulmus* > *Fraxinus* > *Fagus* > *Quercus*. However, the recent finding indicates *Fraxinus* > *Fagus* > *Quercus* > *Ulmus*, and this corresponds to Waring and Running (2007) that disturbances alter forest species composition.

Elm was once the most dominant tree in the forest (\geq dbh 20 cm), but Dutch elm disease had devastated the elm population by 2011 (Brunet et al., 2014). Then it

appears to play a significant role in modifying forest succession trends. DED occurrence in Dalby Söderskog was followed by increasing regeneration of ash. According to Oheimb and Brunet (2007), canopy gaps create ideal conditions for ash recruitment. Ash was known to be relatively robust; yet, it has recently become a primary issue in most of Europe due to a pathogenic fungus that causes ash dieback (McKinney et al., 2011).

4.5. Live tree carbon stock

The Dalby Söderskog mixed broadleaf species contribute to carbon storage by storing 107.7 t C/ha. The capacity of forest ecosystems to store carbon in biomass is affected by species composition, age, and population density (Mendoza-Ponce and Galicia 2010).

Ash is the most important species with total carbon stock contribution (36.7 %), then beech (33.2 %), followed by oak (22.7%), Elm (4.5%), and the lowest contribution of other species (2.9%). Beech is the most abundant shade-tolerant tree species in European temperate forests and may be additionally favoured by adverse effects of ash dieback in eutrophic forests, where Ash usually shows a faster initial growth during regeneration (Emborg, Christensen and Heilmann-Clausen, 2000). European Ash (*Fraxinus excelsior*) is a fast-growing tree species (Fraxigen, 2005), common in different forest types and can regenerate naturally through seed fall.

Meanwhile, pedunculate oak was ranked third. In Dalby Söderskog, it is mainly represented by old individuals (Brunet et al., 2014). Oak distribution in small diameter classes is very poor due to canopies closed due to other dominating species, making no gaps for the light-demanding oak to grow in. Consequently, regeneration of oak drastically declined because of mortality and low regeneration due to canopy closure.

Elm species noticed low carbon storage due to the forest is severely impacted by Dutch elm disease, which has devastated the whole forest standings since the late 1980s. Furthermore, according to Brunet et al. (2014), Elm was limited to the shrub layer, surviving as shrubs or small trees. Other species such as maple and cherry all exhibited low abundance and carbon stocks. It may be due to the low frequency of the tree species in the adjacent forests and a quick establishment by other more persistent and competitive tree species.

If 50% of biomass is carbon (Houghton, 2007; Thomas and Martin, 2012), mean biomass carbon density in the Swedish primary forests ranged between 9 t C/ha and 91 t C/ha. Our findings show that the Dalby Söderskog stored more carbon than the average Swedish forest. However, compared to 28 t C/ha and 229 t C/ha in other research on carbon storage in Nordic and Baltic primary forests (Vucetich et al., 2000; Kenina et al., 2019; Nord-Larsen et al., 2019), 107.7 t C/ha is within that range.

Our research demonstrates the substantial carbon stores associated with largediameter trees in the region. Still, there is the potential for significant losses in aboveground carbon with giant trees prone to biotic and abiotic causes if any additional disturbance occurs soon.

4.6. Live tree versus dead tree carbon

Even though biotic and abiotic factors affect the forest, the results show that living tree carbon is higher than deadwood carbon in this circumstance. The Dalby Söderskog obtained above ground carbon (without soils) of 175.7 t C/ha with 107.7 t C/ha stored in the living biomass pool and 68.3 t C/ha in the deadwood carbon pool. This forest has a relatively high total carbon stock.

According to findings, highest carbon stock of live trees stored in diameter class is more than 50 cm, 71.2 t C/ha, and for coarse wood debris in diameter class of less than 60 cm, 50.7 t C/ha. The higher share of deadwood in lower dbh classes is likely related to the high elm mortality that previously dominated these smaller dbh classes, as shown in the carbon stocked along diameter gradient (Fig. 15). This simplifies, most small diameter classes less than 60 cm are abundant in the form of CWD, indicating that the stand still maintains an abundance of large trees that are consistently productive and have numerous second generations upraised.

The succession pattern changes, the dispersal is allowed to run its course, and mortality will occur at different rates among different size classes, influencing regeneration differently among species. As a result, it will impact the accumulation of living biomass and cause things to alter in unexpected ways.

The absence of management promoted the existence of developed forest successional growth phases, resulting in a higher level of living biomass, tree density and deadwood volumes. In this scenario, the current disruption for Ash is still early, and the pattern of carbon stored in CWD rather than live trees has considerably less influence. For the time being, Ash continues to outnumber other tree species in terms of distribution, with the distribution of species predicted to shift again in the following decades.

4.7. Study limitations

The main limitation in our study is the lack of decay classes for each deadwood type. The results on biomass and carbon obtained in this study, especially for deadwood, vary as published in other scientific papers; this is because most studies have applied deadwood decomposing reduction classes (1 to 5) when calculating deadwood biomass. The decay class represents the stage of decomposition of dead

wood and is usually determined by visual criteria. The classification of CWD by decay class is crucial since decay affects wood density and subsequent carbon content. Grove et al. (2009), show that log volumes measured in field inventories can be transformed into CWD mass and carbon content based on the decay class and estimate corresponding wood densities. Failing to apply decay classes in calculation could produce errors for measured carbon pools.

In this study, with the absence of a survey about decay classes for each piece of deadwood, we cannot apply the decomposition reduction factor when computing carbon, it's possible that the carbon content of deadwood has been overestimated. Based on field observation on 692 pieces of deadwood, 72% was with absence of barks, and only 18% had intact bark. Based on surveyed deadwood, it can be said deadwood mainly in early decay stages, were still fresh, and the structure is hard. Hence, considering carbon contents in different decay stages would improve the accurate assessment of carbon stocks in deadwood.

Besides that, the resulting biomass and carbon estimate highly depend on the samples and the conceptual approach utilized (i.e., whether biomass functions or biomass expansion factors are used), Neumann et al. (2016). In this study, a constant biomass expansion factor of 0.64 was used, which might probably influence the results obtained. Pietsch and Hasenauer (2002), underlined that constant biomass expansion factors are widely known for overestimating young trees and underestimating biomass for older trees.

Another limitation is that the secondary data from previous research have no data on essential parameters, the height of the live tree. Estimating volume value from the volume production table was used to resolve these issues, which raises some doubts about the volume and biomass of the live tree.

4.8. Conclusion

As a concluding remark, this study found that old-growth forest stores large carbon stocks, implying the maintaining protection of old broadleaved forests is crucial not only for increasing biodiversity dead wood but also for carbon storage. In the specific case of Dalby Söderskog, effects of tree diseases have temporarily increased the relative share of dead wood compared to live tree volumes, in particular for smaller diameter trees. The mixed tree species composition of the forest has, however, buffered the effects of Dutch elm disease and ash dieback, and living tree volume still remains considerably larger than the volumes of dead wood.

5. References

- Angst, G., Messinger, J., Greiner, M., Häusler, W., Hertel, D., Kirfel, K., et al. (2018). Soil organic carbon stocks in topsoil and subsoil controlled by parent material, carbon input in the rhizosphere, and microbial-derived compounds. *Soil Biology and Biochemistry*, vol. 122, pp. 19–30.
- Badalamenti, E., Battipaglia, G., Gristina, L., Novara, A., Rühl, J., Sala, G., Sapienza, L., Valentini, R., and La Mantia, T. (2019). Carbon stock increases up to old growth forest along a secondary succession in Mediterranean island ecosystems. *PLoS ONE*, vol. 14, pp. 1-13.
- Birdsey, R., and Pan, Y. (2015). Trends in management of the world's forests and impacts on carbon stocks. *Forest Ecology and Management*, vol. 355, pp. 83-90.
- Bond-Lamberty, B., Peckham, S.D., Ahl, D.E., and Gower, S.T. (2007). The dominance of fire in determining carbon balance of the central Canadian boreal forest. *Nature*, vol. 450, pp. 89–92.
- Bond-Lamberty, B., and Gower, S.T. (2008). Decomposition and fragmentation of coarse woody debris: revisiting a boreal black spruce chronosequence. *Ecosystems*, vol. 11, pp. 831–840.
- Boyd, I.L., Freer-Smith, P., Gilligan, C.A., and Godfray, H.C. (2013). The consequence of tree pests and diseases for ecosystem services. *Science*, vol. 342.
- Bradford, J.B., Birdsey, R., Joyce, L.A., and Ryan, M.G. (2008). Tree age, disturbance history, and carbon stocks and fluxes in subalpine Rocky Mountain forests. *Global Change Biology*, vol. 14, pp. 2882-2897.
- Brunet, J., and Isacsson, G. (2009). Restoration of beech forest for saproxylic beetles-effects of habitat fragmentation and substrate density on species diversity and distribution. *Biodiversity and Conservation*, vol. 18(9), pp. 2387–2404.
- Brunet, J., Bukina Y., Hedwall, P.O., Holmström, E., and Oheimb, G. von (2014). Pathogen induced disturbance and succession in temperate forests: Evidence from a 100-year data set in southern Sweden. *Basic and Applied Ecology*, vol. 15, pp. 114-121.

- Bujoczek, L., Szewczyk, J., and Bujoczek, M. (2018). Deadwood volume in strictly protected, natural, and primeval forests in Poland. *European Journal Forest Resource*, vol. 137, pp. 401–418.
- Bukina, Y. (2012). Long-term succession and loss of foundation species in a temperate broadleaved forest in southern Sweden. Master Thesis. Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences. Alnarp.
- Burrascano, S., Keeton, W.S., Sabatini, F.M., and Blasi, C. (2013). Commonality and variability in the structural attributes of moist temperate old-growth forests: A global review. *Forest Ecology and Management*, vol. 291, pp. 458-479.
- Canadell, J.G., and Raupach, M.R. (2008). Managing forests for climate change mitigation. *Science*, vol. 320, pp. 1456-1457.
- Chatterjee, A., G. F., Vance, and D. B. Tinker. (2009). Carbon pools of managed and unmanaged stands of ponderosa and lodgepole pine forests in Wyoming. *Canadian Journal of Forest Research*, vol. 39, pp. 1893-1900.
- Christensen, M., Hahn, K., Mountford, E.P., Odor, P., Standovar, T., Rozenbergar, D., Diaci, J., Wijdeven, S., Meyer P., et al. (2005). Dead wood in European beech (Fagus sylvatica) forest reserves. *Forest Ecology Management*, vol. 210, pp. 267-282.
- Clark, D. B., and J. R. Kellner. (2012). Tropical Forest biomass estimation and the fallacy of misplaced concreteness. *Journal of Vegetation Science*, vol. 23, pp.1191-1196.
- Dittmann, S., Thiessen, E., and Hartung, E. (2017). Applicability of different noninvasive methods for tree mass estimation: A review. *Forest Ecology and Management*, vol. 398, pp. 208-215.
- Ekoungoulou, R., Niu, S.K., Loumeto, J.J., Ifo, S.A., Bocko, Y.E., Mikieleko, F.E.K., Guiekisse, E.D.M., Senou, H., and Liu, X.D. (2015). Evaluating the carbon stock in above- and below-ground biomass in a moist central African forest. *Applied Ecology and Environmental Sciences*, vol. 3, pp. 51-59.
- Emborg, J., Christensen, M., and Heilmann-Clausen, J. (2000). The structural dynamics of Suserup Skov, a near-natural temperate deciduous forest in Denmark. *Forest Ecology and Management*, vol. 126, pp. 173-189.
- Fei, S., Morin, R.S., Oswalt, C.M., and Liebhold, A.M. (2019). Biomass losses resulting from insect and disease invasions in US forests. *Proceedings of* the National Academy of Sciences, vol. 116, pp. 17371 – 17376.
- Feldhusen, K., Hammarskjöld, G., Mjureke, D., Pettersson, S., Sandberg, T., Staaf, H., Svensson, U., and Österberg, K., (2004). Sweden's National Inventory

Report 2004, Submitted under the United Nations Framework Convention on Climate Change, *Swedish Environmental Protection Agency*.

- Fraxigen, (2005). Ash Species in Europe: Biological Characteristics and Practical Guidelines for Sustainable Use. Oxford Forestry Institute, University of Oxford, UK. pp. 128.
- Ghosh, S. (2018). State of Forest Report says that India's forest and tree cover has increased by 1 percent. Retrieved from https://india.mongabay.com/2018/02/16/state-of-forest-reportsays-that-indias-forest-and-tree-cover-has-increased-by-1-percent/.
- Goodale, C. L., Apps, M. J. Birdsey, R. A., Field, C. B., Heath, L. S., Houghton, R. A., et al. (2002). Forest carbon sinks in the Northern hemisphere. *Ecological Applications*, vol. 12(3), pp. 891–899.
- Grove, S.J., Stamm, L., and Barry, C. (2009). Log decomposition rates in Tasmanian *Eucalyptus obliqua* determined using an indirect chronosequence approach. *Forest Ecology and Management*. vol. 258, pp. 389–397.
- Hadden, D., and Grelle, A. (2017). Net CO2 emissions from a primary boreonemoral forest over a 10 year period. *Forest Ecology and Management*, vol. 398, pp. 164-173.
- Harmon, M.E., Franklin, J.F., Swanson, F.J., Sollins, P., Gregory, S.V., Lattin, J.D., et al. (1986). Ecology of coarse woody debris in temperate ecosystems. *Advances in Ecological Research*, vol. 15, pp. 133-302.
- Harmon, M. E., and Sexton, J. (1996). Guidelines for measurements of woody detritus in forest ecosystems, University of Washington, pp. 73.
- Herrero, C., Krankina, O., Monleon, V., and Bravo, F. (2014). Amount and distribution of coarse woody debris in pine ecosystems of north-western Spain, Russia and the United States. *Iforest Biogeosciences and Forestry*, vol.7, pp. 53-60.
- Hess, D., and D, Tasa. (2011). McKnight's physical geography: a landscape appreciation. *Prentice Hall*.
- Houghton, R. A. (2007). Balancing the global carbon budget. *Annual Review of Earth and Planetary Sciences*, vol. 35, pp. 313-347.
- Houghton, R. A., F. Hall, and S. J. Goetz. (2009). Importance of biomass in the global carbon cycle. *Journal of Geophysical Research, Biogeosciences*, vol. 114, pp. 1-13.

- IPCC, (2001). The Scientific Basis: Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, New York, USA.
- IPCC, (2006). Guidelines for National Greenhouse Gas Inventories, *Cambridge University Press*, Cambridge.
- Karjalainen, L., and Kuuluvainen, T. (2002) Amount and diversity of coarse woody debris within a boreal forest landscape dominated by *Pinus sylvestris* in Vienansalo wilderness, eastern Fennoscandia. *Silva Fennica*, vol. 36, pp. 147–167.
- Keith, H., Mackey, B.G. and Lindenmayer, D.B. (2009). Re-evaluation of forest biomass carbon stocks and lessons from the world's most carbon-dense forests. *Proceedings of the National Academy of Sciences*, vol. 106(28), pp.11635-11640.
- Kenina, L., I. Jaunslaviete, L. Liepa, D. Zute, and A. Jansons. (2019). Carbon pools in old-growth Scots pine stands in hemiboreal Latvia. *Forests*, vol.10, pp. 1-10.
- Keßler, M., Cech, T.L., Brandstetter, M., and Kirisits, T. (2012). Dieback of ash (*Fraxinus excelsior* and *Fraxinus angustifolia*) in Eastern Austria: Disease development on monitoring plots from 2007 to 2010. Journal of Agricultural Extension and Rural Development, vol. 4(9), pp. 223-226.
- Krankina, O.N., Harmon, M.E., Kukuev, Y.A., Treyfeld, R.F., Kashpor, N.N, Kresnov, V.G., et al. (2002). Coarse woody debris in forest regions of Russia. *Canadian Journal of Forest Research*, vol. 32, pp. 768-778.
- Kumar, M., Kumar, R., Konsam, B., Sheikh, M.A., and Pandey, R. (2019). Aboveand belowground biomass production in *Pinus roxburghii* forests along altitudes in Garhwal Himalaya, India, *Current Science*, vol. 116(9).
- Kuuluvainen, T., and Gauthier, S. (2018). Young and old forest in the boreal: critical stages of ecosystem dynamics and management under global change. *For Ecosystem*, vol. 5(26).
- Lamlom, S. H., and Savidge, R. A. (2003). A reassessment of carbon content in wood: variation within and between 41 North American species. *Biomass* and Bioenergy, vol. 25, pp. 381-388.
- Lindquist, B. (1938). Dalby Söderskog, en skånsk lövskog i forntid och nutid. *Acta Phytogeographica Suecica*, vol. 10, pp. 1-273.
- Liski, J., Lehtonen, A., Palosuo, T., Peltoniemi, M., Eggers, T., Muukkonen, P., and Mäkipää, R. (2006). Carbon accumulation in Finland's forests 1922– 2004 – an estimate obtained by combination of forest inventory data with

modelling of biomass, litter and soil. *Annals of Forest Science*, vol.63, pp. 687-697.

- Liu, J., Chen, J.M., Cihlar, J., and Park, W.M. (1997). A process-based boreal ecosystem productivity simulator using remote sensing inputs. *Remote Sensing of Environment*, vol. 62, pp. 158-175.
- Luyssaert, S., Schulze, E., Börner, A., Knohl, A., Hessenmöller, D., Law, B.E., Ciais, P., and Grace, J. (2008). Old-growth forests as global carbon sinks. *Nature*, vol. 455, pp. 213-215.
- Malmer, N., Lindgrem, L., and Persson, S. (1978). Vegetational succession in a south Swedish deciduous wood. *Vegetation*, vol. 36, pp. 17-29.
- Martin, J. A., Fuentes-Utrilla, R., Gil, L., and Witzell, J. (2010). Ecological factors in Dutch elm disease complex in Europe - a review. *Ecological Bulletins*, vol. 53, pp. 209-224.
- McKinney, L.V., Nielsen, L.R., Collinge, D.B., Thomsen, I.M., Hansen, J.K. and Kjær, E.D. (2014). The ash dieback crisis: genetic variation in resistance can prove a long-term solution. *Plant Pathology*, vol. 63, pp. 485–499.
- McKinney, L.V., Nielsen, L.R., Hansen, J.K., and Kjær, E.D. (2011). Presence of natural genetic resistance in *Fraxinus excelsior (Oleaceae)* to *Chalara fraxinea (Ascomycota)*: an emerging infectious disease. *Heredity*, vol.106. pp. 788–797.
- Mendoza-Ponce, A., and Galicia, L. (2010). Aboveground and belowground biomass and carbon pools in highland temperate forest landscape in Central Mexico. *Forestry*, vol. 83, pp. 497-506.
- Muukkonen, P. (2007). Generalized allometric volume and biomass equations for some tree species in Europe. *European Journal of Forest Research*, vol.126, pp. 157-166.
- Neumann, M., Moreno, A., Mues, V., Härkönen, S., Mura, M., Bouriaud, O., Lang, M., Achten, W.M., Thivolle-Cazat, A., Bronisz, K., Merganic, J., Decuyper, M., Alberdi, I., Astrup, R., Mohren, F., and Hasenauer, H. (2016). Comparison of carbon estimation methods for European forests. *Forest Ecology and Management*, vol. 361, pp. 397-420.
- Nilsson, S.G., Niklasson, M., Hedin, J., Aronsson, G., Gutowski, J.M., Linder, P., Ljungberg, H., Mikusinski, G., and Ranius, T. (2002). Densities of large living and dead trees in old-growth temperate and boreal forests. *Forest Ecology Management*, vol. 161, pp. 189-204.
- Nord-Larsen, T., Vesterdal, L., Bentsen, N.S., and Larsen, J.B. (2019). Ecosystem carbon stocks and their temporal resilience in a semi-natural beech-dominated forest. *Forest Ecology and Management*, vol. 447, pp. 67-76.

- Öder, V., Petritan, A.M., Schellenberg, J., Bergmeier, E., and Walentowski, H. (2021). Patterns and drivers of deadwood quantity and variation in midlatitude deciduous forests. *Forest Ecology and Management, vol. 487*, pp. 118977.
- Oheimb, G. von., and Brunet, J. (2007). Dalby Söderskog revisited: long-term vegetation changes in a south Swedish deciduous forest. *Acta Oecologica*, vol. 31, pp. 229 242.
- Pan, Y., R. A. Birdsey, J. Fang, S. Piao, R. Houghton, P. E. Kauppi, A. Rautiainen, W. A. Kurz, et al. (2011). A large and persistent carbon sink in the world's forests. *Science*, vol. 333, pp. 988-993.
- Passovoy, M.D., and Fule, P.Z. (2006). Snag and woody debris dynamics following severe wildfires in northern Arizona ponderosa pine forests. *Forest Ecology and Management*, vol. 223, pp. 237-246.
- Pautasso, M., Aas, G., Queloz, V., and Holdenrieder, O. (2013). European ash (*Fraxinus excelsior*) dieback – A conservation biology challenge. *Biological Conservation*, vol. 158, pp. 37-49.
- Pietsch, S.A., and Hasenauer, H. (2002). Using mechanistic modeling within forest ecosystem restoration. *Forest Ecology and Management*, vol. 159, pp. 111-131.
- Pihlgren, A., Hallingbäck, T., Aronsson, M., Dahlberg, A., Edqvist, M., Johansson, G., Krikorev, M., and Thor, G. (2010). The new Swedish Red List 2010. In Swedish.
- Pinkard, E. A., Battaglia, M., Roxburgh, S., and O'Grady A. P. (2011). Estimating forest net primary production under changing climate: adding pests into the equation, *Tree Physiology*. vol. 31, pp. 686–699.
- Potter, C and Urquhart, J, (2017). Tree disease and pest epidemics in the Anthropocene: A review of the drivers, impacts and policy responses in the UK, Forest Policy and Economics, *Elsevier*, vol. 79, pp. 61-68.
- Pugh, T.A., Lindeskog, M., Smith, B., Poulter, B., Arneth, A., Haverd, V., and Calle, L. (2019). Role of forest regrowth in global carbon sink dynamics. *Proceedings of the National Academy of Sciences*, vol. 116, pp. 4382 – 4387.
- Ruks, J. (2020). Effects of ash dieback and Dutch elm disease on forest structure in Dalby Söderskog 2012-2020. Master Thesis. Southern Swedish Forest Research Centre, Swedish University of Agricultural Sciences. Alnarp.
- Seedre, M., J. Kopáček, P. Janda, R. Bače, and M. Svoboda. (2015). Carbon pools in a montane old-growth Norway spruce ecosystem in Bohemian Forest:

Effects of stand age and elevation. *Forest Ecology and Management*, vol. 346, pp. 106-113.

- Seedre, M., Shrestha, B. M., Chen, H. Y. H., Colombo, S., and Jõgiste, K. (2011). Carbon dynamics of North American boreal forest after stand replacing wildfire and clearcut logging. *J. Forest Resources Japan*, vol. 16(3), pp. 168–183.
- Siitonen, J. (2001). Forest management, coarse woody debris and saproxylic organisms, Fennoscandian boreal forests as an example. *Ecological Bulletins*, vol. 49, pp. 11-41.
- Sollins, P. (1982). Input and decay of coarse woody debris in coniferous stands in western Oregon and Washington. *Canadian Journal of Forest Research*, vol. 12, pp. 18–28.
- Spies, T.A., Franklin. J.F., and Thomas, T.B. (1988). Coarse woody debris in Douglas-fir forests of western Oregon and Washington. *Ecology*, vol. 69, pp. 1689-1702.
- Stener, L.G. (2013). Clonal differences in susceptibility to the dieback of *Fraxinus* excelsior in southern Sweden. Scandinavian Journal of Forest Research, vol. 28(3), pp. 205 – 216.
- Stenlid, J., Oliva, J., Boberg, J., and Hopkins, A.J. (2011). Emerging diseases in European forest ecosystems and responses in society. *Forests*, vol. 2, pp. 486-504.
- Sturtevant, B.R., Bissonette, J.A., Long, J.N., and Roberts, D.W. (1997). Coarse woody debris as a function of age, stand structure, and disturbance in boreal Newfoundland. *Ecological Applications*, vol 7, pp. 702–712.
- Thomas, S. C., and A. R. Martin. (2012). Carbon content of tree tissues: A synthesis. *Forests*, vol. 3, pp. 332-352.
- Vucetich, J.A., Reed, D.D., Breymeyer, A., Degórski, M., Mroz, G.D., Solon, J., Roo-Zielińska, E., and Noble, R.D. (2000). Carbon pools and ecosystem properties along a latitudinal gradient in northern Scots pine (*Pinus sylvestris*) forests. *Forest Ecology and Management*, vol. 136, pp. 135-145.
- Waller, M.P. (2013). Drought, disease, defoliation and death: forest pathogens as agents of past vegetation change. *Journal of Quaternary Science*, vol. 28, pp. 336-342.
- Waring, R.H., and Running, S.W. (2007). Forest Ecosystem Analysis at Multiple Time and Space Scales. *Elsevier Academic Press*.
- Wirth, C., and Lichstein, J.W. (2009). The imprint of species turnover on oldgrowth forest carbon balances – insights from a trait-based model of forest

dynamics, 81–113 in Wirth, C., Gleixner, G. and Heimann, M. (eds), *Old-Growth Forests*, vol. 207, pp. 81-113.

Woodall, C.W., and Westfall, J.A. (2009). Relationships between the stocking levels of live trees and dead tree attributes in forests of the United States. *Forest Ecology Management*, vol. 258, pp. 2602–2608.

Acknowledgements

Throughout the writing of this dissertation, I have received a great deal of support and assistance. My heartfelt thanks to all who assisted me in the completion of this thesis. I would like to express my utmost gratitude to my supervisor, Jörg Brunet, for his invaluable supervision, support, and tutelage during the course. I would like to thank my examiner, Per Ola Hedwall, for his thoughtful comments and suggestions for improvement. Next, I would like to express my gratitude to my family and friends for their unconditional support, encouragement, and assistance throughout this time. Finally, the European Union Erasmus Mundus Scholarship Program, which provided support for my education. Without it, I would never have been capable of having knowledgeable mesmeric ride in Europe.

Appendix



Figure A17. Distribution of downed dead wood (logs) in different diameter classes.



Figure A18. Distribution of snags in different diameter classes.



Figure A19. Distribution of stumps in different diameter classes.

Variable	Ν	Mean	SE	StDev	Minimum	Q1	Median	Q3	Maximum
			Mean						
Fagus	50	109.9	21.8	153.9	0.0	0.0	23.0	207.0	589.0
Fraxinus	50	119.0	22.8	161.2	0.0	0.0	20.5	205.6	659.9
Ulmus	50	16.0	2.8	19.6	0.0	0.0	9.9	21.0	103.0
Quercus	50	81.4	15.3	108.4	0.0	0.0	0.0	138.9	351.0
Others	50	10.3	3.1	21.8	0.0	0.0	0.0	14.0	111.5
Total volume	50	336.7	29.6	209.6	15.0	197.2	316.2	453.5	874.8

Table A4. Descriptive Statistics. Volume of live trees by species.

Variable		Ν	Mean	SE Mean	StDev	Minimum	Q1	Median	Q3	Maximum
Log m3/ha	vol	50	174.4	22.1	156.4	7.9	67.6	124.3	241.6	779.6
Snag m3/ha	vol	50	49.3	16.6	117.3	0.0	0.0	9.3	33.7	621.7
Stump m3/ha	vol	50	4.1	1.5	10.6	0.0	0.0	0.0	2.8	62.2
CWD m3/ha	vol	50	227.9	27.7	196.1	7.9	87.7	162.9	303.2	805.4
CWD biomass t	/ha	50	145.8	17.7	125.5	5.1	56.1	104.3	194.1	515.5
CWD car tC/ha	bon	50	68.5	8.3	59.0	2.4	26.4	49.0	91.2	242.3

Table A5. Summary of statistics for CWD.

Sample plot		Snag vol	Stump vol	CWD vol	Living trees	All wood		
no.	Log vol m3/ha	m3/ha	m3/ha	m3/ha	m3/ha	m3/ha		
1	117.13	7.63	0.00	124.76	342.71	467.47		
2	60.51	30.42	3.67	94.60	744.72	839.32		
3	61.24	21.24	1.45	83.93	494.09	578.02		
4	551.63	161.10	0.00	712.73	302.82	1015.54		
5	156.83	4.52	0.00	161.35	655.93	817.28		
6	70 59	6.12	0.00	76 72	461 57	538.28		
7	104.16	0.00	0.00	104.16	210.82	314 98		
,	465.81	140.60	0.00	606.41	210.02	901.99		
9	120.00	1 50	0.00	121 50	/53.18	574 77		
11	60.17	15.62	1 53	77 21	433.18 874.84	952.15		
12	19 65	10.02	0.00	07.01	200 71	100 62		
14	46.05	49.20	11.97	97.91	125.25	488.03		
14	75.69	0.00	2.50	07.77	135.55	223.12		
17	253.13	2.86	2.56	258.56	33.00	291.56		
18	216.15	44.11	0.00	260.26	809.24	1069.49		
23	110.98	13.90	10.00	134.89	416.91	551.80		
27	256.38	40.65	1.45	298.49	97.40	395.89		
28	125.15	10.51	6.98	142.64	199.97	342.62		
29	46.63	0.00	6.78	53.41	357.62	411.04		
30	241.34	15.44	0.00	256.78	634.98	891.76		
31	178.63	23.12	0.00	201.75	233.12	434.88		
32	195.37	58.74	1.45	255.55	389.44	644.99		
33	153.98	0.00	12.74	166.71	395.22	561.94		
34	96.61	174.14	0.00	270.75	421.34	692.09		
36	350.32	0.00	2.56	352.88	75.87	428.75		
39	9.23	132.85	0.00	142.09	517.54	659.62		
41	14.57	0.00	0.00	14.57	595.02	609.59		
42	38.47	0.00	0.00	38.47	337.28	375.75		
44	319.08	0.00	0.00	319.08	274.52	593.60		
45	158.83	175.94	0.00	334.78	408.99	743.77		
46	171.29	88.17	0.00	259.46	134.91	394.37		
47	471.47	0.00	62.19	533.66	340.13	873.79		
49	118.10	12.34	0.00	130.44	304.80	435.24		
51	24.02	14.50	0.00	38.52	263.23	301.75		
52	77.79	0.00	0.00	77.79	292.68	370.47		
54	158.45	6.05	0.00	164.50	366.86	531.36		
55	304.41	9.03	4.07	317.51	285.36	602.86		
57	294.77	31.35	2.46	328.58	240.90	569.48		
58	19.48	0.00	0.00	19.48	746.18	765.65		
59	76.30	2.22	0.00	78.52	220.68	299.20		
62	449.98	0.00	39.70	489.68	110.44	600.12		
64	16.80	512.18	0.00	528.98	454.57	983.55		
65	242.38	9.48	0.00	251.86	188.78	440.64		
66	123.35	3.19	6.95	133.50	57.00	190.50		
67	179.47	0.00	2.51	181.98	246.82	428.80		
68	69.76	15.10	2.51	87.37	53.54	140.91		
69	779.61	10.49	0.00	790 10	15.00	805 10		
70	86.78	0.00	1 45	88.73	79.54	167 77		
72	167.09	621 72	16.62	805.43	32.00	837.42		
72	224 55	0.00	<u> </u>	228 67	327.60	556.20		
74	7 01	0.00	0.00	7 01	51/ 01	521 02		
Total Mean	174.42	49.32	4.11	227.86	336.70	564.56		

Table A6. Wood volumes in 50 sample plots in Dalby Söderskog.

Middle dbh, cm	12.5	17.5	22.5	27.5	32.5	37.5	42.5	47.5	52.5	57.5	62.5	67.5	72.5	77.5	82.5	87.5	92.5	>97.5	Sum
Quercus, no stems	8	3	6	9	10	24	50	52	89	121	94	97	97	90	89	57	57	68	1021
Quercus, m3 sum			2	5	7	28	81	109	236	366	310	355	355	378	378	268	268	387	3533
Quercus, m3 per stem			0.33	0.56	0.70	1.17	1.62	2.10	2.65	3.02	3.30	3.66	3.66	4.20	4.25	4.70	4.70	5.69	
Fagus, no stems	362	204	99	50	51	52	62	64	72	56	56	42	41	36	36	22	21	23	1349
Fagus, m3 sum	22	35	34	31	51	71	107	146	207	193	212	181	176	180	180	121	115	175	2237
Fagus, m3 per stem	0.06	0.17	0.34	0.62	1.00	1.37	1.73	2.28	2.88	3.45	3.79	4.31	4.29	5.00	5.00	5.50	5.48	7.61	
Fraxinus, no stems	674	572	440	268	173	132	104	100	94	87	63	42	36	31	20	15	5	14	2870
Fraxinus, m3 sum	50	102	150	151	150	161	178	225	270	296	237	175	164	151	103	81	30	103	2777
Fraxinus, m3 per stem	0.07	0.18	0.34	0.56	0.87	1.22	1.71	2.25	2.87	3.40	3.76	4.17	4.56	4.87	5.15	5.40	6.00	7.36	
Ulmus, no stems	3010	1520	841	437	352	247	220	216	149	159	97	56	55	43	43	20	20	20	7505
Ulmus, m3 sum	187	232	260	236	304	310	366	479	411	520	354	227	223	202	202	104	104	131	4852
Ulmus, m3 per stem	0.06	0.15	0.31	0.54	0.86	1.26	1.66	2.22	2.76	3.27	3.65	4.05	4.05	4.70	4.70	5.20	5.20	6.55	
Other, no stems	316	120	137	103	51	33	13	5	8	5	4	1	0	0	0	0	0	2	798
Other, m3 sum	15	17	43	56	41	40	21	12	20	16	14							7	302
Other, m3 per stem	0.05	0.14	0.31	0.54	0.80	1.21	1.62	2.40	2.50	3.20	3.50							3.50	

Table A7. Summary of the complete tree inventory in Dalby Söderskog in 1970 (Lindgren 1971). Calculated volumes include stems and branches down to 3 cm at the top.