



Swedish University of Agricultural Sciences
Faculty of Veterinary Medicine and Animal Science

Fine mapping of autoimmune diseases in chicken

Utlwanang Mosupiemang



Swedish University of Agricultural Sciences
Faculty of Veterinary Medicine and Animal Science
Department of Animal Breeding and Genetics

Fine mapping of autoimmune diseases in chicken

Utlwanang Mosupiemang

Supervisors:

Susanne Kerje, UU, Sweden

Johann Sölkner, BOKU, Austria

Examiner:

Göran Andersson, SLU, Department of Animal Breeding and Genetics

Credits: 30 HEC

Course title: Degree project in Animal Science

Course code: EX0556

Programme: Erasmus Mundus Programme - European Master in Animal Breeding and Genetics

Level: Advanced, A2E

Place of publication: Uppsala

Year of publication: 2012

Name of series: Examensarbete / Swedish University of Agricultural Sciences,
Department of Animal Breeding and Genetics, 384

On-line publication: <http://epsilon.slu.se>

Key words: Chicken, autoimmune disease, animal models, systemic sclerosis, vitiligo

CONTENTS

ABSTRACT	1
BACKGROUND.....	2
The chicken as an animal model	2
Chicken models of autoimmune diseases.....	2
<i>Chicken model of systemic sclerosis</i>	2
<i>Other animal models of SSc</i>	3
<i>Chicken model of vitiligo</i>	4
<i>Other animal models of vitiligo</i>	4
<i>Systemic sclerosis</i>	5
<i>Cytokines and chemokines (receptors) in scleroderma</i>	8
<i>Suppressor of cytokines signalling 1 (SOCS 1)</i>	8
<i>Chemokine ligand 1 (CCL1) and chemokine receptor 8 (CCR8)</i>	8
<i>Treatment</i>	11
THE CHICKEN GENOME	11
MAPPING STRATEGIES	12
<i>Linkage Mapping</i>	12
PRELIMINARY RESULTS	13
MATERIALS AND METHODS	14
Animals	14
UCD 200 pedigree.....	14
Phenotyping.....	14
Blood sampling and DNA extraction	15
Sequencing for detection SNPs in potential candidate genes	15
Genotyping using pyrosequencing	16
STATISTICAL ANALYSIS	16
<i>Single Marker Analysis</i>	16
RESULTS.....	17
SNP identification in candidate genes for systemic sclerosis	17
Pyrosequencing	17
QTL confirmation in the SLxBL F3-5 generations.....	17
DISCUSSION	19
FUTURE PROSPECTS	20
CONCLUSION	21
ACKNOWLEDGEMENTS	21
APPENDIX	22
SNPs in QTL used in pyrosequencing	28
<i>Chromosome 2 marker RS13673471</i>	28
<i>Chromosome 12 marker RS14975805</i>	28
<i>Chromosome 14 marker RBL1140</i>	28
<i>Chromosome 19 marker RS14123499</i>	28
R script	28
REFERENCES.....	29

ABSTRACT

Autoimmune diseases arise due to tolerance breakdown or failure of the body's immune system to establish tolerance leading to the body failing to recognize its own tissues and organs as self. Systemic sclerosis is a disease of the connective tissue characterized by immunological abnormalities and increased deposition of extracellular matrix proteins in the skin. Vitiligo is a dermatological disorder characterized by loss of melanocytes from the cutaneous epidermis. The aim of this project was to confirm the previously identified QTLs in the Smyth line (SL) chicken model of vitiligo in F₃-F₅ generations of the SL and Brown line (BL) cross and systemic sclerosis in the University of California at Davies (UCD 200) model in the 2nd backcross generations (BC₂) of UCD 200 and red jungle fowl cross (RJF). We confirm the association of the disease with our marker genotypes on chromosome 3 but not on chromosomes 9 and 21 in the SL and BL crosses from F₃-F₅ generations. We did not confirm the disease association with chromosomes 2, 12, 14 and 19 in the BC₂ generation from the UCD x RJF cross because the pedigree was not large enough for the analysis. 68% of the individuals in the BC₂ pedigree expressed the diseases with males showing 100% liability. We recorded a mortality rate of approximately 50% in the BC₂ population. We did not observe any significant mortality in the intercrosses between the SL and BL.

Keywords: chicken, autoimmune disease, animal models, systemic sclerosis, vitiligo

BACKGROUND

Chickens and humans shared a common ancestor approximately 310 million years ago (Hedges 2002, Hillier *et al.* 2004 and Burt 2005). The red jungle fowl (*Gallus gallus*) is considered to be the ancestor of all domestic chickens (West and Zou 1988, Fumihito *et al.* 1994, Wong 2001, Kanginakudru 2008). The chicken (*Gallus gallus domesticus*) was domesticated over 8 000 years ago in South East Asia. It was originally used for religious purposes, leisure, cock fighting and the Romans used the chicken as oracles (Crawford 1996). It was later used as an agricultural animal with specialized breeds generated for different purposes. It is a source of protein (meat and eggs) with billions of eggs used each year (Burt 2005). Some chickens are kept as pets such as the Bantam and the Silkies since they are more docile (Clucks & Chooks 2008).

The chicken as an animal model

Animal models are important in providing clues for understanding human diseases. Evidence of the use of chicken as an animal model described by Aristotle was the use of chicken in embryology. Harvey (1628) studied blood circulation using chickens while Pasteur (1880) discovered transmission of infections in chickens through attenuating a virulent pathogen of chicken cholera. Today, the chicken is an important experimental system for developmental biology, immunology, microbiology, genetics, virology and cancer (Jimenez *et al.* 1984, Brown *et al.* 2003, International Chicken Genome Sequencing Consortium 2004 and Stern 2005). Animal models for understanding diseases were based on whether the diseases are induced by exogenous compounds or genetically inherited as a trait (Van de Water *et al.* 1995).

Experimental animal models that spontaneously develop autoimmune diseases reflect the human situation more closely compared with models where the disease has been induced (Erf 2010). Even though murine models have been broadly used in biomedical research for autoimmune diseases they are very rare and currently there is no model that spontaneously develops autoimmune vitiligo (Erf 2010).

The chicken has several advantages as an animal model for studies of biological processes and they include a rich genetic diversity, large population size, ease of breeding, high rates of recombination (Kerje 2003, Siegel *et al.* 2006) and this has been enhanced by the construction of the chicken genome sequence and identification of 2.8 million SNPs (International Chicken Genome Sequencing Consortium 2004 and International Chicken Polymorphism Map Consortium 2004).

Chicken models of autoimmune diseases

Chicken model of systemic sclerosis

Complex pathophysiology of scleroderma/systemic sclerosis (SSc) has led to the development of animal models, which gives a closer look at pathophysiology of the disease such as

functions of various cell types like monocytes, T-lymphocytes, fibroblasts and mast cells (Haustein 2002).

The University of California at Davies lines 200 and 206 (UCD 200 and 206) developed by Bernier in 1942 have become ideal models for the study of SSc since they show the entire manifestation of clinical, histopathological and serological spectrum of SSc while other animal models (tight skin mice) of SSc lack vascular damage (Sgonc 1999 and Wick *et al.* 2006). In this chicken model, F₁ generation show no incidence of the disease while F₂ generation had a disease incidence of 4%.

Other animal models of SSc

Other animal models of SSc include the tight skin mouse (*Tsk*). The *tsk* mouse has thickened skin that is tightly bound to the subcutis and deep muscular tissue. Compared to human scleroderma, the *tsk* mice lack dermal sclerosis and fibrosis is deeper than in humans but biochemical and molecular abnormalities similar to human SSc were noted (Yamamoto 2010).

In bleomycin-induced scleroderma, bleomycin (an isolate from fungus *Streptomyces verticulus*) is used. Bleomycin is usually used as an anti-tumour antibiotic and has several biochemical properties which include blocking the cell cycle at G₂, produces free radicals that induces apoptosis and exerts effects on skin-constituted cells such as fibroblasts, keratinocytes, endothelial cells and immunocytes. It also up-regulates collagen and TGF- β mRNA expression (in lungs of rats and skin fibroblasts of humans) and augments gene expression of extra cellular mass proteins (ECM) together with fibrotic cytokines (Yamamoto 2000; 2009; 2010).




Fig. 1. Possible pathogenesis of bleomycin induced systemic sclerosis. MMP is matrix metalloproteinase NO is nitric oxide. Yamamoto 2010.

In the fibrosis mouse model of TGF- β /CTGF, mice injected with TGF- β showed granulation and fibrosis of the skin (Yamamoto 2005). TGF is a powerful stimulus for *in vitro* formation of collagen and induces fibrosis and angiogenesis in newborn mice. It is suggested that TGF- β has an important role in inducing granulation and fibrotic tissue formation whereas CTGF maintains fibrosis (Kanzler *et al.* 1999; Takehara 2003)

Chicken model of vitiligo

The Smyth line (SL), developed by Dr J Robert Smyth in the 1970s is well established as an animal model of vitiligo (Smyth 1980). SL chickens show post hatch loss of melanin synthesizing melanocytes in feather and choroid tissue. The SL chicken is the only animal model that mimics the entire spectrum of clinical and biological manifestations of the corresponding human disease (Wick *et al.* 2006).

Liability of vitiligo in the Smyth line is 70-95% within the contemporary group and de-pigmentation in the feathers is seen at 6-20 weeks of age with animal showing complete de-pigmentation in adulthood (Erf 2007). However, the parental Brown line (BL101) with matched major histocompatibility complex (MHC) has less than 2% incidence of vitiligo while the Light Brown (LBL101) is resistant against development of vitiligo.

Incidences of the disease in this animal model are predictable and the lesions are easily accessible since the feather tissue regenerates. Smyth (1989) characterized the disorder as spontaneous vitiligo-like loss of melanin producing cells in the feathers and the choroid after hatching.

Other animal models of vitiligo

Several species having naturally occurring de-pigmentation have been proposed as models for vitiligo. One example is *Grey* horses such as Lipizzaner, Arabian and Andalusian. These horses are normally coloured at birth and those carrying the dominant *Grey* allele show de-pigmentation after birth which increases with age until they appear completely white at maturity (Boissy and Lamoreux 1988), nevertheless a duplication of the *STX17*-region was denoted to be the causative mutation leading to premature loss of pigment and melanoma in these horses (Pielberg *et al.* 2008), and therefore this is not a good model for autoimmune disease as no autoimmune component is involved.

The Sinclair miniature swine model of vitiligo develops spontaneous regression of cutaneous malignant melanoma lesions (Hook *et al.* 1982). This pig is considered a good model for spontaneous melanoma. De-pigmentation is associated with the tumours together with development of uveitis and vitiligo and there is evidence of immune system involvement in tumour regression and development of vitiligo (Erf 2010). Vitiligo in the water buffalo was observed to resemble vitiligo in humans in cytological aberrations of melanocytes but no autoimmune component has been confirmed (Cerundolo *et al.* 2007).

The vitiligo mouse (*C57Bl/6^{vit/vit}*) has in addition to progressive loss of hair and ocular pigmentation with age also contact allergens as observed in human vitiligo (Boissy *et al.* 1987). However, the phenotype was mapped to the microphthalmia (*MITF*) locus (*mi^{vit/vit}*) (Lamoureux *et al.* 1992) and therefore this animal model was excluded as a model for autoimmune vitiligo.

The Barred Plymouth Rock and White Leghorn chicken lines were thought to be

models of vitiligo (Bowers *et al.* 1992). In Barred Plymouth Rock chickens, the black and white band patterns results from autophagocytic degeneration of melanocytes, which are not present in the white bands. The *barring* phenotype was lately linked to two non-coding mutations in the *CDKN2A* and two missense mutations in a highly conserved region (Hellström *et al.* 2010). The *barring* gene was found to inhibit melanin deposition in the feathers, beak, shanks and the eyes of the chicken (Erf 2010). The white feathers in the White Leghorns are known to be caused by a mutation in the *PMEL17* gene (Kerje *et al.* 2004). Due to the fact that all de-pigmentation phenotypes described above are caused by mutations in single genes eliminates all of them as models for autoimmune vitiligo.

AUTOIMMUNE DISEASES

Autoimmune diseases arise as a result of failure of the body to recognize its own cells and tissues allowing the immune system to respond against something which is usually not a threat or non-foreign. This normally happens when immune tolerance is broken or not well established. Immunological tolerance is the failure to mount an immune response to an antigen which helps the body not to attack its own tissues and other antigens (<http://www.herb.co.za/herb-remedies/immune-system01.htm>).

Central tolerance is a key feature in autoimmune diseases. It is a mechanism by which developing T-cells and B-cells are rendered non-reactive to healthy tissues and it is induced at the primary site of the lymphocytes (Hogquist *et al.* 2005). Self antigens are transported to the thymus where clonal deletion or tolerance of self-reactive thymocytes is induced (Sprent 1995). Autoimmune disorders fall into two categories: systemic, which damages many organs and localized which attacks specific organs and tissues even though it can extend beyond the targeted tissue to other body organs (Ravel and Descotes 2004 and <http://www.aacc.org>). The precise cause of autoimmune diseases is generally unknown but it is suggested that there appears to be some genetic predisposition to develop autoimmune disease in many cases and environmental factors have a role in triggering the disorders.

Systemic sclerosis

Systemic sclerosis is a heterogeneous disorder that affects the connective tissue of the skin and internal organs such as the lungs, intestinal track, heart and kidney (Haustein 2002). It is characterized by vasomotor disturbances, fibrosis, subcutaneous and skin atrophy as well as muscle and arterial occlusion of internal organs, polyarthritis and presence of antinuclear antibodies, peri-vascular mononuclear cell-infiltration and increased deposition of extra cellular matrix, microvascular damage, immune system disturbance and massive collagen deposition (Gershwin *et al.* 1981, Sgonc 1999, Yamamoto 2002, Jimenez and Derk 2004, Schwartz *et al.* 2009, Rabquer *et al.* 2009).

Aetiology of SSc is assumed to be immunologic mechanisms, genetic risk factors and environmental triggers as well as interaction between genetic risk factors and environmental triggers (Jin *et al.* 2007, Rula and Hajj-Ali 2008, Maron *et al.* 2010). In humans, familial clustering of SSc is rare (Van de Water and Gershwin 1985) and this might be because familial cases of the disease have not been well documented or environmental factors are very

important and this can be proven in the UCD 200 and 206 lines since they have been heavily selected for the disease.

In humans, SSc is characterized by fatigue, intolerance to cold, pain, deformity and loss of function of the hand as well as change in facial appearance (Medsger 1991). Pathohistological findings indicated that fibroblasts, endothelial cells and lymphocytes are key main players in three major clinical observations which are fibrosis of cutaneous and viscera, vascular alterations and immunological alterations (Wick *et al* 2006, Yamamoto 2005, 2009). Progressive thickening and fibrosis of the skin is usually noted and in subclinical cases internal organs are affected (Fleischmajer *et al.* 1977, Haustein 2002 and Wick *et al.* 2006). In humans, SSc starts with oedema of the fingers preceded by Raynaud's phenomenon (Yamamoto 2009). Raynaud's phenomenon is discolouration of the fingers or toes due to reduced blood flow caused by vasoconstriction. Fibrosis can be caused by profibrotic cytokines including TGF-beta, interleukin 4 (IL-4), and platelet derived growth factor (PDGF) (<http://www.aacc.org>).

In the UCD 200 chickens, SSc is characterized by swelling, erytheroma and necrosis of the comb (self dubbing) (Fig. 1) (Van de Water and Gershwin 1985). Van de Water and Gershwin (1985) also observed swelling of the peripheral joints as well as polyarthritis, swelling of the upper neck sac after four weeks of age. Other visceral organs involved include oesophagus (64%), small intestine (60%), lung (60%), kidney (65%), heart (30%) and testis (66%).

Antinuclear-antibodies have been detected in 65% of the UCD 200 chickens at six months of age, and it was noted that such chickens also produced rheumatoid factor at four months of age (Van de Water and Gershwin 1985).

A comparative study of skin biopsies from UCD 200 chickens and biopsies from human SSc patients revealed that endothelial cell apoptosis induced by anti-endothelial cell antibody (AECA) - dependent cellular cytotoxicity is a principal event in the pathogenesis of SSc (Sgonc 1999). Histopathological findings of the skin involve inflammatory infiltrates of lymphocytes, monocytes, plasma, histiocytes around the blood vessels and sweat glands. Infiltrating immunocytes releases fibrogenic cytokines, which play a very crucial role in the initiating events in SSc (Jimmenez *et al.* 1986). Comparative details of SSc in both humans and chickens are described in Table 1.



Fig. 2. UCD 200 chickens showing swelling and necrosis of the comb (Photos. S Kerje)



<http://www.goldbamboo.com/pictures-t7460.html>

Fig. 3. Systemic sclerosis patients showing polyarthritis in fingers equivalent to feet necrosis in chickens.

Treatment

Several vitamins, pharmaceuticals, hormones, drugs and surgical procedures are reported to be successful in the treatment of systemic sclerosis and these treatments have been well accepted by the public (Medsger 1991).

Table 1. Comparison of clinical, biochemical, immunological, and pathologic features of human and avian SSc, source Wick et al. 2006.

	Human SSc	Avian SSc
Clinical features		
<i>Disease presentation</i>	Subtle, middle age	Acute, early in life
<i>Skin fibrosis</i>	Present	Present
<i>Esophageal fibrosis</i>	Present	Present
<i>Lung fibrosis</i>	Present	Present
<i>Kidney involvement</i>	Present	Present
<i>Heart involvement</i>	Present	Present
<i>Polyarthritis</i>	Present	Present
<i>Eye involvement</i>	Debated	Absent
Autoimmune features		
<i>Autoantibodies</i>		
ANA	Present	Present
Anti-Scl-70	Present	Absent
Anticentromere	Present	Present
Anticytoplasmic	Present	Present
<i>Autoreactive T cells</i>	Present	Present
Aetiology		
<i>Genetic susceptibility</i>	Necessary not sufficient	Necessary and sufficient

<i>Environmental factors</i>	Hypothesized	Not important
Pathology		
<i>Fibroblast alterations</i>	Present	Present
<i>Endothelial alterations</i>	Present	Present
<i>Smooth muscle alterations</i>	Present	Present

Cytokines and chemokines (receptors) in scleroderma

Cytokines are a large family of glycoproteins that regulates fundamental biological processes, which includes immunity, wound healing, haematopoiesis and development of the nervous system and they are essential components of adaptive and innate immune systems (Krebs and Hilton 2001; Alexander and Hilton 2003). Chemokines are a small family of cytokines that participate in immune and inflammatory responses through chemo-attraction and activation of leukocytes (Baggiolini 1998). Chemokines function as leukocyte activators and chemo-attractants in inflammatory responses (Luster 1998). In aetiology of scleroderma, there is growing evidence that overproduction of extracellular matrix by activated fibroblasts results from the interaction between endothelial cells, macrophages, lymphocytes and fibroblasts through mediators such as cytokines, chemokines and growth factors (Yamamoto 2009).

Suppressor of cytokines signalling 1 (SOCS 1)

The suppressor of cytokine signalling 1 is a key negative regulator of cytokine signalling and immune response (Egan *et al.* 2003, Hanada *et al.* 2003, Fenner *et al.* 2006). The SOCS1 family of proteins forms part of the negative feedback system that regulates cytokine signalling transduction by binding to the cell surface receptors and activating intracellular signal transduction such as the JAK-STAT pathway (Krebs and Hilton 2001; Alexander and Hilton 2004). The SOCS1 protein target signal transducers for proteosomal destruction (Alexander and Hilton 2003) and expression can be induced by a subset of cytokines IL-1, IL-3, erythropoietin and interferon gamma (Hanada *et al.* 2003).

Chemokine ligand 1 (CCL1) and chemokine receptor 8 (CCR8)

CCL1 is a small glycoprotein secreted by activated T cells belonging to a family of chemokines. CCL1 attracts monocytes, NK cells, immature B cells and dendritic cells by interacting with a cell surface of chemokine receptors, CCR8 (Millner and Krangel 1992, Roos *et al.* 1997, Hapel and Haque 2002, Haque *et al.* 2004). CCR8 is a G-protein-coupled receptor that is unique for the CCL1 and viral monocyte inflammatory protein-1. It was found to be a functional endothelial receptor mediating endothelial chemotaxis in response to CCL1 and vCCL1 (Dairaghi *et al.* 1999).

Other cytokines involved in scleroderma include TGF- β , released by activated macrophages or lymphocytes, chemically attracts fibroblasts and increases the synthesis of extracellular mass *e.g.* collagen type I and type III and fibronectin (Yamamoto 2009). TGF- β increases levels of TGF- β receptors in fibroblasts leading to increased TGF- β production and hence high deposition of extracellular mass and fibrosis (Yamamoto 2009).

Connective tissue growth factor (CTGF) is induced in fibroblasts after activation by TGF- β (Frazier *et al.* 1996, Yamamoto 2009). CTGF protein stimulates DNA synthesis and up-regulates collagen, fibronectin and integrin expression in fibroblasts (Frazier *et al.* 1996).

An existing disproportion between type I and II cytokine responses have been reported in the pathogenesis of scleroderma. Interleukin 13 (IL-13) has the capability to suppress pro-inflammatory cytokines production in monocytes and macrophages and enhance the growth and differentiation of B cells and promote immunoglobulin synthesis (Yamamoto 2009).

Pro-inflammatory chemokines are associated with the initiation of fibrosis and sclerosis. This led to the suggestion that chemokines and chemokine receptors may have a critical role as mediators of inflammation and fibrosis (Yamamoto 2009).

Autoimmune vitiligo

Vitiligo is an acquired dermatological disorder characterized by loss of epidermal melanocyte leading to de-pigmentation of the skin (Wang and Erf 2004). It is thought that vitiligo is due to a combination of several factors which, include genetic susceptibility, immune system and environmental components including several kinds of stress (emotional) accumulation of toxic melanin precursor in melanocyte (DOPA, dopachrome, 5 6-dihydroxyindole) and disturbance of melanocyte homeostasis (impaired intra and extra cellular calcium) (Wick *et al.* 2006 and Kakourou 2009).

The disease is known to affect up to 2% of the world population (Erf *et al.* 2001, Namazi 2005, Wick *et al.* 2006 and <http://www.aocd.org>). Vitiligo is often associated with other autoimmune diseases such as hypothyroidism (4-8%), alopecia areata (2-3%) and 5-15% uveitis often resulting in blindness. People with vitiligo are more prone to develop Addison's disease, pernicious anaemia and diabetes mellitus, type I (Erf 2010).

The disorder usually appears as focal pattern where the disease is limited to one or few areas, segmental pattern where de-pigmentation occurs only in one side of the body and as generalized pattern which is the most common and occurs symmetrically in both parts of the body.



Fig. 4. Patients with vitiligo showing loss of pigmentation. Photos (<http://www.psoriasiscafe.org/vitiligo-pictures.htm>)

In both chickens and humans, vitiligo is characterized by spontaneous loss of pigmentation in association with uveitis and underlying hair as well as oral mucosa (Lamont *et al.* 1982 and Alkhateeb *et al.* 2003). There is an increased frequency of association of vitiligo and several antibodies suggesting hyperactive immune system in humans and in some patients autoantibodies are directed against melanocytes or melanocyte proteins (Lamont *et al.* 1982

Alkhateeb *et al.* 2003). Evidence of strong association with anti-melanocyte antibodies in both animals and humans has been reported (Norrirs *et al.* 1988). An intrinsic heritable melanocyte defect with abnormal and irregular surfaces predisposes the SL chickens to the pigment disorder (Boissy *et al.* 1983).

The immunopathology of Smyth line vitiligo has been shown to support the involvement of cell mediated immune response in destruction of the melanocytes causing white feathers in the chickens (Fig. 4) (Wang and Erf 2003; Wang *et al.* 2004). The environmental component in the expression of autoimmune vitiligo was shown by vaccination of chickens with a live Turkey herpes virus vaccine (HVT), which triggered the expression of SL vitiligo (Erf *et al.* 2003). HVT show a strong tropism for epithelium of feather follicles, however some studies have revealed that killed HVT had no effect in the expression of vitiligo (Erf *et al.* 2001, 2003). Vaccinations with Newcastle disease virus (NDV) and Infectious bronchitis virus (IBV) did not trigger vitiligo suggesting that viral infections and antiviral immune activity are not responsible for the expression of autoimmune vitiligo (Erf 2010).



Fig. 5. Phenotypes in SL x BL intercross chickens with vitiligo (Photos. S. Kerje).

Vitiligo in human starts with irregularly shaped melanosomes containing pigmented membrane extensions, hyperactive melanization and selective autophagocytosis of melanosomes. Both of the aspects of immunity, humoral and cellular, have been implicated to have a role in the pathogenesis of vitiligo (Boissy *et al.* 1984). In the SL chickens, vitiligo also starts with the defect in melanocytes followed by an autoimmune response, which involves humoral and cellular reactions that eliminate abnormal pigment cells (Rezaei *et al.* 2007). The neural theory is associated with segmental vitiligo while the autoimmune hypothesis is thought to be associated to generalised vitiligo (Shajil *et al.* 2006).

Tissue destruction by the autoimmune cells in humans and animals is associated with lack of regulatory function within the immune system, hyperactive immune response and altered immune response to endogenous and exogenous factors (Erf 2010).

It has been shown that the presence of retroviral genes is linked to the expression of vitiligo in the SL chicken model and this is because of the structural and sequence similarity to exogenous viruses linked to immune dysregulation and their tissue specific expression (Sreekumar *et al.* 2000).

Increased T cells in the feather pulp of the SL chickens have been noted together with the inflammatory leukocytes before and during the onset of vitiligo (Erf 2010). In all (100%)

of the SL chickens, antibodies against melanocytes have been detected in serum (Rezaei *et al.* 2007). Melanocyte death occurs by apoptosis and is induced by cytotoxic T cells (Erf *et al.* 2001 and Wang *et al.* 2004). In mammalian melanocytes the target antigen for the antibodies is the tyrosinase related protein (TRP-1). Anti-pigment cell antibodies, which recognize the same melanocyte antigens as in humans have been identified in horses, cats and dogs.

Treatment

Treatment and re-pigmentation of vitiligo in patients can be achieved with immunosuppressive agents such as ultraviolet radiation in combination with Psolaren (PUVA), corticosteroids and cytotoxic drugs.

THE CHICKEN GENOME

The chicken genome sequence has enhanced the usefulness of chicken as an animal model (Edwards *et al.* 2005). Analyses have shown that chickens and humans share about 60 percent of their genes whereas 88 percent is shared between humans and rodents. This has made orthology relationships between chicken and humans more complex and non detectable because some proteins and domain families are under-represented in the chicken as compared with mammals. The chicken genome, however, provide new insights on evolution, structure and gene content of mammalian genomes (International Chicken Genome Sequencing Consortium 2004).

The chicken genome is 1.05 Gbp (International Chicken Genome Sequencing Consortium 2004) divided into 39 pairs of chromosomes (nine macrochromosomes and 30 microchromosomes). Macrochromosomes measure from 50-200 Mb in size while microchromosomes have an average of 12 Mb in size (Axelsson *et al.* 2005) Microchromosomes have higher G+C content, more CpG islands as well as high gene density, shorter introns, few repetitive sequences and high recombination rates. The female is the heterogametic sex carrying Z and W chromosomes while males are homogametic ZZ (Burt 2007).

The chicken genome is estimated to have 20 - 23 000 genes (<http://genome.wellcome.ac.uk>) and approximately 30-40% of the mammalian genome in size. The difference in size is attributed to reduction of interspersed repeats, segmental duplications and pseudogenes of which the chicken has 10% as compared to 50% in humans (International Chicken Genome Sequencing Consortium 2004, Burt 2007). Only 44% of the conserved sequences represent protein coding genes while conserved non-coding sequences are located far from well-defined genes and may have important regulatory functions (International Chicken Genome Sequencing Consortium 2004).

2.8 million SNPs have been reported in the chicken genome database (International Chicken Polymorphism Map Consortium 2004; International Chicken Genome Sequencing Consortium 2004; Wong *et al.* 2004). Aligned segments of the chicken and human genomes occur in long syntenic blocks with low rates of translocations and more frequent intra-chromosomal rearrangement from the last common ancestor (International Chicken Genome Sequencing Consortium 2004).

MAPPING STRATEGIES

Quantitative Trait Loci (QTL) mapping is described as the application of techniques to identify the genes controlling a certain trait. Identification of QTLs is critical for understanding the basis of complex traits (Broman 2001). Mapping complex traits has more challenges since it is not possible to follow all genomic regions underlying complex traits without knowledge of how they segregate (Doerge 2002). Even if statistical associations between QTL and markers have been found, genomic regions are sometimes large and expensive to undertake and there are also some genetic interactions such as epistasis and non-genetic factors to take into account (Doerge 2002, Zou 2009).

According to Broman (2001) experimental populations need to be fixed for relevant traits and this is achieved by repeated mating of siblings. A series of crosses is necessary of which backcross design is the simplest. Usually, an F₂ generation is used and the most favourable situation in an F₂ design is when all genes underlying the trait of interest are diallelic and alternative alleles fixed in both parental lines (Haley and Knott 1994; Perez and Varona 2000; Andesson 2001). QTL experiments can be designed in a way that enable analysis of complex phenotypes be achievable and this requires large sample of individuals that represent the whole population and provide a number of observable recombinations (Doerge 2001).

Two approaches have been assumed in QTL mapping studies and they are the infinitesimal, which assumes that traits are determined by the unlimited number of genes, each with a small effect, and the finite model which presumes a restricted or finite number of genes with a major effect (Hayes 2008). Two common methodologies for QTL mapping include the candidate gene approach (assumes that a gene could harbour a mutation that results in genetic or physiological variation of a trait) and linkage mapping which uses recombination and DNA markers to construct a linkage map (Hayes 2008).

Linkage mapping

Linkage mapping is determining the order of genes or markers on a chromosome and the relative distance between those genes based on the relative recombination frequencies. Genetic maps are essential tools for mapping and dissection of complex quantitative trait loci and allow for systematic genome scan for genes affecting all kinds of traits (Miles *et al.* 2009).

Linkage mapping is very essential in the identification of candidate genes that may not have been studied in detail as it allows defining of chromosomal regions that harbour the genes affecting the phenotype (Wick *et al.* 2006). Wick *et al.* (2006) described linkage mapping as one way of excluding candidate genes since it discriminates genes not in linkage with the trait locus, which is better than by functional studies. Linkage analysis looks for regions in the genome with a higher proportion of shared alleles among affected individuals, which depicts that there is a disease predisposing allele (Carlson *et al.* 2004). However, linkage has not worked well when trying to identify the genes underlying complex diseases (Reich 2006).

PRELIMINARY RESULTS

In a previous study (Ek *et al.* unpublished), a backcross generated after crossing pure line UCD 200 birds with F₁ birds from a cross between UCD200 and red jungle fowl was used to identify the genes underlying comb necrosis in the UCD 200 line. A whole genome scan including 384 SNPs uncovered 4 genomic regions associated with comb necrosis on chromosome 2, 12, 14 and 19 (Fig. 6).




Fig. 6. QTL regions identified in the backcross generation after matings between F₁ and pure line UCD 200. Chromosomes 2 and 14 were highly significant whereas 12 and 19 were suggestive. QTLs on chromosome 2 and 12 were identified for comb necrosis at 21 days of age while chromosomes 2, 14 and a suggestive QTL on 19 were identified for comb necrosis at 175 days of age.

In the autoimmune vitiligo animal model a Hidden Markov algorithm was used to estimate the QTL genotypes since it was more efficient than gridQTL in handling markers with mixed information content (Ek *et al.* unpublished). The whole pedigree was genotyped with 384 SNP markers. Genome wide QTL scan of individual QTL effects using Haley-Knott regression and Flexible Intercross Analysis (FIA) revealed marker association on chromosomes 3, 9 and 21 with the vitiligo phenotype (Fig. 7).




Figure 7. QTL regions implicated in autoimmune vitiligo in the SL chicken. Chromosome 3 was highly significant ($P < 0.0001$) in both Haley-Knott regression and FIA whilst chromosome 9 was significant

in FIA and suggestive using Haley-Knott regression and chromosome 21 was suggestive ($P < 0.2$) in FIA.

The purpose of this project is to

- Confirm the QTLs for autoimmune vitiligo in the F₃, F₄ and F₅ intercross generations in a cross between SL and BL.
- Validate QTL association with comb necrosis in the 2nd backcross generation in a pedigree based on a cross between UCD 200 and red jungle fowl.
- Identify SNPs in the candidate genes, *SOCS1*, *CCR8* and *CCLI* underlying the already identified QTLs for comb necrosis on chromosome 2, 12, 14 and 19 through sequencing of the UCD 200 and red jungle fowl parents from the cross.

MATERIALS AND METHODS

Animals

UCD 200 pedigree

Eight red jungle fowls (four males and four females) and eight UCD 200 individuals (four males and four females) were mated to produce an F₁ generation. 12 UCD 200 (four males and eight females) were crossed with F₁ generation birds to generate a large backcross pedigree of 598 individuals. BC₁ males were bred to UCD 200 females to produce 38 BC₂ individuals hatched in three batches.

Smyth line pedigree

14 Smyth line (SL) and 12 Brown line (BL) parental chickens were crossed to produce 57 F₁ individuals. F₁ were intercrossed to generate 496 F₂ individuals hatched in seven batches. F₂ was subsequently intercrossed to produce 45 F₃ individuals which were intercrossed to produce 102 F₄ individuals hatched in three batches and F₄ were further intercrossed to generate 90 F₅ individuals hatched in three batches.

Phenotyping

All individuals in the BC₂ generation were visually observed and scored as affected, indicating disease, or non-affected for comb necrosis at adult age (more than 175 days of age). Individuals with comb necrosis have red, swollen and inflamed combs with part of the comb missing. Individuals scored as non-affected has no inflammation and a normally formed comb (Fig. 1).

The vitiligo phenotype were recorded as binary traits, affected or non-affected. Affected individuals have lost pigmentation resulting in whole or partly white feathers. Non-affected individuals have a fully pigmented plumage. All individuals in the F₃ to F₅ generations were visually observed and scored according to the above mentioned criteria (Fig. 5).

Blood sampling and DNA extraction

A total of about 2 ml blood was collected from the brachial vein for serum, plasma and whole blood with 5.5% EDTA solution as anticoagulant. Samples were collected from 19 individuals from the 2nd backcross (F1 x UCD 200) chickens at adult age and from 90 individuals from F₅ intercross generation from the SL x BL cross at 9 weeks of age.

DNA was extracted by the salt precipitation method using ethanol described by Miller *et al.* (1988) and dissolved in 100 µl 1 x TE solution (10 mM Tris pH 8.0, 0.1 mM EDTA). 10 µl of blood was used for DNA extraction as all blood cells are nucleated in birds. The amount and quality of DNA was measured using the Nanodrop ND 1000 Spectrophotometer (Saveen Werner AB, Malmö, Sweden) and the DNA was diluted to a concentration of 50 ng/µl.

Sequencing for detection SNPs in potential candidate genes

Primers were designed to amplify all exons and introns of the *CCR8*, *CCLI* and *SOCS1* candidate genes. Reference sequences were retrieved from the UCSC genome browser and the chicken reference genome sequence WUGSC 2.1/galGal3, 2006 (<http://genome.ucsc.edu>)

Gene Fisher software (Giegerich *et al.* 1996) version 2.0 was used for primer design. The sizes of the fragments amplified in the PCR are 781 bp for *SOCS1*, 1827 bp for *CCLI* and 3350 bp for *CCR8*. Internal primers were designed for *CCR8* and *CCLI* regions as sequencing from the ends does not cover the entire fragment. All primers can be found in the Appendix (Table 8).

Amplification of the genes using parental DNA as template was carried out in Applied Biosystems 2720 thermal cyclers (Applied Biosystems, Foster City, CA, USA). Hot StarTaq polymerase (Qiagen, Hilden, Germany) with a touchdown PCR temperature cycling was used to allow optimal amplification. The total volume of the PCR reaction was 10 µl including 1 x PCR buffer (Qiagen), 2.5 mM MgCl₂, 200 µM of dNTPs, 20 pmol of forward and reverse primer, 0.1 units of Hot StarTaq polymerase (Qiagen) and about 50 ng of DNA. The thermocycling was as carried out as follows, initial activation of the polymerase at 94 C for 10 min, 14 cycles of denaturing at 94 C for 30 sec, annealing temperature starting at 65°C decreasing 1 degree per cycle to 52 C for 30 sec, extension at 72 C for 30 sec, followed by 26 cycles of denaturing at 94 C for 30 sec, annealing temperature of 52 C for 30 sec, extension at 72 C for 30 sec and a final extension at 72 C for 5 min.

DNA fragments including *CCLI* and *SOCS1* genes were separated in a 2% agarose gel and fragments including *CCR8* was separated in 1% agarose gel in 0.5 x TBE and visualised under ultraviolet light after EtBr staining.

The QIAQuick Gel extraction Kit (Qiagen) was used for extraction of DNA fragments containing the amplified genes from the agarose gel. Elution was done with 40 µl of elution buffer (Qiagen) for fragments containing *CCLI* and *SOCS1* and with H₂O for fragments containing *CCR8*. Sequencing of the amplified products was done at the Rudbeck laboratories in Uppsala.

CodonCode Aligner software, Version 3.5.6 (CodonCode Corporation, Dedham, MA, USA) was used to edit, align and compare the DNA sequences.

Genotyping using pyrosequencing

The DNA fragments used as template for pyrosequencing were amplified using three primers. One primer was designed to anneal a few bp 5' of the SNP position and the other a bit further away containing an added tail to fit with the tail primer carrying the biotin label. The resulting amplified fragment contains biotin that is used to fish out the fragment prior to pyrosequencing using the PyroMark Q96 machine (Qiagen) (See Appendix Fig. 8)

The SNPs selected for genotyping were previously identified to be informative in the cross and genotyped in all individuals in the pedigree.

PCR for amplification of SNPs genotyped in the UCD x JF BC₂ generation

Amplification of the SNPs that were genotyped was carried out with HotStarTaq polymerase protocol (Qiagen). The PCR reaction was carried out in a total volume of 25 µl containing of 1 x PCR buffer (Qiagen), 2.5 mM MgCl₂, 20 pmol dNTPs, 20 pmol of each primer except the tailed primer that had a concentration of 2 pmol, 0.1 units of HotStarTaq polymerase (Qiagen) and about 50 ng of DNA. The thermocycling was carried out as follows, initial activation of the polymerase at 94 C for 10 min followed by 14 cycles of denaturing at 94 C for 30 sec, annealing temperature starting at 65°C decreasing 1 degree per cycle to 52 C for 30 sec, extension at 72 C for 30 sec, followed by 26 cycles of denaturing at 94 C for 30 sec, annealing temperature of 52 C for 30 sec, extension at 72 C for 30 sec and a final extension at 72 C for 5 min.

PCR for amplification of SNPs genotyped in the SL x BL F₃-F₅ generations

SNPs were amplified using Ampli-TaqGold polymerase (Applied Biosystems). PCR reaction was prepared to 25 µl including the reagents below; 1 x PCR buffer (Applied Biosystems), 2.5 mM MgCl₂, 20 pmol dNTPs, 20 pmol of each primer except the tailed primer that had a concentration of 2 pmol, 0.1 units of Ampli-TaqGold polymerase (Applied Biosystems) and about 50 ng of DNA. The thermocycling was carried out as follows, initial activation of the polymerase at 94 C for 4 minutes followed by 14 cycles of denaturing at 94 C for 30 sec, annealing temperature starting at 65°C decreasing 1 degree per cycle to 52 C for 30 sec, extension at 72 C for 30 sec, followed by 26 cycles of denaturing at 94 C for 30 sec, annealing temperature of 52 C for 30 sec, extension at 72 C for 30 sec and a final extension at 72 C for 5 minutes.

Primer sequences for all markers genotyped can be found in Appendix Table 9. Inheritance check was done manually to verify whether the alleles in the individuals are in concordance with their parental alleles.

STATISTICAL ANALYSIS

Single Marker Analysis

Genotypes of SL x BL intercrosses in the F₃-F₅ generations were analysed using generalized linear model in R software. This study used a binomial trait, that is either affected or unaffected, and therefore a binomial distribution was assumed. Generalized linear models are

usually used for the analysis of binomial distributions. The ANOVA method is used when there are more than two marker genotypes.

The following model was assumed

$$y = \text{sex} + \text{batch2} + \text{ad} + \text{d} + \text{e}$$

where ad = additive effects d = dominance and e = interaction

RESULTS

Table 2. Incidence of comb necrosis in the UCD 200 2nd back cross generation.

Sex	No. of affected		No. of unaffected		Frequency of affected
	21 days	175 days	21 days	175 days	
Males		8		0	100
Females		5		6	42
Total		13		6	68

Table 3. Incidence of vitiligo in the SL x BL intercross, F3-F5 generation.

Sex	No. of affected			No. of unaffected			Frequency of affected
	F ₃	F ₄	F ₅	F ₃	F ₄	F ₅	
Males	10	13	7	15	31	38	
Females	6	6	3	13	37	38	
Total	16	19	10	28	68	76	
	36	22	13	64	78	87	

SNP identification in candidate genes for systemic sclerosis

Comparison of sequences from JF 848, JF875, *SOCS1* and *CCLI* did not show any presence of SNPs. We could only obtain a partial sequence from *CCR8*.

Pyrosequencing

Genotype data were obtained for all seven markers genotyped. Only two individuals failed to give genotypes for marker RS13673471 for Chromosome 2. See Appendix Fig. 10, 11 and 12.

QTL confirmation in the SLxBL F3-5 generations

The analysis gave suggestive evidence (P=0.1) for association between the marker located on chromosome 3, Table 4. No evidence of association was detected for markers on chromosome 9 and 21 (Table 5 and 6).

Table 4. Analysis of variance for chromosome 3.

Coefficients				
	Estimate	std. Error	z value	Pr(> z)
Intercept	-1.5254	0.4690	3.252	0.00114 **
sexM	0.3860	0.4156	0.929	0.35307
batch21	0.7285	0.4205	1.733	0.08318
ad	0.6163	0.3656	1.686	0.09187
dom	0.2071	0.5994	-0.346	0.72969

Signif. codes	0 '***'	0.001 '**'	0.01 '*'	0.1 ' ' 1

* Z-values are used to compare samples to standard normal distributions (with $\mu = 0$ $\sigma = 1$). Error is deviation from normal values.

Table 5. Analysis of variance for chromosome 9.

Coefficients				
	Estimate	std. Error	z value	Pr(> z)
Intercept	-1.0431	0.7341	-1.421	0.1553
sexM	0.3939	0.4120	0.956	0.3390
batch21	0.7433	0.4164	1.785	0.0743
ad	0.2215	0.4930	0.449	0.6532
dom	-0.9583	1,3060	-0.734	0.4631

Signif. codes	0 '***'	0.001 '**'	0.01 '*'	0.1 ' ' 1

* Z-values are used to compare samples to standard normal distributions (with $\mu = 0$ $\sigma = 1$). Error is deviation from normal values.

Table 6. Analysis of variance for chromosome 21.

Coefficients				
	Estimate	std. Error	z value	Pr(> z)
Intercept	-1.0463	1.7525	-0.597	0.5505
sexM	0.4076	0.4095	0.995	0.3196
batch21	0.7250	0.4129	1.756	0.0791
ad	-0.5736	0.9077	-0.632	0.5274
dom	-0.9086	3.3648	-0.270	0.7871

Signif. codes	0 '***'	0.001 '**'	0.01 '*'	0.1 ' ' 1

* Z-values are used to compare samples to standard normal distributions (with $\mu = 0$ $\sigma = 1$). Error is deviation from normal values.

DISCUSSION

A backcross pedigree structure (BC) was created by crossing the F₁ males from JF x UCD 200 cross to UCD 200 to increase the incidence of the disease from 4% observed in the F₂ to 50% in a backcross pedigree structure. Expression of comb necrosis indicating systemic sclerosis is alleged to be non-specific to any sex (Gershwin *et al.* 1981). This study shows that 100% of males in the 2nd backcross generation showed comb necrosis at day 200 while females showed an incidence of 42% making an overall liability of the 2nd backcross individuals to 68%. This frequency is higher in comparison to the frequency reported by Ek *et al.* (unpublished) where 95% of the males and 27% of the females in the first backcross (BC) generation were found to have comb necrosis and a study by Gershwin *et al.* (1981) who reported 95% of comb necrosis in all chickens. One possible explanation for this is that the BC₂ generation used in the study was small and perhaps not representative and phenotyping of the chickens may have been carried out at different ages. Another possible reason could be that the BC₂ chickens are old and the ones that are still alive are less sick. This may also explain the difference in frequency between the sexes, possibly females are more sensitive or maybe there are about unequal number of males and females still alive.

SSc in chickens can also appear in several forms ranging from limited skin involvement to severe or diffuse cutaneous (LeRoy *et al.* 1988). In most cases, fibrosis and thickening of the skin is observed while in some cases involvement can be confined to the digits (Fritzler and Kinsella, 1980) (Fig. 2). Mortality in the UCD 200 is 20% at one month, 40% at four months and 55% at five months of age (Gershwin *et al.* 1981). We observed slightly equivalent mortality rates (49%) at six months (200 days) in our BC₂ individuals. Most of these animals died at less than one week of age precisely 1-3 days post hatch. The precise causes of these deaths are unknown since we did not carry out autopsy.

Sequencing of the *CCR8* gene only provided partial sequence (from both UCD 200 and JF). This may be explained by the possibility that the DNA from the red jungle fowl parentals might be degraded and the gene is relatively large, 3350 bp, and therefore it is difficult to amplify a fragment of that length. One possibility could be that there may be some SNPs in the primer site that may cause inefficient annealing as low amount of the fragment could be amplified. Five SNPs can be found in the *CCR8* gene in the draft chicken sequence. They are positioned at 2:42719138, 2:42719308, 2:42719563, 2:42719645, 2:42720032 and the ones at positions 2:42719308 and 2:42719563 bp are synonymous A/G ([rs15966955](#)) and C/G ([rs13671498](#)) while non-synonymous SNPs were at positions 2:42719645, G/A ([rs14171803](#)) and 2:42720032 C/T ([rs15966952](#)) and a deletion G/- at position 2:42719138 ([rs15090893](#)).

Comparing these SNPs: in a G/- deletion there is G/G in RJF genotype, for SNP C/T RJF has C/C genotype, for SNP C/G RJF has genotypes G/G while layers have A/A, in a C/G SNP broilers have genotypes G/G, RJF has C/C and the Silkie has G/G. For A/G SNP RJF has genotypes A/A while the Silkie has G/G. (See appendix, Fig. 8 for comparative details). In the synonymous variations, amino acids remain at leucine for C/T switch and histidine for G/A transition while in non-synonymous variations the amino acids in C/G switch change from glycine to alanine whilst in A/G shift leucine is changed to proline. The roles that these SNPs may have in the candidate genes are shown in Appendix Fig. 9.

In the autoimmune vitiligo model, we noted frequencies of 16% in males and 7% in females of the F₅ generation. This is lower than expected because previous analysis showed a prevalence of 40% in the F₂ generation. Nonetheless, we are still expecting more chickens to show the disease as they grow older. It is known that the disease is more prevalent in the adolescence stage up to early adulthood (Erf 2010). We realized a reduction, in incidences of the disease from generation F₃ to F₅ at 21 weeks of age however, this should be verified as the chickens grow.

F₅ generation may not have been at the right time and more individuals may get the disease at a later stage.

We used one marker per QTL region in the statistical analysis. Since the analysis program used required a haplotype, *i.e.* two markers, the genotyping marker was duplicated in the analysis. We did not observe any significant association in any of the 3 chromosomes analysed, only a suggestive association was found between the phenotype and the marker on chromosome 3. These results were not surprising because in the previous generation analyzed, the QTL on chromosome 3 was significant in both Haley-Knott and FIA ($P < 0.0001$) whereas the QTL on chromosome 9 was only significant in FIA ($P < 0.05$) and the one on chromosome 21 was suggestive ($P < 0.2$) (See Fig. 6). (Ek *et al.* unpublished). Simulations and analyses of experimental data showed that the power of FIA was equal or better than Haley-Knott regression and insensitive to allele fixation in the founders. This method provides good estimates of QTL as well (Rönnegård *et al.* 2008).

Genotyping families with additional markers may increase the information content hence improving the likelihood of detecting true susceptibility loci. Results from simulation studies suggested that breeding populations in QTL mapping should be more or equal to 100 individuals (Darvasi and Soller 1995). Our study populations for F₃ and F₄ are lower than the suggested size except for F₅, and this may lower the power of mapping. Rönnegård *et al.* (2008) showed that using few parental lines to generate a large F₂ population increases the power of QTL detection while using a large parental population increase the chance of having more alleles with different effects if they are not fixed.

FUTURE PROSPECTS

A large population of UCD x JF BC₂ individuals should be created to enable and increase the power of the analysis.

In the SLxBL F₅ generation several more markers should be genotyped under the QTLs and males recombinant for different regions should be selected and crossed with pure line SL females to generate a back-cross to narrow down the QTL region. Darvasi and Soller (1995) showed that advanced intercross lines provide a three-five fold reduction in the confidence interval of QTL map location after eight generations *i.e.* in F₁₀ as compared to an F₂ or a backcross.

When additional potential candidate genes have been identified, their functions can be studied using the tissues that have been collected and are currently stored in freezers. Expression analysis could be carried out after extraction of RNA from the collected tissues (See list of collected tissues in Appendix Table 7).

CONCLUSION

Generally, causes of autoimmune diseases are unknown and the search for ultimate causes requires animal models. Such animal models are important in understanding the progression of diseases in humans since it is possible to set up specific crosses in animals which is not possible in humans. Several animal models have been proposed based on whether the disease is induced or inherited. Completion of the chicken genome sequence has enhanced the chicken as an animal model for understanding human diseases. Despite these advantages, animal models are however not absolute solutions to elucidating human diseases as animals and humans have physiological differences as different species and have different protein functions. Furthermore, there are differences in the immune systems between mammals and birds since birds have the bursa. In mammals tolerance is induced in the thymus (for T cells) and bone marrow (for B cells) while in birds maturation and tolerance are induced in the bursa. However, identified genes may give a hint to the pathways of their involvement in disease progression.

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude to Dr. Susanne Kerje, Department of Medical Biochemistry and Microbiology, Uppsala University, for her supervision and guidance throughout the work. She has enlightened my scope and understanding in both laboratory work and writing. I also show my gratitude to Ulla Gustafson (Department of Animal Breeding and Genetics, SLU) for her technical support in the laboratory work. I sincerely thank Weronica Ek for assisting with statistical analysis. I am also thankful to Anna-Stina Sahlqvist for working with us in this project.

I am grateful to Dr. Birgitta Malmfors and Sofia Folestam, Department of Animal Breeding and Genetics, for their support with the logistics. My appreciation also goes to my examiner professor Göran Andersson (Department of Animal Breeding and Genetics, SLU). I thank him for accepting to work with me. Finally my deepest gratitude goes to European Master in Animal Breeding and Genetics (EMABG) for the financial support.

APPENDIX




Fig. 8 pyrosequencing assay for genotyping of SSc and vitiligo individuals.




Fig. 9 Comparison of the CCR8 SNPs in different chicken strains




Fig.10. SNP variations in the CCR8 and proteins families coded for.




Fig.11 Pyrogram of a homozygous GG SNP genotype




Fig.12 Pyrogram of a heterozygous AG SNP genotype




Fig.12 Pyrogram of a homozygous AA SNP genotype




Fig.13 Expected or theoretical genotype of the AA individual




Fig.14 Expected or theoretical genotype of the heterozygous AG individual




Fig.15 Expected or theoretical genotype of the homozygous GG individual.

Table 7. Tissue samples collected form SSc and vitiligo chickens at slaughter.

Disorder		
Tissue	SSc	Vitiligo
Subcutaneous fat	√	√
Spleen	√	√
Testis	√	√
Breast muscle	√	√
Liver	√	√
Ovary	√	√
Heart muscle(apex)	√	√
Skin	√	√
Comb	√	X
Feather	√	√
Thyroid	√	√
Eyes	√	X

Table 8. Primers used for sequencing of the candidate genes in SSc.

Gene	Primer	Primer sequence
CCR8	CCR8_rev	TACCCATGACTCTTACCTCGTA
	CCR8_int1443_rev	GCATACACAGAGTGCACA
	CCR8_int2074_rev	CAAAAGAGGACTGTGAGGA
	CCR8_fwd	ATAACATCTGCTACGGAGAGGA
	CCR8_int2020_fwd	CACAGTGCCTTCAACAGA
CCL1	CCL1_fwd	GTGGTTTGTTCGGTGGCAGCA
	CCL1_int1066_rev	CTGCTGGAAGATTGGGAA
	CCL1_int995_fwd	CACAGTACGCCTCTCTCA
	CCL1_rev	TGCATTGACCCTGCATTGAATCCA
SOCS1	SOCS1_fwd	CCGTAGCCACTGACGCTATGGA
	SOCS1_rev	TAAAAACCCAGGAAGCTGGA

Table 9. Primers used in genotyping of SNPs through pyrosequencing

Chrom.	Forwad primer	Reverse primer
3	CACGACGTTGTAAAACGACA ACTTG CATCAA TGCCAGTGA	ATGAGTCTCTCTTGT CRATCA CACA
9	CACGACGTTGTAAAACGACTTAGCA GCA TTGTTCTAGTTCCG	CAA AAAACCCACATCTAGTTTC
21	CACGACGTTGTAAAACGACAGGCTG GGAGTTGGTGCACAG	GGGCTCAGGAAGCACGCAC
2	CACGACGTTGTAAAACGACCACTGA GTTTGT TGAATATGGCT	CATGTC ACTATAACTTCACCT GC
12	CACGACGTTGTAAAACGACTGCTGT GACTGG GTTGGAGG	CGCTGCCTGAGCAGAGCTGT
14	AACAGCAAGAGTAATATAAGAT	CACGACGTTGTAAAACGACTG ACTT AACACCCCATTTCTCTA
19	TCACACATGCAAACCAGACAC	CACGACGTTGTAAAACGACG GT ACAGCGCACATGGAAAAG

SNPs in QTL used in pyrosequencing

Chromosome 2 marker RS13673471

CAGATGCTAGTCAGGCTCACTGAGTTTGTGGAATATGGCTTGCATACAGTGATGG
TTACTTAAACACCTGCTCCTCCATTGCTTCCTGTGAATATTCAGT[C/T]GCAGGTGA
AGTTATAGTGACATGCCAGGAGAAGGTAGCAGCTTAAACCAGTACAGTGTTTCA
ACATTTATTGATTTATAAGAAATCACAGAAGTGCACAT

Chromosome 12 marker RS14975805

GAAATGCTGTGACTGGGTTGGAGGCTCCCAGCAGCAGGCAGGAGGTCTGACACA
GCGGGGATGAACAACTTCCGCTGGTACAAAAGGGATGACAGAACA[T/C]ACAG
CTCTGCTCAGGCAGCGGTGCCCATCAGAACCCCTTGGGTyCATCCAGGCCCTTA
GCAAAGCCACTCGGCAGCCCTGAGCCyGGCGCACGGCCCCA

Chromosome 14 marker RBL1140

TGAAGGGTTCAAGAGTTAACAGTCAAAAATTGGCAGTGATCATTTTAATTTTTGA
TTTCTGACCAGCAGTATGCTGCAACAGCAAGAGTAATATAAGATC[A/G]TTGCTA
GGCAAATCGTGGCCCAAAGAGCAGAGAAGGTAGAGAAATGGGGTGTAAAGTCA
CAGAGAACGATAACATCCAGGGCATACTCTGAAGAAGGTG

Chromosome 19 marker RS14123499

TGTGCCGTGACCTTAGCAAAAAGTGGTTCGTCTCGGGAGCGCATGGAAATTCTCCC
ATTCTAATCAAGAGGAATTAACCTTCACACATGCAAACCAGACAC[G/A]CCTCTT
CTGAAAACCTCAGTGCTAACAACGTGCGGTACATTAACCTTGATTAATCCTAACGT
AATCTTTTCCATGTGCGCTGTACCGATGCCTTCTCCACC

R script

```
phen=as.matrix(read.table("phen.out"))
batch=as.matrix(read.table("batch_sex.out"))

prob=as.matrix(read.table("KH_chr21.out",skip=2))

prob=subset(prob,phen[,2]!='-999')
batch=subset(batch,phen[,2]!='-999')
phen=subset(phen,phen[,2]!='-999')

prob=prob[594:738,]
batch=batch[594:738,]
phen=phen[594:738,]

y=as.numeric(phen[,2])
sex=as.factor(batch[,2])
batch=as.matrix(batch[,3:6])
```

```
write(t(batch),ncol=4,file='temp_batch.txt')
batch=as.matrix(read.table("temp_batch.txt"))
ad=prob[,2]-prob[,3]
dom=prob[,4]+prob[,5]

lm1=lm(y~sex+batch+ad+dom)

glm1=glm(y~sex+batch+ad+dom,family=binomial)
```

REFERENCES

- Alkhateeb A., Fain P.R., Thody A., Bennet D.C. and Spritz R.A (2003) Epidemiology of vitiligo and associated autoimmune disease in Caucasian probands and their families. *Pigment Cell Research*. Volume 16, Issue 3: pp 208- 214.
- Alexander W.S and Hilton D.J. (2004) The role of suppressors of cytokine signaling (socs) proteins in regulation of the immune response. *Annu. Rev. Immunol.* 22: 503 –529
- American Association for Clinical Chemistry (2010) Autoimmune Disorders. www.aacc.org/
- American Osteopathic College of Dermatology (2010) Dermatologic disease database. <http://www.aocd.org/>
- Andersson L (2001) Genetic dissection of phenotypic diversity in farm animals. *Nature Reviews Genetics* 2: 1-11.
- Axelsson E, Webster M.T, Smith N.G.C, Burt D.W and Ellegren H. (2005) Comparison of the chicken and turkey genomes reveals a higher rate of nucleotide divergence on microchromosomes than macrochromosomes. *Genome Research* 15: 120- 125.
- Baggiolini M. (1998) Chemokines and Leukocyte traffic. *Nature* 392: 565-568
- Broman K.W. (2001) Review of statistical methods for QTL mapping in experimental crosses. *Lab Animal*. Volume 30 Issue 7: 44-52
- Brown W.R.A., Hubbard S.J., Tickle C. and Wilson S.A. (2003) The chicken as a model for Large-scale analysis of Vertebrate gene function. *Nature reviews*. Volume 4: 87
- Boissy R.E., Moelmann G.E. and Lener A.B (1987) Morphology of melanocytes in hair bulbs and eyes of Vitiligo mice. *American Journal of Pathology*. Volume 127: Issue 2: 380-388.

Boissy R.E., Lamont S.J. and Smyth R.J jr (1984) Persistence of abnormal melanocytes in Immunosuppressed chickens of the Autoimmune 'DAM' line. *Cell and Tissue Research* 235: 663-668.

Burt D.W. (2007) Emergence of the chicken as the model organism: Implications for Agriculture and Biology. *Poultry Science Association Inc* 86: 1460-1471.

Burt D. W. (2005) Chicken Genome: Current status and future opportunities. *Genome Research*. 15: 1692-1698.

Carlson CS, Eberle MA, Kruglyak L, Nickerson DA (2004) Mapping complex disease loci in whole-genome association studies. *Nature* 429: 446-452.

Chensue, S. W., Lukacs, N. W., Yang, T. Y., Shang, X., Frait, K. A., Kunkel, S. L., Kung, T., Wiekowski, M. T., Hedrick, J. A., Cook, D. N., Zingoni, A., Narula, S. K., Zlotnik, A., Barrat, F. J., O'Garra, A., Napolitano, M., and Lira, S. A (2001) Aberrant in Vivo T Helper Type 2 Cell Response and Impaired Eosinophil Recruitment in Cc Chemokine Receptor 8 Knockout Mice *Journal of Experimental. Medicine* 193:573–584.

Cerundolo R. De Caprariis D. Esposito L. Maiolino P Restucci B and Roperto F (2007) Vitiligo in two Water buffaloes: Histological, histochemical, and ultrastructural investigations. *Pigment Cell Research*. Volume 6, Issue 1: 23-28.

Clucks & Chooks (2008) *The History of the chicken*.

<http://www.henkeeping.co.uk/>

Coffman C. J., Doerge R. W., Wayne M. L and McIntyre L. M, (2003) Intersection tests for single marker QTL analysis can be more powerful than two marker QTL analysis. *BMC Genetics*. 4: 10

Crawford R.D (1996) Origin and history of poultry species, in: Crawford R.D. (Ed.), *Poultry Breeding and Genetics*, Elsevier, New York, 1990, USA pages 6792-6795.

Dairaghi D.J, Fan R.A, McMaster BE, Hanley M.R, and Schall T.J (1999) HHV8-encoded vMIP-I selectively engages chemokine receptor CCR8. Agonist and antagonist profiles of viral chemokines. *Journal of Biological Chemistry* (1999) 274: 21569-21574.

Darvarsi A and Soller M. (1995) Advanced Intercross Lines, an Experimental Population for Fine Genetic Mapping. *Genetics*. Volume 141, issue 3: 1199- 1207.

De Vries M. E, Cao H, Wang J., Xu L, Kelvin A. A., Ran L, Chau L. A, Madrenas J, Hegele R A and Kelvin D J. (2003) Genomic Organization and Evolution of the CX3CR1/CCR8 Chemokine Receptor Locus. *The Journal of Biological Chemistry*: volume 278. Number 14: 11985- 11994.

Dietrich H.M. (1989) Housing breeding and selecting chickens of the obese strain (OS) with spontaneous autoimmune thyroiditis. *Laboratory Animals*. 23: 345-352.

Doerge R.W, (2002) Mapping and Analysis of Quantitative Trait Loci in Experimental Populations. *Nature Reviews*. Volume 3: 43- 51.

Erf G. F, (2010) Animal Models. in Picardo M and Taieb (eds) *Vitiligo*. Published by Springer-Verlag Berlin Heidelberg. Berlin, Germany.

Egan P.J., Lawlor K.E., Alexander W.S. and Wicks I.P. (2003) Suppressor of Cytokine Signalling-1 regulates acute inflammatory arthritis and T-Cell activation. *The Journal of Clinical Investigation*. Volume 111 Issue 6: 915-924.

Erf G., Bersi T.K., Wang X., Sreekumar P.G and Smyth R J., jr. (2002) Herpesvirus Connection in the Expression of Autoimmune Vitiligo in Smyth Line Chickens. *Pigment Cell & Melanoma*. Volume 2, Issue 1: 40-46.

Erf G.F., Bersi T.K., Wang X, (2001) Herpesvirus connection in the expression of Autoimmune Vitiligo in the Smyth line chickens. *Pigment Cell Research*. 14: 40-46.

Fenner J.E., Starr R., Cornish A.L., Zhang J.G., Metcalf D., Schreiber R.D., Sheehan K., Hilton D.J., Alexander W.S., Hertzog P.J, (2006) Suppressor of cytokine signalling 1 regulates the immune response to infection by a unique inhibition of type 1 interferon activity. *Nature immunology*. Volume 7. Issue 1: 33-39.

Fritzler M.J, Kinsella T.D and Garbutt E, (1980) The CREST syndrome: A distinct serologic entity with anticentromere antibodies. *Volume 69, Issue 4: 520-526*.

Fumihito A, Miyake T., Sumi Shin-Ichiro, Takada M., Ohno S. and Kondo N, (1994) one subspecies of Red Jungle Fowl (*Gallus gallus*) Suffices as the matriarchic ancestor of all domestic breeds. *Proceedings of the National Academy of Sciences of the United States of America*. Volume 91, Number 26: 12505-12509.

Gershwin M.E., Abplanalp H., Castles J.J., Ikeda R.M., van Der Water J., Eklund J. and Haynes D, (1981) Characterization of spontaneous disease of White Leghorn chickens resembling Progressive Systemic Sclerosis. *J. Exp Med*. Volume 153.

Giegerich R, Meyer F and Schleiermacher C, (1996) GeneFisher - software support for the detection of postulated genes. *Proceedings of International Conference on Intelligent Systems for Molecular Biology*. Volume 4: 68-77

Haley C.S and Knott S.A (1994). Mapping quantitative trait loci in crosses between outbred lines using least squares. *Genetics*. Volume 236, issue 3: 1195-1207.

Hanada T, Yoshida H, Kato S, Tanaka K, Masutani K, Tsukada J, Nomura Y, Mimata H, Kubo M and Yoshimura A. (2003) Suppressor of Cytokine Signaling-1 Is Essential for

Suppressing Dendritic Cell Activation and Systemic Autoimmunity. *Immunity*. Volume 19, Issue 3: 437-450.

Haustein U.F. (2002) Systemic Sclerosis- Scleroderma. *Dermatology online Journal*. Volume 8 Number 1.

Havey W (1628) on the motion of the heart and blood in animals. *Modern History Sourcebook*
<http://www.fordham.edu/halsall/mod/1628harvey-blood.html>

Hedges S.B (2002) The origin and evolution of model organisms. *Nature Reviews Genetics*. Volume 3: 838-849.

Hellström A R., Sundström E, Gunnarsson U, Bed'Hom B, Tixier-Boichard M., Honaker C F, Anna-Stina Sahlqvist, Jensen P, Kämpe O, Siegel P B., Kerje S and Andersson L. (2010) Sex-linked barring in chickens is controlled by the CDKN2A/B tumor suppressor locus. *Pigment Cell Melanoma Res.* Volume 9999. Issue 9999.

Hogquist K.A, Baldwin T, Jameson S (2005). Central tolerance: learning self-control in the thymus. *Nature Reviews Immunology*. Volume 5 Issue 10: 772–82

Hook R.R jr, Berkelhammer J and Oxenhandler R.W. (1982). Melanoma. *Sinclair Swine Melanoma. American Journal of Pathology*. Volume 108 : 130-133.

International Chicken Genome Sequencing Consortium (2004). Sequence and comparative analysis of the chicken genome provide unique perspectives on vertebrate evolution. *Nature* 432:695–716.

International Chicken Polymorphism Map Consortium. 2004. A genetic variation map for chicken with 2.8 million single-nucleotide polymorphisms. *Nature* 432:717–722.

Jimenez S.A and Derk C.T (2004) Molecular Pathways toward an Understanding of the Pathogenesis of Systemic Sclerosis. *Annals of Internal Medicine*. Volume 140, Issue 1: 37-50.

Jin Y., Mailloux C.M, B.S., Gowan K., Riccardi S.L., LaBerge G., Bennett D. C., Fain P. R., and Spritz, R.A. (2007). *NALP1* in Vitiligo-Associated Multiple Autoimmune Disease. *The New England Journal of Medicine*. volume 356: number 12. 1216-1225.

John S, Shephard N, Liu G, Zeggini E, Cao M, Chen W, Vasavda N, Mills T, Button A, Hinks A, Eyre S, Jones K.W, Olie W, Silman A, Gibson N, Worthington J and Kennedy G.C. (2004) Whole-Genome Scan, in a Complex Disease, Using 11,245 Single-Nucleotide Polymorphisms: Comparison with Microsatellites. *The American Journal of Human Genetics*. Volume 75, Issue 1; 54-64.

Kakourou T. (2009). Vitiligo in children. *World Journal of Paediatrics*. Volume 5 Issue 4 pp 265- 268.

Kanginakudru S, Metta M, Jakati R.D, and Nagaraju J. (2008) Genetic evidence of Indian red jungle fowl corroborates multiple domestication of modern day chicken. *BMC Evolutionary Biology*. Volume 8:174.

Kanzler S, Lohse A.W., Keil A, Henninger J, Dienes H P., Schirmacher P, Rose-John S, Meyer Zum Büschenfelde K H, and Blessing M (1999) TGF-1 in liver fibrosis: an inducible transgenic mouse model to study liver fibrogenesis. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. Volume 276. Issue 4: 1059-1068.

Kerje S. (2003) Mapping genes affecting Phenotypic traits in Chickens. Phd dissertation, Uppsala University, Uppsala, Sweden.

Kerje S., Sharma P., Gunnarsson U., Kim H., Bagchi S., Fredriksson R., Schütz K., Jenson P., von Heijne G., Okimoto R and Andersson L (2004) The *Dominant white*, *Dun* and *Smoky* color variants in Chicken are Associated with insertion/Deletion Polymorphism in the *PMEL17* gene. *Genetics* 168, 1507-1518.

Lamont S.J. Boissy R.E. and Smyth J.R. jr (1982) Humoral immune response and expression of postnatal Amelanosis in Dam Line chickens. *Immunological investigations*. Volume 11 issue 2 pp 121- 127.

Lamoreux M.L, Boissy R.E, Womack J.E and Nordlund J.J (1992) The *vit* Gene Maps to the *mi* (Microphthalmia) Locus of the Laboratory Mouse. *The Journal of Heredity*. Volume 83, Issue 6: 435 - 439.

LeRoy E.C, Black C, Fleischmajer R, Jablonska S, Krieg T, Medsger T.A Jr, Rowell N, Wollheim F. (1988) Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *Journal of Rheumatology* Volume 15, issue 2: 202-205.

Medsger T.A Jr. (1991) Treatment of systemic sclerosis. *Annals of Rheumatic disease*. Volume 50: 877- 886.

Miles L.G., Isberg S.R., Glenn T.C., Lance S.L., Dalzell P., Thomson P.C. and Moran C (2009). A genetic linkage map for the salt water crocodile (*Crocodylus porosus*). *BMC Genomics* 10: 339

Miller M.D, Krangel MS (1992). "The human cytokine I-309 is a monocyte chemoattractant". *Proceedings of the National Academy of Sciences*. U.S.A. Volume 89 Issue 7: 2950–2954.

Miller S.A, Dykes D.D and Polesky H.F (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research*

Namazi M.R. (2005). Phenytoin as a novel anti-vitiligo weapon. *Journal of Autoimmune Diseases*. volume 2:11

Norris D.A., Kissinger R.M. Naughton G.M. and Bystryn J.C (1988). Evidence for Immunologic Mechanisms in Human Vitiligo: Patients' Sera Induce Damage to Human Melanocytes In Vitro by Complement-Mediated Damage and Antibody-Dependent Cellular Cytotoxicity. *Journal of Investigative Dermatology*. Volume 80 : 783-789.

Pasteur, L. 1880. De l'attenuation du virus cholera des poules. *Compte rend. Acad. se.* 91: 673-680 In *Milestones in Microbiology: 1556 to 1940*, translated and edited by Thomas D. Brock, ASM Press. 1998, p126

Pe´rez-Enciso M and Varona L (2000) Quantitative Trait Loci Mapping in F2 crosses between outbred Lines. *Genetics*. Volume 155, issue 1: 391-405.

Pielberg G. R, Golovko A, Sundström E, Curik I, Lennartsson J, Seltenhammer M H, Druml T, Binns M, Fitzsimmons C, Lindgren G, Sandberg K, Baumung R, Vetterlein M, Strömberg S, Grabherr M, Wade C, Lindblad-Toh K, Ponte ´n F, Carl-Henrik Heldin, Johann Sölkner & Leif Andersson (2008). A cis-acting regulatory mutation causes premature hair graying and susceptibility to melanoma in the horse. *Nature Genetics*, 40: 1004-10009.

Rezai N., Gavalas N.G. Weetman A.P. and Kemp E.H. (2007). Autoimmunity as an aetiological factor in vitiligo. *Journal of the European Dermatology and virology*. Volume 21: Issue 7 pp 865 - 876.

Rönnegård L, Besnier F and Carlborg Ö. (2008) An Improved Method For Quantitative Trait Loci And Identification Of Within Line Segregation In F2 Intercross Designs. *Genetics* 178: 2315-2326.

Roos R.S, Loetscher M, Legler DF, Clark-Lewis I, Baggiolini M, Moser B (July 1997). Identification of CCR8 the receptor for the human CC chemokine I-309. *Journal of Biological Chemistry*. Volume 272, Issue 28: 17251–17254.

Royo J. L, Hidalgo M and Ruiz A (2007) Pyrosequencing protocol using a universal biotinylated primer for mutation detection and SNP genotyping. *Nature Protocols*, volume 2: 1734- 1739.

Rula A and Hajj-Ali. (2008). Systemic sclerosis. *The Merck Manual Online Medical Library*. <http://www.merck.com/mmpe/sec04/ch032/ch032h.html#sec04-ch032-ch032h-215>

Schwartz R.A., Dziankowska-Bartkowiak B., Zalewska A and Sysa-Jedrzejowska A (2009). Systemic sclerosis, *emedicine*, Medscape. E-medicine web page.

Sgonc R. (1999). The Vascular Perspective of Systemic Sclerosis: Of Chickens, Mice and Men. *International Archives of Immunology and Allergy*. 120: 169-176

Shajil E.M, Marfatia Y.S and Begum R. (2006) Acetylcholine esterase levels in different clinical types of vitiligo in Baroda, Gujarat. *Indian Journal of Dermatology*. Volume 51, Issue

4: pp 289- 291.

Siegel P.B., Dodgson J.B. and Andersson L. (2006). Progress from Chicken Genetics to the Chicken Genome. Invited Reviews. *Poultry Science* 85: 2050-2060.

Smyth J.R. (1989). The Smyth chicken: a model for autoimmune amelanosis. *Poultry Biology* 2: 1-19.

Sprent J. (1995) Central tolerance of T cells. *Int reviews, Immunology*. Volume 13. Issue 2: 95-105.

Sreekumar G.P., Smyth J.R., Ambady S and Ponce De Leon F.A (2000). Analysis of the Effect of endogenous viral genes in Smyth Line chicken model for autoimmune vitiligo. *American Journal of Pathology*. Volume 156: Issue 3 pp 1099 - 1107.

Stern C.D. (2005). The chick: A great model system even becomes greater. *Developmental cell*. Volume 8, Issue 1: pp 9-17.

Takehara K. (2003) Pathogenesis of systemic sclerosis. *The Journal of Rheumatology*. Volume 30: 755-759.

Van der Water J, Jimenez S.A and Gershwin E.M. (1995). Animal Models of Scleroderma: Contrasts and comparisons. *Internal Reviews of Immunology*. Issue no 2-4: 201-216

Van der Water J and Gershwin E.M. (1985). Animal model of Human disease. Avian Scleroderma. An inherited Fibrotic disease of White Leghorns Chicken Resembling Progressive Systemic Sclerosis. *American Journal of Pathology*. Volume 120, Issue 3: 478-482.

West B and Zou B.X, (1988). Did Chickens go North? New Evidence for Domestication. *Journal of Archaeological Science*. Volume 15: 515-533.

Wick G., Andersson L., Hala K., Gershwin M.E., Gisela E.F., Lamont S.J. and Sgoc R (2006). Avian Models with Spontaneous Autoimmune Diseases. *Advances in Immunology*. Volume 92.

Wick G., Sundick R.S and Albin B. (1974). The Obese strain of chickens (OS): An animal Model with Spontaneous Autoimmune Thyroiditis. *Clinical Immunology and Immuno Pathology*. Volume 3 Issue 2: 272-300.

Wang X. and Erf G.F. (2004). Apoptosis in feathers of Smyth line chickens with autoimmune vitiligo. *Journal of Autoimmunity*. Volume 24, issue no 1: 21-30.

Yamamoto T, Eckes B, Krieg T. (2000) Bleomycin increases steady-state levels of type I collagen, fibronectin and decorin gene expression in human skin fibroblasts.

Archives of Dermatological Research 292: 556–561.

Yamamoto T. (2005) Characteristics of Animal Models for Scleroderma. *Current Rheumatology Reviews* 1: 101- 109.

Yamamoto T. (2009). Scleroderma - Pathophysiology. Review article. *European Journal of Dermatology* Volume 19 Issue1: 14 - 24.

Yamamoto T. (2010) Animal model of systemic sclerosis. *Journal of Dermatology* 37: 26-41.

Zhu C, Gore M, Bucker E.S and Yu J. (2009) Status and Prospects of Association Mapping in Plants. *The Plant Genome*. Volume 1, Issue 1: 5-20.

Zou F. (2009) QTL mapping in Intercross and Backcross Populations. *Cardiovascular Genomics*. *Methods in Molecular Biology*. Volume 573: 157- 173.